

WYOMING GAME AND FISH DEPARTMENT

FISH DIVISION

ADMINISTRATIVE REPORT

TITLE: Instream Flow Studies on Packstring Creek, a Bonneville Cutthroat Trout (*Oncorhynchus clarki utah*) Stream.

PROJECT: IF-PE97-07-9705

AUTHOR: Paul D. Dey and Thomas C. Annear

DATE: March 1997

ABSTRACT

Studies were conducted in 1996 on Packstring Creek to determine instream flows necessary for maintaining Bonneville cutthroat trout (BRC) habitat and populations. Studies complemented ongoing habitat and population monitoring of BRC streams (Remmick et al. 1994). A habitat retention model was used to derive a single year-round instream flow water right recommendation of 0.7 cfs

INTRODUCTION

Wyoming Bonneville cutthroat trout (*Oncorhynchus clarki utah*) populations occur primarily in the Thomas Fork and Smiths Fork watersheds. These major Bear River tributaries and associated waterways were surveyed for physical, chemical, and biological characteristics between 1966 and 1977 (Miller 1977). Binns (1981) reviewed the distribution, genetic purity, and habitat conditions for Bonneville cutthroat trout populations. Recent population and habitat survey results are in Remmick (1981, 1987) and Remmick et al. (1994).

In general, populations are limited by seasonally low flows, lack of riparian cover, thermal pollution arising in conjunction with low flows and reduced riparian vegetation, and silt pollution (Binns 1981). Livestock grazing on both private and public lands is an important contributor to degraded habitat conditions. However, despite habitat challenges, significant populations remain throughout the native range and a recent review categorized Wyoming's populations as "95% secure, stable" (Duff 1996).

Population status in other Bonneville basin states including Utah, Idaho and Nevada is less secure. Therefore, Bonneville cutthroat trout were recently petitioned for listing under the Endangered Species Act but are not listed at this time. Status review was initiated in response to concerns expressed by the Idaho Fish and Game Department, the Desert Fishes Council and the Utah Wilderness Association. This species is considered "rare" in Wyoming by the Wyoming Game and Fish Department (WGFD 1977).

A 5-year management plan for Wyoming, developed by the Wyoming Game and Fish Department (WGFD) in coordination with the U.S. Forest Service (USFS) and U.S. Bureau of Land Management (BLM), outlines management goals and provides criteria for listing Bonneville cutthroat trout as threatened (Remmick et al. 1994). The plan's purpose is to outline management practices to prevent listing by moving toward wider distribution and higher populations. The plan recommends that status decisions be made after five-years of population and habitat monitoring. Habitat protection by acquiring instream flow water rights will help prevent additional population declines.

Resource management practices could be significantly affected by listing Bonneville cutthroat trout as Threatened or Endangered. Instream flow water right identification and acquisition on Bonneville cutthroat trout streams is important to help avoid listing. Therefore, since 1993 the WGFD has filed for instream flow water rights on most non-ephemeral tributaries with documented BRC presence in the Thomas and Smiths Fork drainages. Studies in 1996 focused on Poker Hollow Creek, North Fork Smiths Fork Creek, Lander Creek, Trespass Creek, Little White Creek, and Packstring Creek.

Study objectives were to 1) investigate the relationship between discharge and physical habitat quantity and quality for Bonneville cutthroat trout and, 2) determine an instream flow regime necessary to maintain or improve Bonneville cutthroat trout populations.

METHODS

Study Area

Packstring Creek is a small headwater tributary that flows generally southeast before entering Salt Creek near Salt Flat (Figure 1). The Packstring Creek basin is under Bridger-Teton National Forest administration. Grass, sedge and scattered willow occur in a narrow riparian zone. Some aspen occur in hillside pockets and at higher elevations. Stream gradient estimated from a USGS 7.5 minute quadrangle map averages about 3.8%. Channel type was rated as A4 (Rosgen 1985) indicating a headwater stream with relatively high gradient but occurring in a geologic deposition area so that channel substrate is predominately small cobble, gravel and sand.

Fisheries

The term "fishery" refers to fish populations and their associated habitat in a defined area. Instream flow water rights are intended to maintain or improve habitat conditions so that viable fisheries are maintained or improved. Trout populations, particularly in small mountain streams, normally fluctuate widely. In a western Oregon stream studied for 11 years, density of age 0 cutthroat trout (fry, <2 inches) varied from 8 to 38 per 100 m² and density of age 1 cutthroat trout (juveniles, 4-4.5 inches) ranged from 16 to 34 per 100 m² (House 1995). In this example, population fluctuations occurred despite the fact that habitat conditions were not degraded and appeared to be relatively stable. The author suggested that small changes in peak winter flows between years would have accounted for shifts in overwinter survival between age-classes.

Similar population fluctuations occur in Sand Creek, a Crook County, Wyoming stream that experiences relatively little discharge variation (Mueller 1987). Sand Creek brown trout population density ranged from 646 trout/mile to 4,060 trout/mile in a three year period. Biomass estimates for the same period ranged between 48 and 142 pounds per acre.

Long-term trout population maintenance in small streams depends on periodic strong year classes produced in good flow years. Without benefit of periodic favorable flows, populations in some streams would decline or disappear. The WGFD instream flow strategy recognizes the inherent variability of trout populations and thus defines the "existing fishery" as a dynamic feature. Instream flow recommendations are based on a goal of maintaining habitat conditions that provide the opportunity for trout numbers to fluctuate within existing natural levels.

During survey work in 1995, Forest Service biologists collected Bonneville cutthroat trout ranging between 4.5 and 10.5 inches from the mainstem and small tributary in T29N, R119W, S22. On August 27, 1996 approximately 262 feet of stream were electrofished to yield 13 BRC ranging in length from 2.1 to 6.5 inches. The resulting population estimate was 288 trout per mile (27.1 lbs/acre).

Habitat Modeling

After visually surveying approximately 0.5 stream miles, a study area was located at T29N, R119W, S27, SE1/4 (Figure 1 and 2). Three perpendicular transects across hydraulic control riffles were used to determine a flow level that maintains certain hydraulic criteria (see below). Dates and flow levels during which data were collected are listed in Table 1.

Instream flow filing recommendations derived from this site were applied to a 1.3 mile-long reach extending downstream from the confluence of two forks at the boundary between sections 22 and 27 of T29N, R119W to the confluence with Salt Creek at T29N, R119W, S26. The land through which the proposed segment passes is under Bridger-Teton National Forest administration.

The small headwater nature of Packstring Creek implies that deep water adult habitat is not abundant. Instead, small areas are suitable for moderate sized fish under normal flow conditions. It is likely that small tributaries in the Thomas and Smiths Fork drainages act as recruitment sources to the larger creeks. Spawning and nursery habitat in these tributaries appear plentiful during years with normal flow. Fry may passively drift down into the larger streams shortly after emergence or adults may emigrate as small pools no longer meet their energetic or predator avoidance needs. From a drainage-wide perspective, maintaining BRC populations should hinge on maintaining hydraulic connectivity between the small headwater tributaries and the larger streams. Therefore, our general approach with the small headwater tributaries was to simply determine a flow level that maintains habitat levels and retains flowing water through riffle areas. Such flow levels should ensure that normal processes of immigration and emigration can occur. Additional studies and supplemental instream flow filings may be considered in the future if fisheries data and management goals indicate such actions are needed and appropriate.



Figure 2. Habitat Retention transect 2.

Table 1. Dates and discharges instream flow data were collected from Packstring Creek in 1996.

Date	Discharge (cfs)	Data
August 27	0.3	Population Estimate
September 9	0.3	Habitat Retention

Habitat Retention Method

A Habitat Retention method (Nehring 1979, Annear and Conder 1984) was used to identify a maintenance flow by analyzing data from the three hydraulic control riffle transects added on September 9. A maintenance flow is defined as the continuous flow required to maintain specific hydraulic criteria in stream riffles. Year-round criteria maintenance in riffles ensures that habitat is also maintained in other habitat types such as runs or pools (Nehring 1979). In addition, maintenance of identified flow levels may ensure passage between habitat types for all trout life stages and maintain adequate benthic invertebrate survival.

A maintenance flow is realized at the discharge for which any two of the three criteria in Table 2 are met for all riffle transects in a study area. The instream flow recommendations from the Habitat Retention method are applicable year round.

Table 2. Hydraulic criteria for determining maintenance flow with the Habitat Retention method.

Category	Criteria
Mean Depth (ft)	Top Width ^a X 0.01
Mean Velocity (ft/s)	1.00
Wetted Perimeter (%) ^b	50

a - At average daily flow or mean depth = 0.20, whichever is greater

b - Percent of bank full wetted perimeter

Simulation tools and calibration techniques used for hydraulic simulation in PHABSIM (Physical Habitat Simulation) are also used with this technique. The PHABSIM method uses empirical relationships between physical variables (depth, velocity, and substrate) and suitability for fish to derive weighted usable area (WUA; suitable ft² per 1000 ft of stream length) at various flows. In habitat retention, only hydraulic control riffles are examined and no direct reference is made to suitability of identified depths and velocities for trout life stages. Instead, the AVDEPTH model under PHABSIM is used to simulate average cross section depth, wetted perimeter and velocity for a range of flows on each cross section. The flow that maintains 2 out of 3 criteria for all three transects is then readily apparent.

RESULTS AND DISCUSSION

Habitat Retention Analysis

Habitat retention analysis indicates that 0.7 cfs is required to maintain hydraulic criteria at all riffles (Table 3). Maintenance of naturally occurring flows up to this flow is necessary at all times of the year.

Table 3. Simulated hydraulic criteria for three Packstring Creek riffles. Average daily flow = 1.4 cfs. Bank full discharge = 9.1 cfs.

	Mean Depth (ft)	Mean Velocity (ft/s)	Wetted Perimeter (ft)	Discharge (cfs)
Riffle 1	0.45	2.67	8.0	9.1
	0.25	1.07	5.1	1.3
	0.24	0.98^a	5.0	1.1
	0.22	0.93	4.9	1.0
	0.21	0.72	4.8	0.8
	0.19^a	0.70	4.8	0.7^b
	0.18	0.72	4.7	0.6
	0.17	0.67	4.5	0.5
	0.16	0.61	4.3	0.4
	0.15	0.52	4.0^a	0.3
Riffle 2	0.13	0.43	3.6	0.2
	0.45	3.09	7.0	9.1
	0.27	1.32	3.5^a	1.2
	0.27	1.26	3.4	1.1
	0.26	1.19	3.4	1.0
	0.24	1.07	3.3	0.8
	0.22	1.02^a	3.2	0.7^b
	0.22	0.92	3.1	0.6
	0.20^a	0.84	3.1	0.5
	0.19	0.75	3.0	0.4
Riffle 3	0.16	0.67	2.9	0.3
	0.14	0.54	2.7	0.2
	0.56	2.04	8.4	9.1
	>0.29	>0.81^a	>5.3	>1.2
	0.28	0.77	5.3	1.1
	0.26	0.75	5.2	1.0
	0.25	0.68	4.9	0.8
	0.25	0.64	4.6	0.7
	0.25	0.60	4.2^a	0.6^b
	0.25	0.56	3.7	0.5
0.23	0.51	3.5	0.4	
0.22	0.44	3.2	0.3	
0.19^a	0.35	3.1	0.2	

a - Hydraulic criteria met

b - Discharge at which 2 of 3 hydraulic criteria are met

Trout populations are naturally limited by extreme winter conditions (October through March; Needham et al. 1945, Reimers 1957, Butler 1979, Kurtz 1980, Cunjak

1988, Cunjak 1996). Formation of frazil ice (suspended ice crystals formed from super-chilled water) in high gradient stream reaches can be both a direct mortality source through gill abrasion and subsequent suffocation or an indirect mortality source as resultant anchor ice limits habitat, causes localized de-watering, and extracts excessive metabolic demands on fish forced to seek ice-free habitats (Brown et. al 1994). Pools downstream from high gradient frazil ice-forming areas can accumulate anchor ice when woody debris or surface ice provides anchor points for frazil crystals (Brown et. al 1994, Cunjak and Caissie 1994). Such accumulations may result in mortalities if emigration is blocked by low winter flows or ice dams.

In high gradient systems, groundwater influx areas, ice covered pools not close to frazil sources, heavy snow cover with stream bridging, and pools with little woody debris offer refugia from frazil ice (Brown et al. 1994). Lower gradient streams and narrow streams are more likely to have insulating surface ice cover or at higher elevations, heavy snow cover and bridging. Packstring Creek's high elevation, relatively narrow width and moderate slope likely means that heavy snow fall will result in snow bridging. Frazil ice formation is likely a concern mainly in early winter before sufficient insulating snow is present or in late winter when snow melt becomes superchilled by flowing over snow and ice before entering the stream. Therefore, natural winter flow levels up to the identified 0.7 cfs should be maintained to maximize access to and availability of frazil-ice-free refugia. Any artificial reduction of natural winter stream flows could increase trout mortality and effectively reduce the number of fish the stream could support.

The 0.7 cfs identified by the Habitat Retention Method may not always be present during the winter. Because the existing fishery is adapted to natural flow patterns, occasional shortfalls during the winter do not imply a need for additional storage. Instead, they illustrate the necessity of maintaining all natural winter stream flows, up to 0.7 cfs, to maintain existing trout survival rates.

INSTREAM FLOW RECOMMENDATIONS

Based on the analyses and results outlined above, the instream flow recommendations in Table 4 will maintain the existing Packstring Creek Bonneville cutthroat trout fishery. These recommendations apply to a 1.3 mile Packstring Creek segment extending downstream from the confluence of two forks at the boundary between sections 22 and 27 of T29N, R119W to the confluence with Salt Creek at T29N, R119W, S26. The land through which the proposed segment passes is under Bridger-Teton National Forest administration. Because data were collected from representative habitats and simulated over a wide flow range, additional data collection under different flow conditions would not significantly change these recommendations.

Table 4. Instream flow recommendations to maintain the existing Packstring Creek trout fishery.

Time Period	Instream Flow Recommendation (cfs)
October 1 to September 30	0.7

This analysis does not consider periodic requirements for channel maintenance flows. Because this stream is unregulated, channel maintenance flow needs are adequately met by natural runoff patterns. If regulated in the future, additional studies and recommendations may be appropriate for establishing channel maintenance flow requirements.

LITERATURE CITED

- Annear, T.C. and A.L. Conder. 1984. Relative bias of several fisheries instream flow methods. *North American Journal of Fisheries Management* 4:531-539.
- Binns, N.A. 1981. Bonneville cutthroat trout *Salmo clarki* utah in Wyoming. Wyoming Game and Fish Department, Fisheries Technical Bulletin No. 5.
- Brown, R.S., S.S. Stanislowski, and W.C. Mackay. 1994. Effects of frazil ice on fish. In *Proceedings of the Workshop on the Environmental Aspects of River Ice, Saskatoon, Sask., 18-20 August 1993. Edited by T.D. Prowse. NHRI Symp. Ser. No.12. National Hydrology Research Institute, Saskatoon, Sask. pp.261-278.*
- Butler, R. 1979. Anchor ice, its formation and effects on aquatic life. *Science in Agriculture, Vol XXVI, Number 2, Winter, 1979.*
- Cunjak, R.A. 1988. Physiological consequences of overwintering in streams; the cost of acclimatization? *Canadian Journal of Fisheries and Aquatic Sciences* 45:443-452.
- Cunjak, R.A. 1996. Winter habitat of selected stream fishes and potential impacts from land-use activity. *Canadian Journal of fisheries and Aquatic Sciences* 53(Suppl.1):267-282.
- Cunjak, R.A. and D. Caissie. 1994. Frazil ice accumulation in a large salmon pool in the Miramichi River, New Brunswick: ecological implications for overwintering fishes. In *Proceedings of the Workshop on the Environmental Aspects of River Ice, Saskatoon, Sask., 18-20 August 1993. Edited by T.D. Prowse. NHRI Symp. Ser. No.12. National Hydrology Research Institute, Saskatoon, Sask. pp.261-278.*
- Duff, Donald. 1996. Conservation assessment for inland cutthroat trout. U.S. Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah. Pages 35-73.
- House, R. 1995. Temporal variation in abundance of an isolated population of cutthroat trout in western Oregon, 1981-1991. *North American Journal of Fisheries Management* 15:33-41.
- Kurtz, J. 1980. Fishery management investigations. - a study of the upper Green River fishery, Sublette County, Wyoming (1975-1979). Completion Report. Wyoming Game and Fish Department, Fish Division, Cheyenne.
- Miller, D.D. 1977. Comprehensive survey of the Bear River drainage. Wyoming Game and Fish, Administrative Report.
- Mueller, J.W. 1987. Electrofishing results in Sand Creek, Hospital Gulch Station, Sand Creek Country Club, 1986. Wyoming Game and Fish Department Administrative Report.
- Needham, P., J. Moffett, and D. Slater. 1945. Fluctuations in wild brown trout populations in Convict Creek, California. *Journal of Wildlife Management* 9:9-25.

- Nehring, R. 1979. Evaluation of instream flow methods and determination of water quantity needs for streams in the state of Colorado. Colorado Division of Wildlife, Fort Collins.
- Reimers, N. 1957. Some aspects of the relation between stream foods and trout survival. California Fish and Game 43:43-69.
- Remmick, R. 1981. A survey of native cutthroat populations and associated stream habitats in the Bridger-Teton National Forest. Wyoming Game and Fish Department, Administrative Report.
- Remmick, R. and N.A. Binns. 1987. Effect of drainage wide habitat management on Bear River cutthroat trout (*Salmo clarki utah*) populations in the Thomas Fork drainage, Wyoming. Wyoming Game and Fish Department, Administrative Report.
- Remmick, R., K. Nelson, G. Walker, and J. Henderson. 1994. Bonneville cutthroat trout inter-agency five year management plan (1993-1997).
- Rosgen, D. 1985. A stream classification system. IN: Riparian Ecosystems and Their Management; Reconciling Conflicting Uses. Proceedings of the First North American Riparian Conference, April 16-18, Tucson, Arizona. GTR-RM120, pp. 91-95.
- WGFD. 1977. Current Status and Inventory of Wildlife in Wyoming. 133 p.