

ATLAS OF BIRDS, MAMMALS, AMPHIBIANS, AND REPTILES IN WYOMING



USFWS National Digital Library



USFWS National Digital Library



USFWS National Digital Library



USFWS National Digital Library

**Wyoming Game and Fish Department
Nongame Program
Statewide Wildlife and Habitat Management Section
Wildlife Division
December 2016**

**ATLAS OF BIRDS, MAMMALS,
AMPHIBIANS, AND REPTILES IN WYOMING**

**Wyoming Game and Fish Department
Nongame Program
Statewide Wildlife and Habitat Management Section
Wildlife Division**

December 2016

**Compiled and edited by:
Andrea Orabona, Courtney Rudd, Nichole Bjornlie,
Zack Walker, Susan Patla, and Bob Oakleaf**


**Wyoming Game and Fish Department
Nongame Program
260 Buena Vista
Lander, WY 82520**

Suggested Citation:
Orabona, A. C., C. K. Rudd, N. L. Bjornlie, Z. J. Walker, S. M. Patla, and R. J. Oakleaf. 2016. Atlas of Birds, Mammals, Amphibians, and Reptiles in Wyoming. Wyoming Game and Fish Department Nongame Program, Lander, USA.

TABLE OF CONTENTS

Figure 1. Latilongs in Wyoming	i
Introduction.....	1
Methods	2
Explanation of Records.....	4
Name and Status, WBRC.....	4
Mgmt. Status	6
Habitat	6
Comments.....	6
Distribution	7
Birds – Symbol Definitions.....	7
Mammals – Symbol Definitions.....	8
Amphibians and Reptiles – Symbol Definitions	8
Updates to the Atlas.....	8
Table 1. Seasonal Status Definitions.....	9
Table 2. Abundance Definitions.....	9
Table 3. Vertebrate Life Forms	10
Table 4. Management Status.....	11
Acknowledgements	12
Literature Cited.....	12
Birds.....	17
Mammals.....	130
Amphibians and Reptiles	162
Index	174
Index of Common Names	175
Index of Scientific Names	196
Appendix I.....	217
Rare and Unusual Bird Sighting Form	218
Rare and Unusual Bird Sighting Form – Photograph Only.....	220
How to Document Sightings of Rare and Unusual Birds	221
Mammal Observation Record Form	222
Appendix II.....	223
Documentation Requested for All Sightings (Birds)	224
Documentation Requested for First Latilong Sightings (Birds).....	225
Appendix III.....	226
Habitat Codes and Descriptions	227

Latitude and Longitude Degree Blocks in Wyoming



INTRODUCTION

In 1979, the Wyoming Game and Fish Department (Department) and the Bighorn Audubon Society compiled a working draft of the *Wyoming Avian Atlas*. The first edition of the *Wyoming Avian Atlas* was published in 1982 under the authorship of Robert (Bob) Oakleaf, Helen Downing, Bert Raynes, Meg Raynes, and Oliver K. Scott. Updates were provided each year in the Threatened, Endangered, and Nongame Bird and Mammal Investigations Annual Completion Reports. In 1981, Scott Findholt, Bob Oakleaf, and Bill Long published a *Working Draft of the Wyoming Mammal Atlas*. Updates were provided in some Annual Completion Reports.

In 1991, the Department published a revision of both atlases, titled the *Draft Distribution and Status of Wyoming Birds and Mammals*. This publication included all of the information in the *Avian Atlas* and the draft *Mammal Atlas*. Several hundred people, without whom this effort would not have been possible, contributed to the 1991 draft publication. A combined *Bird and Mammal Atlas*, edited by Bob Oakleaf, Bob Luce, Sharon Ritter, and Andrea Orabona Cerovski, was published in 1992.

Annual updates were provided to all persons who received the 1992 *Atlas*. These updates included hundreds of observations of birds and mammals, including many for which little information was previously available. A large number of bat observations were added due to a special project carried out between 1994 and 1996 to document bat distribution in Wyoming.

In 1997, Nongame Program biologist, John Priday, initiated an effort to gather data from a variety of sources to compile current distribution data for the amphibians and reptiles of Wyoming. After initial review within the Department, the decision was made to publish this information with the *Bird and Mammal Atlas*, and the title was changed to the *Atlas of Birds, Mammals, Reptiles, and Amphibians* to reflect this inclusion.

Updates to the 1997 *Atlas* continued as new information became available, and reprints occurred in 1999 and 2004. In 2005, the *Atlas* was made available on the Department's web site, negating the need for paper copies to be mailed. However, original hard copies and electronic files of both the *Atlas* and annual *Atlas* updates remain available from the Wyoming Game and Fish Department's Nongame Bird Biologist in Lander. The *Atlas* and annual *Atlas* updates are available on the Department's web site at <https://wgfd.wyo.gov/Wildlife-in-Wyoming/More-Wildlife>.

Since 2001, the American Ornithologists' Union (AOU) had made numerous modifications to genus and species names and phylogenetic order from their original check-list published in 1983 (AOU 1983, 2016). In addition, many updates to the avian portion of the *Atlas* were added as a result of observers submitting their sightings to the Wyoming Bird Records Committee, on the WyoBirds list serve, and through eBird. Therefore, a new version titled the *Atlas of Birds, Mammals, Amphibians, and Reptiles* was published in July 2004, April 2009, June 2012, and March 2016 to reflect these major changes.

The information in the *Atlas* documents past observations, but it also encourages use of the data to record new observations and distribution records for each species represented. The Department encourages wildlife watchers who want to know where a particular species can be found to use the *Atlas* for this purpose. Although latilongs cover hundreds of square miles, habitat associations, status, and abundance are good indicators of where to look for a particular bird, mammal, amphibian, or reptile. The Life Form codes and comments give further information that we hope will enhance your search for, and enjoyment of, Wyoming's wildlife.

Resource managers, consultants, and wildlife biologists are encouraged to use the *Atlas* to obtain basic information on occurrence and distribution of birds, mammals, amphibians, and reptiles for use in resource management decisions.

METHODS

Information sources for the initial versions of the *Atlas* included the Department's Wildlife Observation System, the Wyoming Natural Diversity Database, the University of Wyoming Museum collection, Wyoming mammals recorded in museum collections in other institutions, and published literature.

A number of published observation records were included in the *Atlas*. Jane Dorn's monumental bibliography of Wyoming bird references (Dorn 1978) was an important source of published literature. Other sources for birds included: Knight (1902), Grave and Walker (1913), Skinner (1925), McCafferty (1930), McCreary (1939), Flack (1976), Frinzel (1964), Pettingill and Whitney (1965), Gniadek (1977), Nye et al. (1979), and Dorn and Dorn (1990).

Sources for the mammal portion of the *Atlas* included: Findley (1951), Hall and Kelson (1959), Burt and Grossenheider (1980), Long (1965), Brown (1967a, 1967b), Pattie and Verbeek (1967), Maxell and Brown (1968), Lechleitner (1969), Clark (1973a, 1973b), Turner (1974), Bissell (1978), Rothwell et al. (1978), Clark and Dorn (1979), Stromberg (1979), Hall and Thomas (1979), Clark et al. (1980), Wyoming Game and Fish Department (1980), Bee et al. (1981), Belitsky (1981), Hall (1981), Clark and Stromberg (1987), Zeweloff and Collett (1988), Baker et al. (2003), Bradley et al. (2014), and Buskirk (2016).

Amphibian and reptile information came primarily from two published sources (Baxter and Stone 1980, Parker and Anderson 2001) and was compiled by John Priday, along with his personal observations from 1997 through 1999. Since 2008, the Department's herpetologists have provided updates and new information for the amphibians and reptiles sections. Common and scientific names have been updated based on current information from the Society for the Study of Amphibians and Reptiles (Crother 2012).

Raptor aerial surveys; intensive waterbird and secretive marsh bird surveys; implementation of the Monitoring Wyoming's Birds program (Leukering et al. 2001) and the Integrated Monitoring in Bird Conservation Regions program (Hanni et al. 2014); bird banding projects; and projects supported by federal State Wildlife Grants, Wyoming Governor's Big Game License Coalition,

and Wyoming Governor's Endangered Species Account funding have resulted in a significant amount of new bird data. Files of state and federal agencies were reviewed for unpublished data. Other files reviewed included the Breeding Bird Survey, coordinated in the United States by the U.S. Geological Survey Patuxent Wildlife Research Center, and records maintained at the Bird Banding Laboratory. Random sightings from biologists and the public, a statewide small mammal survey conducted between 1989 and 1995, bat surveys conducted from 1994-1998 and 2008-2016, and sensitive species inventories conducted on the Shoshone National Forest from 1995-1997 have contributed a great deal of well documented mammal data. Additional recent project funded by State Wildlife Grants, the Wyoming Governor's Endangered Species Account, and other sources have provided opportunities to expand these data, including projects ranging from documenting presence and distribution of wolverines, fishers, and spotted skunks to those focusing on small mammals, including northern flying squirrels, Preble's meadow jumping mice, water voles, and American pikas. The expanding use of technology to detect species, such as remote cameras and acoustic detectors, provides a more comprehensive understanding of the full wildlife community in an area. Information from numerous individuals who recorded and submitted personal observations of birds and mammals continues to be an important and invaluable source of data.

Unfortunately, the sources of some older observations are unknown. Since 1989, however, observations used for updating the *Atlas* have included only those for which written documentation existed (e.g., Wildlife Observation Form, Rare and Unusual Bird Sighting Form, Mammal Observation Record). Source information for each observation is filed with the Department's Nongame Program in the Lander Regional Office.

In 1989, at the encouragement of Sam Fitton, a group of five expert birders and the Department's Nongame Bird Biologist formed the Wyoming Bird Records Committee (WBRC). The goals of the WBRC are threefold. 1) To solicit, organize, and maintain records, documentation, photographs, audio recordings, and any other material relative to the birds of Wyoming. 2) To review records of new or rare species or species difficult to identify and offer an intelligent, unbiased opinion of the validity or thoroughness of these reports. From these reviews, the WBRC will develop and maintain an Official State List of Wyoming's Birds. 3) To disseminate useful and pertinent material concerning the field identification of Wyoming birds in order to assist Wyoming birders in increasing their knowledge and skill. The WBRC is interested in promoting and maintaining quality and integrity in the reporting of Wyoming bird observations, and it treats all bird records as significant historical documents. The WBRC operates under a set of bylaws approved in 1991, and updated in 1998, 2007, 2008, 2015, and 2016.

Prior to 1990, the Department's Nongame Bird Biologist reviewed all rare and unusual bird documentations. In many cases, if observations of rare and unusual birds were accepted by the regional editor of *American Birds* and published in that magazine, the Department also accepted the record. Since 1990, the WBRC has reviewed old records and documentation, and all new records. Some previously accepted records have been deleted due to lack of acceptable documentation. Those records that required review and were accepted by the WBRC are noted

in the “Distribution” section. Periodically, the WBRC also reviews records previously or currently reviewed by the Yellowstone Bird Records Committee to provide consistency across the state. Eventually, the WBRC would also like to review records previously reviewed by the Yellowstone Bird Observation Committee to provide consistency across the entire state.

If one watches birds solely for their own enjoyment, a written description of observations is unnecessary. However, to record an observation of a rare or unusual bird for scientific purposes, written description is essential. This is necessary whether one is a highly experienced observer or a beginning birder. Compilers of regional bird lists and similar documents must maintain records that are scientifically sound. A Rare and Unusual Bird Sighting Form is included in the *Atlas*, as well as an explanation of how to document bird sightings (Appendix I); these are also available from the Department’s Nongame Bird Biologist. Questions or comments about the WBRC should be directed to the Department’s Nongame Bird Biologist.

The validity of records of rare mammalian species was determined by expert review of written and photographic documentation. Mammal records relied almost entirely upon confirmed identification of specimens repositied in collections or turned in to Department personnel or personnel of other agencies. A completed Mammal Observation Record form accompanied many observations. From these forms, recognized experts were able to determine whether a sufficient number of characteristics of the animal and its behavior were described and similar species were eliminated using accepted criteria. A copy of the Department’s Mammal Observation Record form is included in the *Atlas* (Appendix I) or available from the Department’s Nongame Mammal Biologist.

Expert reviewers and contributors included personnel from the Wyoming Game and Fish Department Nongame Program, Trophy Game Program, Herptile Program, and Wildlife Division in the Lander Region; University of Wyoming Zoology Department; Wyoming Natural Diversity Database; Museum of Southwestern Biology; and Denver Museum of Natural History. This review procedure added many valuable records. In particular, Department biologists Laura Beard, Stan Harter, Zack Lange, Dan Thompson, and Leah Yandow contributed updated distribution information for many species within their areas of expertise.

EXPLANATION OF RECORDS

Since the 1992 *Atlas*, we have attempted to minimize the use of codes as much as possible. The following is a brief explanation of the format using the Upland Sandpiper as an example:

Name and Status	WBRC
Upland Sandpiper	(FL)
<i>Bartramia longicauda</i>	
261.0 UPSA	
Summer resident	
Uncommon	
Life form 5	

The first and second lines contain the common and scientific names. For birds, species names and the order in which they occur follow the most current AOU list (AOU 2016). For mammals, species names and order follows Bradley et al. (2014). For amphibians and reptiles, species names follow Crother (2012), and species order follows Baxter and Stone (1980).

For birds only, an (AS) or (FL) under WBRC indicates that the Wyoming Bird Records Committee requests submission of a Rare and Unusual Bird Sighting Form for all sightings of the species (AS), or for the first observation in each latitude/longitude degree block (latilong) and all nesting observations (FL). Species for which a Rare and Unusual Bird Sighting Form is requested appear in Appendix II. Observations of avian species new to Wyoming or an unusual breeding record for any avian species also require submission of a Rare and Unusual Bird Sighting Form.

Observations or specimens of mammals, except those for common and abundant species, require documentation because of the difficulty of distinguishing similar species.

The number on the third line is the species code assigned for the Department's computerized database system, which is used to store and maintain all recorded wildlife observations. For birds, it is the same as the AOU number.

The four-letter code following the species code is a unique alpha code that can be used on data forms or in field notes. The bird codes are those used by bird banders and are listed in the North American Bird Banding Manual Volume 1 (U.S. Fish and Wildlife Service and Canadian Wildlife Service 2015). However, revisions to the manual are only available online (<http://www.pwrc.usgs.gov/bbl/manual/speclist.cfm>). Only full species, as recognized by the AOU, are included. For mammals, amphibians, and reptiles, the first two letters of the genus followed by the first two letters of the species form the four-letter alpha code. In the case of subspecies for some mammals and reptiles, the alpha code is comprised of the first two letters of the genus followed by the first letter of the species and the first letter of the subspecies. One mammal code is identical to a bird code; therefore, the code for the mammal differs from the normal protocol. When codes for more than one mammal or herptile species were identical, a different code was arbitrarily assigned.

The fourth line refers to seasonal status – the time of the year the species occurs in Wyoming. Seasonal status categories and definitions are presented in Table 1 (page 9).

Line five refers to the abundance of the species in Wyoming. Abundance categories and definitions are presented in Table 2 (page 9).

Life Form is a term used to describe where a particular animal lives, feeds, and reproduces. Life Form codes for migrants, winter residents, and accidental species are not included in the *Atlas*. Table 3 (page 10) lists the life form categories for each species according to Thomas (1979). One life form not included in Thomas (1979) was added to Table 3 (#17, human-built structures).

Mgmt. Status

The management or legal status of a species often significantly affects how it is managed and which state or federal agency, or section within an agency, is responsible for management. The management status codes and definitions, and the Division or Program within the Department responsible for management of Wyoming's species are listed in Table 4 (page 11) and are consistent with the Wyoming Game and Fish Commission Regulations.

The Nongame Program is responsible for managing species legally designated as Nongame, which includes Species of Greatest Conservation Need with a designated Native Species Status (NSS) and Neotropical Migratory Birds (NTMB). Using the Native Species Status matrix that first appeared in the Nongame Bird and Mammal Plan (Oakleaf et al. 1996) and has since been updated and included in the Department's State Wildlife Action Plan (Wyoming Game and Fish Department 2010, 2017 [In Press]), the Department has designated Species of Greatest Conservation Need with a Native Species Status of 1, 2, 3, 4, or U (unknown).

The Nongame Program is also responsible for cooperative management with the U.S. Fish and Wildlife Service of the following federally listed Threatened or Endangered species: Wood Stork, Whooping Crane, Piping Plover, Least Tern, Preble's meadow jumping mouse, black-footed ferret, and Canada lynx.

Habitat

For birds, the habitats listed are those in which the species would be expected to breed or nest in Wyoming. For birds that do not breed in Wyoming and for all mammals, amphibians, and reptiles, the habitats listed are those in which the species is most likely to be found. Habitat use was determined from personal knowledge and literature review, especially Ehrlich et al. (1988) and the Birds of North America series for birds. Habitat type descriptions are from the Department's Wildlife Observation System and are listed in Appendix III (Wyoming Game and Fish Department 1992).

Comments

This column describes nesting or breeding sites, food habits, etc. The literature source or observer's name is given for unusual records, avian species observed in three or fewer latilongs, and for historical observations. The sources used for information on food habits and nesting or denning sites were Ehrlich et al. (1988) and The Birds of North America (Rodewald 2015) for birds, and Clark and Stromberg (1987) and Fitzgerald et al. (1994) for mammals. Baxter and Stone (1980) was consulted for information on amphibians and reptiles.

References used to prepare this publication are listed in the Literature Cited section. Locations listed in the original literature source were used to situate observations to latilong. McCreary (1939) was a major source of bird observation locations. Most pre-1965 mammal data were taken from Long (1965). Efforts by the Wyoming Natural Diversity Database and the

Department acquired records for many Wyoming mammal specimens repositied in museum collections throughout the United States. Baxter and Stone (1980) provided distribution information for amphibians and reptiles.

Distribution

Example for Western Small-footed Myotis:

<u> </u>	<u> O </u>	<u> B </u>	<u> B </u>	<u> O </u>	<u> O </u>	<u> O </u>
<u> O </u>	<u> O </u>	<u> B </u>	<u> B </u>	<u> O </u>	<u> h </u>	<u> a </u>
<u> O </u>	<u> </u>	<u> B </u>	<u> O </u>	<u> B </u>	<u> B </u>	<u> O </u>
<u> O </u>	<u> B </u>	<u> B </u>	<u> O </u>	<u> a </u>	<u> B </u>	<u> B </u>

Each underline represents a latitudinal/longitudinal degree block, or latilong, as shown on the map of Wyoming (Figure 1). A latilong encompasses a rectangle covering one degree of latitude by one degree of longitude, and is numbered on the map. For example, latilong 27 is bounded by 41° and 42° latitude and 105° and 106° longitude. Detailed maps, such as those available from the Bureau of Land Management or U.S. Geological Survey, should be used for determining exact boundaries of a specific latilong.

The definitions of the symbols used for birds, mammals, amphibians, and reptiles are as follows:

Birds – Symbol Definitions

- B Nest or young dependent upon parent birds was observed.
- b Circumstantial evidence of nesting.
- O The species has been observed, but there was no evidence of nesting. The observation may have been recorded during any season of the year, but observations are most likely to correspond with seasonal status listed in Table 1.
- H Historical nesting record, prior to 1950. No recent data to indicate the species still nests or occurs in that latilong.
- h Historical record with no indication of nesting.
- ? Record that needs further verification.
- No verified records.
- * Record has been reviewed and accepted by the Wyoming Bird Records Committee (WBRC).
- # Record has been reviewed and accepted by the Yellowstone Bird Observation Committee (YBOC).
- + Record has been reviewed and accepted by both the WBRC and YBOC.

Mammals – Symbol Definitions

- B Nest, dependent young, juvenile animals, lactating or post-lactation females, or males in breeding condition were observed.
- b Animals were observed and, due to limited mobility, breeding is assumed (bats and large ungulates are highly mobile and are not automatically placed in this category).
- O The species has been observed but, due to the mobility of the species' group and lack of factors listed under (B), breeding cannot be assumed (applies to bats and large ungulates).
- a The species has been detected using acoustic recording equipment and additional verification is warranted. Due to the mobility of the species' group and lack of factors listed under (B), breeding cannot be assumed (applies to bats only).
- H Historical breeding record, prior to 1965. No recent data to indicate the species still breeds or occurs in that latilong.
- h Historical record of occurrence before 1965. No recent data to suggest occurrence.
- No verified records.

Amphibians and Reptiles – Symbol Definitions

- B Nest, eggs, or juvenile animals were observed.
- b Animals were observed and, due to limited mobility and the time of year the observation occurred, breeding is assumed.
- O The species has been observed but, due to the lack of factors listed under (B) or (b), breeding cannot be assumed.
- ? A species observation record has not been verified, but occurrence is assumed based on the proposed species range.
- h Historical record of occurrence. No recent data to suggest occurrence.
- No verified records.
- * Record was obtained from the University of Wyoming Museum.
- # Record was obtained from the Wyoming Natural Diversity Database.

UPDATES TO THE ATLAS

Since 2005, the *Atlas* has been available on the Department's web site. This option replaces the previous paper copies of both the *Atlas* and annual updates that were mailed to individuals on a mailing list maintained by the Department's Nongame Program in the Lander Regional Office. The 2016 *Atlas* is available on the Department's website, and can be downloaded using the following link:

https://wgfd.wyo.gov/WGFD/media/content/PDF/Wildlife/Nongame/WILDLIFE_ANIMALATLAS.pdf.

Annual *Atlas* updates will be posted periodically so interested parties can obtain the most current version of the *Atlas* and track changes until the next updated version is available. Updates can be found on the Department's website at: <https://wgfd.wyo.gov/>. Click on the "Wildlife in Wyoming" link and the "More" link. Updates can be found under the "Wyoming Species Atlas" heading.

TABLE 1. SEASONAL STATUS DEFINITIONS.

SEASONAL STATUS	DEFINITION
Resident	A species that inhabits Wyoming year-round.
Summer Resident	A species that inhabits Wyoming during the summer breeding season (June-August) and migrates out of the state during the non-breeding season. Breeding may or may not be documented; however, records should span all of the summer months (especially July).
Winter Resident	A species that inhabits Wyoming during the winter (especially January) but is absent during the summer.
Migrant	A species found in Wyoming only during spring and/or autumn migration. For birds, records are especially lacking for July and January.
Accidental	A species with accidental, unpredictable occurrence of individual animals because its range does not include Wyoming.
Peripheral	A species whose range extends only into a small portion of Wyoming.
Undetermined	A species that occurs in Wyoming but seasonal status is uncertain.
Extinct	A species that no longer exists in Wyoming or elsewhere.

TABLE 2. ABUNDANCE DEFINITIONS.

ABUNDANCE	DEFINITION
Rare	A species that occupies only a small percentage of the preferred habitat within its range or is found throughout its range in extremely low densities. The species or its sign is seldom encountered while using survey techniques that could be expected to indicate its presence.
Uncommon	A species that is common only in limited areas within its range or is found throughout its range in relatively low densities. Intensive surveying is usually required to locate the species or its sign.
Common	A species that inhabits much of the preferred habitat within its range. The species or its sign is usually encountered while using survey techniques that could be expected to indicate its presence.
Abundant	A species that inhabits most of the preferred habitat within its range. The species or its sign is expected to be encountered while using survey techniques that could be expected to indicate its presence.
Unknown	A species whose existence within Wyoming is verified but insufficient information is available to determine abundance. The species is difficult to observe without specialized survey techniques.
Controlled	A species that currently exists in Wyoming only with human assistance.

TABLE 3. VERTEBRATE LIFE FORMS (modified from Thomas 1979).

LIFE FORM NUMBER	REPRODUCES	FEEDS
1	In water	In water
2	In water	On the ground, in shrubs and/or in trees
3	On the ground around water or emergent vegetation	In water, on the ground, in shrubs, or in trees
4	On cliffs, rims, talus, and/or in caves	On the ground or in the air
5	On the ground without specific water; cliffs, rims, or talus associations	On the ground
6	On the ground	In shrubs, trees, or the air
7	In shrubs	On the ground, in water, or in the air
8	In shrubs	In shrubs, trees, or the air
9	Primarily in deciduous trees	In shrubs, trees, or the air
10	Primarily in conifers	In shrubs, trees, or the air
11	In trees	On the ground, in shrubs, in trees, or in the air
12	On very thick branches	On the ground or in water
13	Excavates its own tree cavity	On the ground, in shrubs, in trees, or in the air
14	In a naturally occurring cavity or a cavity excavated by another species	On the ground, in water, or in the air
15	In an underground burrow	On or under the ground
16	In an underground burrow	In water or the air
17	On human-built structures	On the ground, in shrubs, in trees, or in the air

TABLE 4. MANAGEMENT STATUS.

CODE	DEFINITION	MANAGEMENT AUTHORITY
NSS ^a	Native Species Status 1, 2, 3, 4, or U	Wyoming Game and Fish Department
NG ^b	Nongame bird, mammal, or reptile	Wyoming Game and Fish Department Nongame Program
BG	Big game	Wyoming Game and Fish Department Regional Offices
TG	Trophy game	Wyoming Game and Fish Department Trophy Game Program
SG	Small game	Wyoming Game and Fish Department Regional Offices
GB	Game Bird	Wyoming Game and Fish Department Regional Offices
FB	Furbearer	Wyoming Game and Fish Department Regional Offices
PD	Predacious bird Predatory animal	Wyoming Game and Fish Department Regional Offices
F	Species taken for falconry	Wyoming Game and Fish Department Nongame Program
A	Amphibian	Wyoming Game and Fish Department Fisheries Program
NTMB ^c	Neotropical Migratory Bird	Wyoming Game and Fish Department Nongame Program
End.	Federally listed as endangered	U. S. Fish and Wildlife Service
Thr.	Federally listed as threatened	U. S. Fish and Wildlife Service

^a Species of Greatest Conservation Need with a Native Species Status of 1, 2, 3, 4, or U (Wyoming Game and Fish Department 2017, In Press).

^b All nongame birds in Wyoming are protected under the federal Migratory Bird Treaty Act, with the exception of the non-native Mute Swan, Rock Pigeon, Eurasian Collared-Dove, European Starling, and House Sparrow. Nongame mammals that are protected in Wyoming include the black-footed ferret, fisher, Canada lynx, northern river otter, American pika, and wolverine.

^c Neotropical Migratory Bird (breeds in Canada and the United States and migrates to Mexico, Central America, South America, or the Caribbean for the non-breeding season).

ACKNOWLEDGEMENTS

Since the publication of the *Working Draft of the Wyoming Mammal Atlas* in 1981 and the *Wyoming Avian Atlas* in 1982, several hundred individuals have contributed information to update our knowledge of bird, mammal, amphibian, and reptile distribution and status in Wyoming. We would like to sincerely thank these observers for the time and effort they have expended to add to our knowledge of Wyoming's wildlife. We also applaud the efforts of the many observers who have sent information to us which did not result in *Atlas* updates, but added valuable information to the Department's Wildlife Observation System database.

We would also like to recognize the efforts of those dedicated individuals who have taken the time to carefully record their observations of rare and unusual birds and submit them to the Wyoming Bird Records Committee for review. These endeavors have added tremendously to our knowledge of the distribution and status of rare and unusual avian species in Wyoming.

Without the support of the many wildlife watchers, naturalists, and professional biologists in Wyoming, as well as those visiting from other states, it would not be possible to continue to update the information in this document. We are grateful for the help of all contributors.

LITERATURE CITED

- American Ornithologists' Union [AOU]. 1983. Check-list of North American Birds. Seventh edition. American Ornithologists' Union, Washington, D.C.
- American Ornithologists' Union [AOU]. 2016. Fifty-seventh supplement to the American Ornithologists' Union check-list of North American birds. *Auk* 133:544-560.
- Baker, R. J., L. C. Bradley, R. D. Bradley, J. W. Dragoo, M. D. Engstrom, R. S. Hoffman, C. A. Jones, F. Reid, and C. Jones. 2003. Revised checklist of North American mammals north of Mexico, 2003. Occasional Papers, Museum of Texas Technical University, Number 229.
- Baxter, G. T., and M. D. Stone. 1980. Amphibians and Reptiles of Wyoming. Wyoming Game and Fish Department, Cheyenne.
- Bee, J. W., G. E. Glass, R. S. Hoffmann, and R. R. Patterson. 1981. Mammals in Kansas. Museum of Natural History, University of Kansas, Lawrence.
- Belitsky, D. 1981. Small Mammals of the Salt Wells – Pilot Butte Planning Unit. Bureau of Land Management, Wyoming.
- Bissell, S. 1978. Colorado Mammal Distribution, Latilong Study. Colorado Division of Wildlife, Denver.

- Bradley, R. D., L. K. Ammerman, R. J. Baker, L. C. Bradley, J. A. Cook, R. C. Dowler, C. Jones, D. J. Schmidley, F. B. Stangl Jr., R. A. Van Den Bussche, and B. Wursig. 2014. Revised checklist of North American mammals north of Mexico, 2014. Occasional Papers, Museum of Texas Technical University, Number 327.
- Brown, L. N. 1967a. Ecological distribution of mice in the Medicine Bow Mountains of Wyoming. *Ecology* 48:677-680.
- Brown, L. N. 1967b. Ecological sampling of six species of shrews and comparison of sampling methods in the central Rocky Mountains. *Journal of Mammalogy* 48:617-623.
- Burt, W. H., and R. P. Grossenheider. 1980. *A Field Guide to the Mammals*. Houghton Mifflin Company, Boston, MA.
- Buskirk, S. W. 2016. *Wild Mammals of Wyoming and Yellowstone National Park*. University of California Press, Oakland.
- Clark, T. W. 1973a. Distribution and reproduction of shrews in Grand Teton National Park, Wyoming. *Northwest Science* 47:128-131.
- Clark, T. W. 1973b. Local distribution and interspecies interactions in microtines, Grand Teton National Park, Wyoming. *Great Basin Naturalist* 33:205-217.
- Clark, T. W., and R. D. Dorn. 1979. Rare and endangered vascular plants and vertebrates of Wyoming.
- Clark, T. W., V. Saab, and D. Casey. 1980. A partial bibliography of Wyoming mammals. *Northwest Science* 54:55-67.
- Clark, T. W., and M. Stromberg. 1987. *Mammals in Wyoming*. University of Kansas Press, Lawrence.
- Crother, B. I., editor. 2012. Scientific and Standard English Names of Amphibians and Reptiles of North America North of Mexico, with Comments Regarding Confidence in our Understanding. *SSAR Herpetological Circular* 39:1-92.
- Dorn, J. L. 1978. *Wyoming Ornithology: A History and Bibliography with Species and Wyoming Area Indexes*. Bureau of Land Management and Wyoming Game and Fish Department.
- Dorn, J. L., and R. D. Dorn. 1990. *Wyoming Birds*. Mountain West Publishing, Cheyenne, WY.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. *The Birder's Handbook*. Simon and Schuster, Inc., New York, NY.

- Findley, J. S. 1951. Habitat preference of four species of *Microtus* in Jackson Hole, Wyoming. *Journal of Mammology* 32:118-120.
- Fitzgerald, J. P., C. A. Meaney, and D. M. Armstrong. 1994. *Mammals of Colorado*. University Press of Colorado, Niwot, CO.
- Flack, J. A. D. 1976. Bird populations of aspen forest in western North America. *Ornithological Monograph Number 19*.
- Frinzel, J. E. 1964. Avian populations of four herbaceous communities in southeastern Wyoming. *Condor* 66:496-510.
- Gniadek, S. 1977. Some aspects of avian ecology following the 1974 Trail Creek and 1976 Divide fires in Yellowstone National Park. Unpublished Research Report. Yellowstone National Park.
- Grave, B. H., and E. P. Walker. 1913. *The birds of Wyoming, with an explanation of recent changes in their distribution, economic aspects also considered*. University of Wyoming, Laramie.
- Hall, E. R., and K. R. Kelson. 1959. *The Mammals of North America, Volume I*. Ronald Press, New York, NY.
- Hall, E. R. 1981. *The Mammals of North America, Volume II*. John Wiley and Sons, New York, NY.
- Hall, F. C., and J. W. Thomas. 1979. Silvicultural options. Pages 128-147 in J.W. Thomas, editor. *Wildlife Habitats in Managed Forests*. USDA, Forest Service, Agriculture Handbook Number 553.
- Hanni, D. J., C. M. White, N. J. Van Lanen, J. J. Birek, J. M. Berven, and M. A. McLaren. 2014. Integrated Monitoring in Bird Conservation Regions (IMBCR): field protocol for spatially-balanced sampling of landbird populations. Unpublished report. Rocky Mountain Bird Observatory, Brighton, CO. Online: <http://rmbo.org/v3/avian/DataCollection.aspx>.
- Knight, W. C. 1902. *The birds of Wyoming*. University of Wyoming Agricultural Experiment Station Bulletin Number 55.
- Lechleitner, R. 1969. *Wild Mammals of Colorado: Their Appearance, Habits, Distribution, and Abundance*. Pruett Publishing Company, Boulder, CO.
- Leukering, T., M. F. Carter, A. Panjabi, D. Faulkner, and R. Levad. 2001. *Monitoring Wyoming's Birds: The Plan for Count-based Monitoring*. Rocky Mountain Bird Observatory, Brighton, CO.

- Long, C. A. 1965. The mammals of Wyoming. Museum of Natural History, University of Kansas 14:493-758.
- Maxell, M. H., and L. N. Brown. 1968. Ecological distribution of rodents on the high plains of eastern Wyoming. Southwestern Naturalist 13:143-158.
- McCafferty, C. E. 1930. An annotated and distributional list of the birds of Wyoming. M.S. Thesis, University of Wyoming, Laramie.
- McCreary, O. 1939. Wyoming Bird Life. Burgess Publishing Company, Minneapolis, MN.
- Nye, D. L., M. Back, and H. Hinchman. 1979. Birds of the Upper Wind River Valley. USDA Forest Service, Shoshone National Forest.
- Oakleaf, B., A. O. Cerovski, and B. Luce. 1996. Nongame Bird and Mammal Plan. Wyoming Game and Fish Department, Cheyenne.
- Parker, J., and S. Anderson. 2001. Identification Guide to the Herptiles of Wyoming. Wyoming Cooperative Fish and Wildlife Research Unit, Laramie.
- Pattie, D. L., and N. A. M. Verbeek. 1967. Alpine mammals of the Beartooth Mountains. Northwest Science 41:110-117.
- Pettingill, O. S., Jr., and N. R. Whitney, Jr. 1965. Birds of the Black Hills. Special Publication Number 1. Cornell Laboratory of Ornithology, Ithaca, NY.
- Rodewald, P. (Editor). 2015. The Birds of North America. Cornell Laboratory of Ornithology, Ithaca, NY. Online: <https://birdsna.org>.
- Rothwell, R. G., G. Skutches, J. Straw, C. Sax, and H. Harju. 1978. A partial bibliography of the mammals of Wyoming and adjacent states with special reference to density and habitat affinity. Bureau of Land Management Contract Number YA-512-CT8-126.
- Skinner, M. P. 1925. The birds of Yellowstone Park. Roosevelt Wildlife Bulletin 3:1-192.
- Stromberg, M. 1979. Working List of Special Mammals – Wyoming Natural Heritage Program. The Nature Conservancy.
- Thomas J. W. 1979. Wildlife habitats in managed forests, the Blue Mountains of Oregon and Washington. USDA Forest Service, Agricultural Handbook No. 553.
- Turner, R. W. 1974. Mammals of the Black Hills of South Dakota and Wyoming. Museum of Natural History, University of Kansas 60:1-178.

- U. S. Fish and Wildlife Service, and Canadian Wildlife Service. 2015. Species Table and Recommended Band Sizes. Online: <http://www.pwrc.usgs.gov/bbl/manual/speclist.cfm>.
- Wyoming Game and Fish Department. 1980. Densities of animals on potential coal lease sites in south-central Wyoming. Bureau of Land Management Contract Number YA-512-CT7-7.
- Wyoming Game and Fish Department. 1992. Wildlife Observation System. Wyoming Game and Fish Department Biological Services Section, Cheyenne.
- Wyoming Game and Fish Department. 2010. State Wildlife Action Plan. Wyoming Game and Fish Department, Cheyenne.
- Wyoming Game and Fish Department. 2017 (In Press). State Wildlife Action Plan. Wyoming Game and Fish Department, Cheyenne.
- Zeveloff, S. I., and F. R. Collett. 1988. Mammals of the Intermountain West. University of Utah Press, Salt Lake City.

Birds

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Fulvous Whistling-Duck <i>Dendrocygna bicolor</i> 178.0 FUWD Accidental Controlled Life Form	(AS)		Marshes, lakes, rivers.	Feeds on aquatic vegetation. L8: D. Browne, 20 May 2007.	— — — — — — — — O* —
Greater White-fronted Goose <i>Anser albifrons</i> 171.0 GWFG Migrant Rare Life Form	(FL)	GB	Marshes, lakes, agricultural areas.	Feeds on grasses, aquatic vegetation, waste grains, aquatic invertebrates, berries, seeds.	O* — — — — O* O* O* O — — O* — — O* — — O* O* O — — O* — — O — — — — — O* O O*
Snow Goose <i>Chen caerulescens</i> 169.0 SNGO Migrant Uncommon Life Form		GB	Marshes, lakes, agricultural areas.	Feeds on sedges, grasses, aquatic vegetation, waste grains, aquatic invertebrates, insects. Two subspecies are recognized, Greater and Lesser; the Lesser Snow Goose occurs in Wyoming. Once common in Wyoming during migration (McCreary 1939). Includes "Blue Goose".	O O O O O O O O O O — — O O — O O O — — O O O O O O — — O O O
Ross's Goose <i>Chen rossii</i> 170.0 ROGO Migrant Uncommon Life Form	(FL)	GB	Marshes, lakes, agricultural areas.	Feeds on aquatic vegetation, grasses, aquatic invertebrates, waste grains.	O* O* — — O O* — — O* O — — O* — — O* — — — — — — O* — — O* — — O O O O* — — — — O* O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Brant <i>Branta bernicla</i> 174.0 BLBR Accidental Rare Life Form	(AS)		Lakes.	Feeds on aquatic vegetation, grasses, algae, moss, aquatic invertebrates. L3: F.M. Lentsch, 20 November 1992. L10, 17, 26, 27: McCreary (1939). Includes "Black Brant".	— — O* — — — — — — h — — — — — — h — — O* — — — — — — h h — —
Cackling Goose <i>Branta hutchinsii</i> 172.2 CACG Accidental Rare Life Form	(FL)		Marshes, lakes, and rivers in association with other habitats, especially wet-moist meadows, sedge meadows, and agricultural areas.	Feeds on grasses, sedges, grain, insects, crustaceans. L28: G. Johnson, 3 January 2005.	— O* — — — — O*
Canada Goose <i>Branta canadensis</i> 172.0 CAGO Resident Abundant Life Form 3		GB	Marshes, lakes, and rivers in association with other habitats, especially wet-moist meadows, sedge meadows, and agricultural areas.	Nests on the ground or in an old nest of a heron or raptor. Feeds on grasses, sedges, grain, insects, crustaceans.	B B
Mute Swan <i>Cygnus olor</i> 178.2 MUSW Accidental Controlled Life Form	(AS)		Not native to North America; brought from other continents for zoos, parks, and private collections.	Feeds on aquatic vegetation, grasses, aquatic invertebrates, waste grain. Is very aggressive and will out-compete native swans. L19: J. Lawrence, 5 January 1999. L19: B. Walgren, 1 May 2011.	— O* — — — — — — — — — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Trumpeter Swan <i>Cygnus buccinator</i> 181.0 TRUS Resident Uncommon Life Form 3	(FL)	GB-no season NSS2	Marshes, lakes, rivers.	Nests on a muskrat house, a very small island, or a piece of floating bog. Feeds mainly on aquatic vegetation; also insects, aquatic invertebrates. Most of the population winters in Idaho. L7: "B" represents one pair. L19: L. Layton and F. Layton, 11 December 1994. L23: T. Koerner, 6 June 1997. L24: R. Steenberg, 16 November 1997.	<u>B</u> <u>B</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O*</u> <u>O</u> <u> </u> <u>O</u> <u>B*</u> <u>O*</u> <u> </u> <u>O</u> <u> </u> <u>h</u>
Tundra Swan <i>Cygnus columbianus</i> 180.0 TUSW Winter resident Uncommon Life Form		GB-no season	Marshes, lakes, rivers.	Feeds on aquatic vegetation; also aquatic invertebrates, waste grain.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Whooper Swan <i>Cygnus cygnus</i> 179.0 WHOS Accidental Rare Life Form	(AS)		Lakes, marshes, rivers.	Feeds on aquatic vegetation. L1: T. McEneaney, 17 January 2004.	<u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Wood Duck <i>Aix sponsa</i> 144.0 WODU Summer resident Common Life Form 14		GB	Cottonwood-riparian, marshes, lakes, rivers.	Nests in a tree cavity. Feeds mostly on aquatic vegetation; also aquatic invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Gadwall <i>Anas strepera</i> 135.0 GADW Resident Abundant Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground in dense vegetative cover. Feeds mostly on aquatic vegetation; also aquatic invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Eurasian Wigeon <i>Anas penelope</i> 136.0 EUWI Accidental Rare Life Form	(AS)		Marshes, lakes, rivers, mostly below 8,000 feet.	Feeds mostly on grasses, aquatic vegetation; also aquatic invertebrates. L21, 28: McCreary (1939).	<u>O+</u> <u>—</u> <u>O</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>h</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>O*</u> <u>h</u>
American Wigeon <i>Anas americana</i> 137.0 AMWI Resident Abundant Life Form 3		GB	Marshes, lakes, rivers, mostly below 8,000 feet.	Nests on the ground, often far from water. Feeds on aquatic vegetation, grasses, waste grain, aquatic invertebrates. Winter populations are considerably smaller than during other seasons.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
American Black Duck <i>Anas rubripes</i> 133.0 ABDU Accidental Rare Life Form	(AS)		Marshes, lakes.	Feeds on aquatic vegetation, aquatic invertebrates, worms, seeds, tubers. L1: Wilson Bulletin 48:136-137. L20: O.K. Scott. L22, 27: McCreary (1939). Species is difficult to identify and hybridizes with Mallards.	<u>h</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>h</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>h</u> <u>—</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Mallard <i>Anas platyrhynchos</i> 132.0 MALL Resident Abundant Life Form 3		GB	Marshes, lakes, and rivers in association with most habitats below 9,000 feet.	Nests on the ground near water. Feeds primarily on plant material, aquatic vegetation, seeds, green plants, waste grains, aquatic invertebrates, insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Mottled Duck <i>Anas fulvigula</i> 134.0 MODU Accidental Rare Life Form	(AS)		Lakes, ponds, marshes.	Feeds on grass seeds, aquatic vegetation, rice, aquatic invertebrates, few small fishes. L27: S. Billerman, 6 May 2011	— O*
Blue-winged Teal <i>Anas discors</i> 140.0 BWTE Summer resident Common Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground in good vegetative cover. Feeds mostly on seeds and vegetative parts of aquatic vegetation; also aquatic invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Cinnamon Teal <i>Anas cyanoptera</i> 141.0 CITE Summer resident Common Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground in dense cover near a marsh or shallow wetland. Feeds mostly on seeds of aquatic vegetation.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Northern Shoveler <i>Anas clypeata</i> 142.0 NSHO Summer resident Common Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground near water. Feeds on zooplankton, seeds of aquatic vegetation, aquatic invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Northern Pintail <i>Anas acuta</i> 143.0 NOPI Resident Abundant Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground in low or sparse vegetative cover. Feeds mostly on vegetation including seeds, green plants.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Garganey <i>Anas querquedula</i> 139.2 GARG Accidental Rare Life Form	(AS)		Marshes and lakes in association with most habitats below 8,000 feet.	Feeds on aquatic vegetation, aquatic invertebrates. L8: D. Wile, 26 December 2001.	— — — — — — — <u>O*</u> —
Green-winged Teal <i>Anas crecca</i> 139.0 AGWT Resident Abundant Life Form 3		GB	Marshes and lakes in association with most habitats below 8,000 feet.	Nests on the ground in tall vegetative cover near water. Feeds mostly on seeds of aquatic vegetation; also vegetative parts of aquatic vegetation, aquatic invertebrates. Winter populations are considerably smaller than during other seasons.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Canvasback <i>Aythya valisineria</i> 147.0 CANV Summer resident Uncommon Life Form 3		GB	Marshes, lakes, rivers.	Nest is a floating platform, or nests on a muskrat house. Feeds on aquatic vegetation, aquatic invertebrates. Summer populations are considerably smaller than during migration.	B O O O B B O B B B O O O O B B O B B O O B O O B O B O
Redhead <i>Aythya americana</i> 146.0 REDH Summer resident Common Life Form 3		GB	Marshes, lakes, rivers.	Nests on a floating platform in shallow water or on the ground very close to water. Feeds on aquatic vegetation, aquatic invertebrates.	B B B B B O B b B B O O O O B b B B B O O B B B B b B B
Ring-necked Duck <i>Aythya collaris</i> 150.0 RNDU Summer resident Common Life Form 3		GB	Marshes, lakes.	Nests on the ground near water or on a floating structure. Feeds on aquatic vegetation, aquatic invertebrates.	B B O O O O O B O O O O O O O B B b O O O O O O O B O O
Tufted Duck <i>Aythya fuligula</i> 149.1 TUDU Accidental Rare Life Form	(AS)		Lakes.	Feeds on aquatic vegetation, aquatic invertebrates. L8: E. Cole, 1 April 1999. L19: J. Herold and V. Herold, 11 March 1994. L27: M. Gochfield, 10 April 1966, probably an escaped bird.	— — — — — — — O* — — — — — — — — — — — O* — — — — — — — — O* — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Greater Scaup <i>Aythya marila</i> 148.0 GRSC Migrant Rare Life Form	(FL)	GB	Marshes, lakes, rivers.	Feeds mainly on mollusks; also other aquatic invertebrates, aquatic vegetation.	O+ _ _ _ O* _ _ O O O _ _ _ _ O _ O* _ O* _ O _ O* O* _ _ O _
Lesser Scaup <i>Aythya affinis</i> 149.0 LESC Summer resident Common Life Form 3		GB	Marshes, lakes, rivers.	Nests on the ground in open, dry habitat near water. Feeds on aquatic invertebrates, aquatic vegetation. Common during migration.	B B b O O O O B B O O O O O B b b B B O O B O B B B B b
Harlequin Duck <i>Histrionicus histrionicus</i> 155.0 HARD Summer resident Uncommon Life Form 3		GB NSS3	Rivers and lakes in mountainous areas.	Usually nests on the ground under a shrub near a river. Feeds on aquatic invertebrates.	B O _ B _ _ _ B B B _ O _ _ B O O _ O _ _ _ O _ _ O _ _
Surf Scoter <i>Melanitta perspicillata</i> 166.0 SUSC Migrant Rare Life Form	(FL)	GB	Lakes, rivers.	Feeds on aquatic invertebrates.	O+ _ _ _ O* _ O O* O* _ _ _ _ _ _ _ _ O O _ _ O* O* _ _ O* O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
White-winged Scoter <i>Melanitta fusca</i> 165.0 WWSC Migrant Uncommon Life Form	(FL)	GB	Lakes, rivers.	Feeds on aquatic invertebrates.	<u>0</u> <u>—</u> <u>—</u> <u>0</u> <u>0*</u> <u>—</u> <u>0</u> <u>0</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0</u> <u>—</u> <u>0*</u> <u>—</u> <u>0</u> <u>0</u> <u>0*</u> <u>0*</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>0</u> <u>—</u>
Black Scoter <i>Melanitta americana</i> 163.0 BLSC Accidental Rare Life Form	(AS)		Lakes, rivers.	Feeds on aquatic invertebrates.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>0</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u>
Long-tailed Duck <i>Clangula hyemalis</i> 154.0 LTDU Migrant Rare Life Form	(FL)	GB	Lakes.	Feeds on aquatic invertebrates.	<u>0+</u> <u>0</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>0</u> <u>0*</u> <u>0</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>0</u> <u>0*</u> <u>—</u> <u>0*</u> <u>0*</u> <u>—</u> <u>—</u> <u>0</u> <u>—</u>
Bufflehead <i>Bucephala albeola</i> 153.0 BUFF Resident Uncommon Life Form 14		GB	Aspen; cottonwood-riparian; marshes; lakes and rivers associated with lodgepole pine, Douglas fir, and other or mixed coniferous forests. Lower elevation habitats during migration and winter.	Nests in a cavity, usually in a dead tree. Feeds on aquatic invertebrates.	<u>B</u> <u>B</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>B</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>b</u> <u>0</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Common Goldeneye <i>Bucephala clangula</i> 151.0 COGO Resident Common Life Form 14		GB	Aspen; cottonwood-riparian; marshes; lakes and rivers associated with lodgepole pine, Douglas fir, and other or mixed coniferous forests. Lower elevation habitats during migration and winter.	Nests in a cavity, usually in a dead tree. Feeds mostly on aquatic invertebrates; also aquatic vegetation.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Barrow's Goldeneye <i>Bucephala islandica</i> 152.0 BAGO Resident Common Life Form 14		GB	Aspen; cottonwood-riparian; marshes; lakes and rivers associated with lodgepole pine, Douglas fir, and other or mixed coniferous forests. Lower elevation habitats during migration and winter.	Nests in a cavity, usually in a dead tree. Feeds mostly on aquatic invertebrates.	<u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>O</u> <u>B</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Hooded Merganser <i>Lophodytes cucullatus</i> 131.0 HOME Resident Uncommon Life Form 14		GB	Marshes, lakes, rivers.	Nests in a cavity, usually in a dead tree. Feeds mostly on fish; also crustaceans, aquatic invertebrates. Breeding in Wyoming is expected.	<u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u>
Common Merganser <i>Mergus merganser</i> 129.0 COME Resident Common Life Form 3 14		GB	Cottonwood-riparian, marshes, lakes, rivers, especially large rivers.	Nests in a cavity in a deciduous tree, earthen bank, or rock crevice. Feeds mostly on fish.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Red-breasted Merganser <i>Mergus serrator</i> 130.0 RBME Summer resident Uncommon Life Form 3		GB	Lakes, rivers.	Nests on the ground under vegetative cover. Feeds mostly on fish.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Ruddy Duck <i>Oxyura jamaicensis</i> 167.0 RUDU Summer resident Common Life Form 3		GB	Marshes, lakes, rivers.	Nests on the ground in tall emergent vegetation. Feeds on aquatic invertebrates, aquatic vegetation.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u>
Ruddy Shelduck <i>Tadorna ferruginea</i> 141.2 RUSH Accidental Controlled Life Form			Not native to North America; brought from other continents for zoos, parks, and private collections.	Feeds on aquatic invertebrates. L28: G. Johnson and B. Templeton, 18 October 1997, probably released or escaped from captivity.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>
Common Shelduck <i>Tadorna tadorna</i> 141.1 COMS Accidental Controlled Life Form			Not native to north America; brought from other continents for zoos, parks, and private collections.	Feeds on aquatic invertebrates. L2: S. Ahalt, 1995, probably released or escaped from captivity.	<u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Northern Bobwhite <i>Colinus virginianus</i> 289.0 NOBO Resident Uncommon Life Form 5	(AS)	GB	Cottonwood-riparian, riparian shrub, agricultural areas.	Nests on the ground. Feeds on vegetation, seeds, insects. At the periphery of its range. Numerous releases of captive stock statewide, especially in L3. L2: R. Young, 1 June 1997; J. Cicco and L. Cicco, 4 June 1997. L5: R. Adams and R. Adams, October 1992. L9: A. Moscicki and M. Kenney, 17 May 2005.	<u> </u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u>
Chukar <i>Alectoris chukar</i> 288.2 CHUK Resident Common Life Form 5		GB	Mountain-foothills shrublands.	Nests on the ground amid rocks or shrubs. Feeds on seeds; leaves; insects, especially grasshoppers.	<u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>O</u>
Gray Partridge <i>Perdix perdix</i> 288.1 GRPA Resident Uncommon Life Form 5		GB	Basin-prairie shrublands, riparian shrub, grasslands, agricultural areas.	Nests on the ground. Feeds on grain, grasses, forbs, insects. Also called Hungarian Partridge.	<u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u>
Ring-necked Pheasant <i>Phasianus colchicus</i> 309.1 RPHE Resident Common Life Form 5		GB	Agricultural areas, cottonwood-riparian, riparian shrub, sedge meadows.	Nests on the ground in grass or weeds. Feeds on a variety of plants and animals.	<u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u>B</u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Ruffed Grouse <i>Bonasa umbellus</i> 300.0 RUGR Resident Common Life Form 5		GB	Coniferous forests, aspen, mountain-foothills shrublands, riparian shrub (willow, hawthorn, wild plum, dogwood, water birch, bog birch, alder), logged forests.	Nests on the ground. Feeds mostly on buds, leaves, flowers, seeds, fruit; also insects.	<u>B</u> <u>B</u> <u> </u> <u>B</u> <u>B</u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u> </u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Greater Sage-Grouse <i>Centrocercus urophasianus</i> 309.0 GRSG Resident Common Life Form 5		GB NSS4	Basin-prairie and mountain-foothills shrublands, wet-moist meadows, alfalfa, irrigated native meadows.	Nests on the ground under a sagebrush shrub. Feeds on vegetation, especially sagebrush in the winter; forbs; insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
White-tailed Ptarmigan <i>Lagopus leucura</i> 304.0 WTPT Resident Rare Life Form 5	(AS)	GB-no season	Alpine grasslands, alpine willow, spruce-fir forests.	Nests on the ground, occasionally under a shrub. Feeds mainly on buds, leaves, flowers of willows and other shrubs. At the periphery of its range. L1: Condor 24:204 (1927) and Wilson Bulletin 66:60 (20 July 1952). L26: several records. Many records are lacking details in L2, 4, 8, 9, 16, 17.	<u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u> </u>
Dusky Grouse <i>Dendragapus obscurus</i> 297.0 DUGR Resident Common Life Form 5		GB	Coniferous forests, aspen, willow, mountain-park meadows, logged forests.	Nests on the ground. Feeds mostly on leaves, especially conifer needles.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Sharp-tailed Grouse <i>Tympanuchus phasianellus</i> 308.0 STGR Resident Common Life Form 5		GB NSS4	Mixed grass prairie grasslands, mountain-foothills shrublands, willow, irrigated native meadows.	Nests on the ground in grass or under a shrub. Feeds on vegetation, seeds, berries, insects. Includes the "Columbian" Sharp-tailed Grouse (NSS4) in L1, 8, 16, 25; also Sharp-tailed Grouse/Greater Sage-Grouse hybrid in L5.	<u>O</u> <u> </u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u> </u> <u>h</u> <u> </u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u>b</u> <u>B</u>
Greater Prairie-Chicken <i>Tympanuchus cupido</i> 305.0 GPCH Accidental Rare Life Form	(AS)		Tallgrass prairie.	Feeds on seeds, greens, insects. L28: five records since 1900 (McCreary 1939, Wyoming Wildlife 16(2):32, Auk 51:537). More recent observations by B. Luce and G. Robinson.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u>
Wild Turkey <i>Meleagris gallopavo</i> 310.0 WITU Resident Common Life Form 5		GB	Coniferous forests, especially ponderosa pine-Douglas fir, cottonwood-riparian, oak and other mixed deciduous forests; woodland-chaparral; agricultural areas.	Nests on the ground among grasses or shrubs. Feeds on a variety of plants and animals.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>B</u> <u>B</u>
Pied-billed Grebe <i>Podilymbus podiceps</i> 006.0 PBGR Summer resident Common Life Form 3		NG	Marshes and lakes below 8,000 feet. Also seen in streams.	Nest is a floating platform. Feeds on aquatic insects, snails, fish, frogs.	<u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Horned Grebe <i>Podiceps auritus</i> 003.0 HGR Summer resident Uncommon Life Form 3		NG	Marshes, lakes. Also seen in streams.	Nest is a floating platform. Feeds on aquatic insects, crustaceans, mollusks, fish. L2: U. Kepler (one pair in July 1978 at Beck Lake). Breeding range is north of Wyoming. Listed as a summer resident in Yellowstone National Park (Skinner 1925).	<u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u>
Red-necked Grebe <i>Podiceps grisegena</i> 002.0 RGR Summer resident Rare Life Form 3	(AS)	NG	Marshes, shallow lakes edged with reeds and sedges, rivers.	Nest is a floating platform of reeds in shallow water. Feeds on aquatic insects, aquatic invertebrates, fish, amphibians.	<u>O+</u> <u>O*</u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u>O*</u> <u>O*</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>h</u> <u> </u> <u>O*</u> <u>O*</u> <u> </u> <u> </u> <u>h</u> <u>O*</u>
Eared Grebe <i>Podiceps nigricollis</i> 004.0 EAGR Summer resident Common Life Form 3		NG	Marshes and lakes below 8,000 feet. Will also use rivers when not breeding.	Nest is a floating platform of reeds in shallow water near rooted aquatic vegetation. Feeds mostly on aquatic insects, aquatic larvae.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Western Grebe <i>Aechmophorus occidentalis</i> 001.0 WGR Summer resident Common Life Form 3		NG NSSU	Marshes, lakes.	Nest is a floating platform of reeds in shallow water near rooted aquatic vegetation. Feeds mostly on fish.	<u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Clark's Grebe <i>Aechmophorus clarkii</i> 001.1 CLGR Summer resident Uncommon Life Form 3		NG NSSU	Marshes, lakes.	Nest is a floating platform of reeds in shallow water near rooted aquatic vegetation. Feeds mostly on fish.	O O B _ O _ O O _ B _ O _ _ O O B O b O O O B O _ O O O
Rock Pigeon <i>Columba livia</i> 313.1 ROPI Resident Abundant Life Form 4 17		NG	Most habitats below 8,000 feet, especially agricultural areas, cliffs, and urban areas.	Nests on a building ledge, bridge, cliff, or bank. Feeds on grain, seeds, some green leaves. One of four bird species that can be legally killed without a license or permit.	B B B B B B b B B b b B B B b B b B B O B B B b B B
Band-tailed Pigeon <i>Patagioenas fasciata</i> 312.0 BTPI Migrant Rare Life Form	(AS)	NG NTMB	Ponderosa pine, limber pine, pine-juniper, woodland-chaparral.	Feeds on seeds, berries, grain.	O* O _ O* O* _ _ O _ _ _ _ O _ _ O O* _ O _ O* _ _ _ _ _ O _
African Collared-Dove <i>Streptopelia roseogrisea</i> 315.2 AFCD Accidental Controlled Life Form			Origin and native country uncertain; long domesticated and worldwide in captivity. Feral populations occur in parks and woodlands around human habitation.	Feeds on seeds, grain. Range has been expanding since the 1990s. L28: J. Dorn, 19-20 July 1997.	_ O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Eurasian Collared-Dove <i>Streptopelia decaocto</i> 315.9 ECDO Resident Common Life Form 11		NG	Not native to North America; brought from other continents for the pet trade; released in the wild in 1974. Resident populations now inhabit urban areas.	Nests in a coniferous tree in urban areas; may also use a deciduous tree. Feeds on seeds, grain. Range has been expanding since the 1980s. One of four bird species that can be legally killed without a license or permit. L2: first breeding record, D. Oudin and C. Axthelm, 28 August-21 September 2001. L28: first accepted record, G. Johnson, 16 May 1998.	_ <u>B*</u> _ <u>B*</u> _ <u>B</u> <u>O*</u> <u>O</u> <u>O*</u> <u>B</u> <u>O</u> _ _ _ <u>O*</u> _ <u>B*</u> _ <u>O*</u> _ <u>O*</u> <u>B</u> <u>B</u> _ <u>O</u> <u>O</u> _ <u>B*</u>
White-winged Dove <i>Zenaida asiatica</i> 319.0 WWDO Accidental Rare Life Form	(FL)	NTMB	Riparian woodlands and thickets.	Feeds on seeds, grain, some fruit. Range has been expanding in recent years.	_ <u>O*</u> _ _ _ <u>O*</u> _ <u>O*</u> <u>O*</u> <u>O*</u> _ _ _ _ _ <u>O*</u> <u>O*</u> _ <u>O*</u> _ <u>O*</u> _ <u>O*</u> _ <u>O*</u> <u>O*</u> <u>O*</u> <u>O*</u>
Mourning Dove <i>Zenaida macroura</i> 316.0 MODO Summer resident Abundant Life Form 5 7 11 17		GB NTMB	All habitats below 8,500 feet.	Nests in a fork of a tree, on the ground, or on a deserted nest. Feeds on seeds, waste grain.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Passenger Pigeon <i>Ectopistes migratorius</i> 315.0 PASP Extinct Life Form			Deciduous forests.	Fed on nuts, berries, seeds, grain. Species is now extinct; historical records in McCreary (1939).	_ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ <u>h</u> <u>h</u> _ _ _ _ _ _ _

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Yellow-billed Cuckoo <i>Coccyzus americanus</i> 387.0 YBCU Summer resident Uncommon Life Form 3	(FL)	NG NSSU NTMB Thr.	Cottonwood-riparian below 7,000 feet, urban areas.	Nests usually in a shrub. Feeds on insects, especially hairy caterpillars. L8: S. Bassett and R. Bassett, 23 July 1992. L11: J. Allen, 17 July 1992. L23: A. Pedersen, 29 June 1994. Species is listed as federally threatened as of November 2014.	<u> </u> <u> </u> <u>O*</u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>b</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>b</u>
Black-billed Cuckoo <i>Coccyzus erythrophthalmus</i> 388.0 BBCU Summer resident Uncommon Life Form 8	(FL)	NG NSS4 NTMB	Deciduous and mixed coniferous/deciduous forests; open woodlands, especially cottonwood-riparian; urban areas.	Nest is placed horizontally against a tree trunk; also on a log, occasionally in vine tangles. Feeds primarily on hairy caterpillars; also mollusks, fish, small vertebrates, berries. L1: T. McEneaney, 27 July 1994. L28: D. Bradley, 28 August 1987.	<u>O*</u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u>
Lesser Nighthawk <i>Chordeiles acutipennis</i> 421.0 LENI Accidental Rare Life Form	(AS)		Open habitats. Feeds in the air over large, open areas.	Feeds on insects. L28: S. Miller, 30 May 2011.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>
Common Nighthawk <i>Chordeiles minor</i> 420.0 CONI Summer resident Abundant Life Form 6		NG NSS4 NTMB	Open and semi-open habitats below 8,500 feet, including open forests and clearings, prairie-grasslands, sagebrush, rock outcrops, ponderosa pine forests. Feeds in the air over most habitats, especially aquatic and agricultural areas.	Nests on the ground. Feeds on insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Common Poorwill <i>Phalaenoptilus nuttallii</i> 418.0 COPO Summer resident Uncommon Life Form 6		NG NTMB	A variety of habitats below 8,000 feet including pine-juniper, woodland-chaparral, basin prairie and mountain-foothills shrublands, grasslands, agricultural areas.	Nests on the ground. Feeds on insects.	_ <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Black Swift <i>Cypseloides niger</i> 422.0 BLSW Undetermined Unknown Life Form	(AS)		Feeds in the air over coniferous and mixed deciduous/coniferous forests, juniper woodlands, basin-prairie shrublands and mountain-foothills shrublands.	Feeds on flying insects and arthropods. L22: J. Saucier, 15 June 2013. L4: J. Murdock, 4 July 2015.	_ _ _ <u>O*</u> _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ <u>O*</u> _ _ _ _ _ _ _ _
Chimney Swift <i>Chaetura pelagica</i> 423.0 CHSW Summer resident Rare Life Form 4 17	(FL)	NG NTMB	Feeds in the air over many habitats below 7,500 feet, especially urban areas.	Originally nested in a tree hollow, now in a chimney or other suitable human-built structure. Feeds on insects. Species is expanding its range westward. L5: H. Downing, 2 September 1981. L8: B. Raynes and M. Baker, 21 May 1994. L21: K. Kranik, 4 June 2004.	_ _ _ _ <u>O*</u> <u>O</u> _ _ <u>O*</u> _ _ _ _ <u>O</u> <u>O</u> <u>O</u> _ _ _ _ <u>O</u> <u>b</u> <u>O*</u> _ _ _ _ <u>O</u> _ <u>b</u>
Vaux's Swift <i>Chaetura vauxi</i> 424.0 VASW Accidental Rare Life Form	(AS)		Feeds in the air over coniferous and mixed deciduous/coniferous forests.	Feeds on flying insects, some spiders. L15: R. Steenberg, 31 May 2004.	_ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ <u>O*</u> _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
White-throated Swift <i>Aeronautes saxatalis</i> 425.0 WTSW Summer resident Common Life Form 4		NG NTMB	Feeds in the air over most habitats with cliffs below 9,000 feet.	Nests deep in a crack or crevice of a rock wall. Feeds on flying insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u>
Magnificent Hummingbird <i>Eugenes fulgens</i> 426.0 MAHU Accidental Rare Life Form	(AS)	NTMB	Alpine grasslands, mixed oak-conifer forests, mountain canyons.	Feeds primarily on nectar, occasionally on insects. L8: B. Raynes, 2-8 July 1982; American Birds 36(6):1002. L9: W. Moon, 15-22 July 1988.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Ruby-throated Hummingbird <i>Archilochus colubris</i> 428.0 RTHU Accidental Rare Life Form	(AS)	NTMB	Deciduous and mixed deciduous/coniferous forests, open areas with scattered trees.	Feeds on nectar, insects, spiders, tree sap from woodpecker holes. L19: V. Herold, 7-9 June 1999.	<u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Black-chinned Hummingbird <i>Archilochus alexandri</i> 429.0 BCHU Summer resident Peripheral Uncommon Life Form 7	(FL)	NG NSSU NTMB	Basin-prairie shrublands, riparian shrub, canyons, pinyon-juniper woodlands, urban areas, rural developments. A variety of habitats during migration.	Nests on a small limb of a deciduous tree, often near or over a stream. Feeds on nectar, insects. Nesting may occur in southwestern Wyoming.	<u> </u> <u>O</u> <u> </u> <u>O*</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u>b</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Anna's Hummingbird <i>Calypte anna</i> 431.0 ANHU Accidental Rare Life Form	(AS)	NTMB	Woodland-chaparral.	Feeds on nectar, insects. L5: observations by M. Collins, 1 August-15 September 1973; details accepted by P. Hall. L8: D. Birdsall, 19 October 2009.	— — — — — O* O* — — — — — — — — — — — — — — —
Broad-tailed Hummingbird <i>Selasphorus platycercus</i> 432.0 BTLH Summer resident Common Life Form 9 10		NG NTMB	Riparian shrub; mountain-foothills grasslands; coniferous forests; wet-moist meadows within Douglas Fir, Englemann spruce-subalpine fir, other coniferous or mixed forests, and aspen.	Usually nests on a horizontal limb of a deciduous or coniferous tree, near or over a stream. Feeds on nectar, insects.	B B — b b — O B b O O O — O B b B b B B O B b b B B B b
Rufous Hummingbird <i>Selasphorus rufus</i> 433.0 RUHU Summer resident Uncommon Life Form 8 10		NG NSS4 NTMB	Riparian shrub; mountain-foothills grasslands; wet-moist meadows within lodgepole pine, Douglas fir, other coniferous or mixed forests, aspen, and mountain-foothills shrublands.	Usually nests on a drooping limb of a coniferous or deciduous tree. Feeds on nectar, insects. Main breeding range is west of Wyoming. Most observations are presumed to be migrants.	B B O O O — O B O O O — O — O — O O O O O O B O O O O O
Calliope Hummingbird <i>Selasphorus calliope</i> 436.0 CAHU Summer resident Common Life Form 8 10		NG NSS4 NTMB	Coniferous forests, woodland-chaparral, mountain-foothills shrublands, riparian shrub, mountain park-meadows, alpine grasslands. Many habitats during migration.	Nests on a limb of a tree or on a conifer cone. Feeds on nectar, insects.	B B — B B O O B B O b — — — b O b — O — — O O B — O O —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Yellow Rail <i>Coturnicops noveboracensis</i> 215.0 YERA Accidental Rare Life Form	(AS)		Marshes.	Feeds on aquatic invertebrates, insects, fish. L1: T. McEneaney, 19 July 1996. L2: L. Siggins, August 1989.	O* O* _
Black Rail <i>Laterallus jamaicensis</i> 216.0 BLRA Accidental Rare Life Form	(AS)		Marshes, wet meadows.	Feeds on insects, small invertebrates, seeds. L27: T. Dini, 16 May 2010.	_ O*
Virginia Rail <i>Rallus limicola</i> 212.0 VIRA Summer resident Abundance unknown Life Form 3		GB NSSU	Marshes.	Nests on the ground, occasionally over water or mud. Feeds mostly on insects, aquatic invertebrates, seeds.	O O _ O* B _ O B O O _ _ O b _ _ b B O B B b b _ B B b
Sora <i>Porzana carolina</i> 214.0 SORA Summer resident Common Life Form 3		GB	Marshes, willow and other riparian shrub, irrigated native meadows.	Nest is a floating platform, occasionally nests in meadows. Feeds on seeds, insects, aquatic invertebrates.	B B B b B O B B B B B O O b B b B b B O B B B B b B B b

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Whooping Crane <i>Grus americana</i> 204.0 WHCR Summer resident Controlled Life Form 3	(AS)	NG End.	Wet-moist meadow grasslands, sedge meadows, irrigated native and introduced meadows, small grains, marshes.	Nests on the ground. Feeds on fish, small mammals, crustaceans, insects, roots, berries, grain. Birds in western Wyoming are from the Gray's Lake fostering project, except an historical sighting in L1 (Wilson Bulletin 42:198) indicated nesting in Yellowstone National Park; R. Drewien questioned validity. L28: Wyoming Wildlife 14(8):27, 1 bird for 2 weeks, August 1950.	<u>O</u> <u>O</u> _ _ _ _ _ <u>O*</u> <u>O</u> <u>O</u> _ _ _ _ _ <u>O*</u> <u>O*</u> <u>O</u> _ _ _ _ _ <u>O*</u> <u>O</u> _ _ _ _ _ <u>h</u>
Black-necked Stilt <i>Himantopus mexicanus</i> 226.0 BNST Summer resident Uncommon Life Form 3		NG	Marshes, ponds, flooded fields.	Nests on the ground close to water. Feeds primarily on aquatic invertebrates. L19: first breeding record, American Birds 30:983.	<u>O</u> <u>O</u> <u>B</u> _ _ <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ _ <u>O</u> <u>O</u> _ <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> _ _ <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u>
American Avocet <i>Recurvirostra americana</i> 225.0 AMAV Summer resident Common Life Form 3		NG	Marshes, ponds, shorelines.	Nests on the ground close to water among tufts of vegetation. Feeds on invertebrates, insects, some aquatic vegetation.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> _ _ <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u>
Black-bellied Plover <i>Pluvialis squatarola</i> 270.0 BBPL Migrant Uncommon Life Form		NG	Shorelines, aquatic areas.	Feeds on insects, some invertebrates.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ _ <u>O</u> _ _ <u>O</u> <u>O</u> _ <u>O</u> <u>O</u> <u>O</u> _ _ <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b																																								
American Golden-Plover <i>Pluvialis dominica</i> 272.0 AMGP Migrant Rare Life Form	(FL)	NG NTMB	Shorelines, aquatic areas.	Feeds on insects, some invertebrates.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>O</td></tr> <tr><td>O</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>O</td><td>O</td><td>—</td><td>—</td></tr> <tr><td>O</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O</td><td>O*</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O</td><td>O*</td></tr> </table>	—	—	—	—	—	—	O*	—	—	O	O	—	O*	—	—	—	O	O	—	—	O	—	—	—	—	—	O	O*	—	—	—	O*	—	—	—	—	—	—	O	O*
—	—	—	—	—	—	O*	—	—	O																																				
O	—	O*	—	—	—	O	O	—	—																																				
O	—	—	—	—	—	O	O*	—	—																																				
—	O*	—	—	—	—	—	—	O	O*																																				
Snowy Plover <i>Charadrius nivosus</i> 278.0 SNPL Summer resident, Peripheral Rare Life Form 3	(AS)	NG NSSU	Shorelines, aquatic areas.	Nests on the ground among tufts of grass. Feeds on insects, some invertebrates. L24: first breeding record, F. Luke, 1990.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td></tr> <tr><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>O*</td><td>B</td><td>B*</td><td>O*</td><td>O</td><td>O*</td><td>—</td><td>—</td><td>—</td></tr> </table>	—	—	—	—	—	—	—	—	—	O*	—	—	O*	—	—	—	O*	—	—	—	—	—	O*	—	—	—	O*	—	—	—	—	O*	B	B*	O*	O	O*	—	—	—
—	—	—	—	—	—	—	—	—	O*																																				
—	—	O*	—	—	—	O*	—	—	—																																				
—	—	O*	—	—	—	O*	—	—	—																																				
—	O*	B	B*	O*	O	O*	—	—	—																																				
Semipalmated Plover <i>Charadrius semipalmatus</i> 274.0 SEPL Migrant Uncommon Life Form		NG	Shorelines, aquatic areas.	Feeds primarily on invertebrates.	<table border="0"> <tr><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td></tr> <tr><td>O</td><td>O</td><td>O</td><td>—</td><td>O</td><td>O</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td></tr> <tr><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td><td>O</td></tr> </table>	O	O	O	O	O	O	O	O	O	O	O	O	O	—	O	O	—	—	—	—	—	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O
O	O	O	O	O	O	O	O	O	O																																				
O	O	O	—	O	O	—	—	—	—																																				
—	O	O	O	O	O	O	O	O	O																																				
O	O	O	O	O	O	O	O	O	O																																				
Piping Plover <i>Charadrius melodus</i> 277.0 PIPL Migrant Rare Life Form	(AS)	NG Thr. End.	Shorelines, aquatic areas.	Feeds primarily on invertebrates. Federally listed as Threatened, except in the Great Lakes watershed in IL, IN, MI, MN, NY, OH, PA, WI, and Ontario where it is federally listed as Endangered.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>O*</td></tr> <tr><td>—</td><td>b</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>O</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>O</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td><td>O</td></tr> </table>	—	—	—	—	—	—	O*	—	—	O*	—	b	—	—	—	—	—	—	—	—	O	—	—	—	—	—	O*	—	—	O	—	—	—	—	—	—	O*	—	—	O
—	—	—	—	—	—	O*	—	—	O*																																				
—	b	—	—	—	—	—	—	—	—																																				
O	—	—	—	—	—	O*	—	—	O																																				
—	—	—	—	—	—	O*	—	—	O																																				

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Killdeer <i>Charadrius vociferus</i> 273.0 KILL Summer resident Abundant Life Form 3		NG NTMB	Shorelines and aquatic areas associated with most habitats below 8,500 feet.	Nests on the ground, usually exposed. Feeds primarily on insects, some invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Mountain Plover <i>Charadrius montanus</i> 281.0 MOPL Summer resident Uncommon Life Form 5		NG NSSU NTMB	Shortgrass and mixed grass prairies, great basin-foothills grasslands, sagebrush-grasslands.	Nests on the ground, somewhat exposed. Feeds on insects, especially grasshoppers.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Upland Sandpiper <i>Bartramia longicauda</i> 261.0 UPSA Summer resident Uncommon Life Form 5	(FL)	NG NSSU NTMB	Eastern great plains grasslands, dryland grass pastures.	Nests in a depression on open ground, usually concealed by grass. Feeds on insects, terrestrial invertebrates, seeds.	<u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>B</u> <u>B</u> <u>—</u> <u>—</u> <u>—</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>B</u>
Whimbrel <i>Numenius phaeopus</i> 265.0 WHIM Migrant Rare Life Form	(FL)	NG	Marshes, ponds, lakes, shorelines.	Feeds on insects, berries, aquatic invertebrates.	<u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O*</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>—</u> <u>O*</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Long-billed Curlew <i>Numenius americanus</i> 264.0 LBCU Summer resident Uncommon Life Form 3		NG NSS3 NTMB	Sagebrush-grasslands; eastern great plains, great basin-foothills, mountain foothills, and wet-moist meadow grasslands; irrigated native meadows; with aquatic areas nearby. Also other agricultural areas and shorelines.	Nests on the ground near water, sometimes in a moist hollow. Feeds on insects, aquatic invertebrates. Locally common in some areas.	B B B B b B O B B b O O O O B b B b B O B B B b O B O O
Hudsonian Godwit <i>Limosa haemastica</i> 251.0 HUGO Migrant Rare Life Form	(AS)	NG NTMB	Shorelines, aquatic areas.	Feeds on insects, worms, aquatic invertebrates. L5: H. Downing and P. Hall, 9 September 1970; H. Downing, M. Collins, and P. Hall, 17 August 1976. L12: McCreary (1939). L19: F. Luke and others, 29 April 1990; G. Lawrence, 21 August 1992.	O* _ _ _ O* O _ _ _ O _ h _ _ _ _ _ _ O* _ _ _ _ _ _ _ O* O
Marbled Godwit <i>Limosa fedoa</i> 249.0 MAGO Migrant Uncommon Life Form		NG	Wet-moist meadow grasslands, marshes, aquatic areas, shorelines, irrigated native meadows.	Feeds mainly on aquatic invertebrates, insects.	O O O O O O O O O O O O _ O O O O O O O O O O O O O O O
Ruddy Turnstone <i>Arenaria interpres</i> 283.0 RUTU Migrant Rare Life Form	(FL)	NG	Shorelines, aquatic areas.	Feeds primarily on insects; also aquatic invertebrates, berries. L19: J. Lawrence, J. Herold, and V. Herold, 15 July 1993.	O _ _ _ O _ O _ _ O _ _ _ _ _ _ _ _ O* _ _ _ _ _ O _ _ _

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Baird's Sandpiper <i>Calidris bairdii</i> 241.0 BASA Migrant Common Life Form		NG NTMB	Shorelines, aquatic areas.	Feeds on insects; also some spiders.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Least Sandpiper <i>Calidris minutilla</i> 242.0 LESA Migrant Common Life Form		NG	Shorelines, aquatic areas.	Feeds on insects, aquatic invertebrates, some seeds.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
White-rumped Sandpiper <i>Calidris fuscicollis</i> 240.0 WRSA Migrant Rare Life Form	(FL)	NG NTMB	Shorelines, aquatic areas.	Feeds on insects, some seeds.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Buff-breasted Sandpiper <i>Calidris subruficollis</i> 262.0 BBSA Migrant Rare Life Form	(AS)	NG NTMB	Shorelines, aquatic areas.	Feeds mainly on insects.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Pectoral Sandpiper <i>Calidris melanotos</i> 239.0 PESA Migrant Uncommon Life Form		NG NTMB	Shorelines, aquatic areas.	Feeds primarily on insects; also some spiders, worms, seeds.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Semipalmated Sandpiper <i>Calidris pusilla</i> 246.0 SESA Migrant Uncommon Life Form		NG NTMB	Shorelines, aquatic areas.	Feeds on insects, some aquatic invertebrates.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Western Sandpiper <i>Calidris mauri</i> 247.0 WESA Migrant Uncommon Life Form		NG	Shorelines, aquatic areas.	Feeds almost entirely on insects, occasionally some invertebrates.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Short-billed Dowitcher <i>Limnodromus griseus</i> 231.0 SBDO Migrant Rare Life Form	(AS)	NG NTMB	Shorelines, aquatic areas.	Feeds mostly on aquatic invertebrates, insects, seeds. Probably more frequent than records indicate; difficult to separate from Long-billed Dowitcher.	<u>0</u> * <u>0</u> <u>0</u> <u>0</u> * <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> * <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> * <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> * <u>0</u> * <u>0</u> <u>0</u> * <u>0</u> * <u>0</u> *

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Long-billed Dowitcher <i>Limnodromus scolopaceus</i> 232.0 LBDO Migrant Common Life Form		NG	Shorelines, aquatic areas.	Feeds mostly on aquatic invertebrates, insects, seeds.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Wilson's Snipe <i>Gallinago delicata</i> 230.0 WISN Summer resident Common Life Form 3		GB	Wet-moist meadows, sedges, marshes, irrigated native meadows, willow and other mixed riparian shrub.	Nests in a small clump of grass under low vegetation. Feeds mostly on insects, earthworms. A few individuals remain throughout the winter.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
American Woodcock <i>Scolopax minor</i> 228.0 AMWO Accidental Rare Life Form	(AS)		Cottonwood-riparian, willow and other mixed riparian shrub, marshes.	Feeds on earthworms, insects. L8: J. Kirol, 16 April 1977. L17: D. Nelson and others, 19-21 and 29 December 1994 and 12 January 1995. L19: J. Lawrence, 7 November 1996.	<u> </u> <u>h</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u>h</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>
Spotted Sandpiper <i>Actitis macularius</i> 263.0 SPSA Summer resident Common Life Form 3		NG	Rocky shorelines of rivers and lakes.	Nests on an elevated site on the ground close to water. Feeds mainly on flying insects; also some invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Solitary Sandpiper <i>Tringa solitaria</i> 256.0 SOSA Migrant Common Life Form		NG	Marshes, shorelines.	Feeds primarily on aquatic insects.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Greater Yellowlegs <i>Tringa melanoleuca</i> 254.0 GRYE Migrant Common Life Form		NG	Shorelines, aquatic areas.	Feeds on small fish, insects, some invertebrates.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>
Willet <i>Tringa semipalmata</i> 258.0 WILL Summer resident Common Life Form 3		NG	Wet-moist meadow grasslands, marshes, irrigated native meadows, shorelines.	Nests on the ground, commonly on exposed beach or shore. Feeds on aquatic insects, some invertebrates.	<u>B</u> <u>b</u> <u>B</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>B</u> <u>b</u> <u>0</u> <u>0</u> <u>B</u> <u>0</u> <u>0</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>0</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>0</u> <u>B</u> <u>0</u>
Lesser Yellowlegs <i>Tringa flavipes</i> 255.0 LEYE Migrant Common Life Form		NG	Shorelines, aquatic areas.	Feeds primarily on terrestrial and aquatic insects.	<u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Wilson's Phalarope <i>Phalaropus tricolor</i> 224.0 WIPH Summer resident Common Life Form 3		NG NTMB	Marshes, lakes, shorelines.	Nests in a lined scrape on damp ground near water. Feeds mostly on aquatic invertebrates, seeds of aquatic plants.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Red-necked Phalarope <i>Phalaropus lobatus</i> 223.0 RNPH Migrant Uncommon Life Form		NG NTMB	Marshes, lakes, shorelines.	Feeds on aquatic invertebrates, some seeds.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Red Phalarope <i>Phalaropus fulicarius</i> 222.0 REPH Accidental Rare Life Form	(AS)	NTMB	Shorelines, aquatic areas.	Feeds mostly on aquatic insects; also some aquatic invertebrates.	<u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u> </u>
Pomarine Jaeger <i>Stercorarius pomarinus</i> 036.0 POJA Accidental Rare Life Form	(AS)		Large lakes.	Feeds primarily on rodents (lemmings); also seabird eggs. L12: J. Herold and O.K. Scott near border of L12 and L19, 15-16 May 1980.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Parasitic Jaeger <i>Stercorarius parasiticus</i> 037.0 PAJA Accidental Rare Life Form	(AS)		Large lakes.	Feeds on birds; also some small mammals, carrion. L5: H. Downing and M. Collins, 21 June 1985. L10: S. Fitton and T. Fitton, 24-25 November 1988.	— — — — — O* O — — — — — O* — — — — — O — — — — — O*
Long-tailed Jaeger <i>Stercorarius longicaudus</i> 038.0 LTJA Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on small mammals, birds, arthropods, insects. L27: J. Maley, 3 May 2012.	— O*
Long-billed Murrelet <i>Brachyramphus perdix</i> 023.0 LBMU Accidental Rare Life Form	(AS)		Large lakes.	Feeds on fish and aquatic invertebrates. L7: J. Hagen, 18 November 2003. L19: J. Gaither, B. South, and J. Herold, 23 July 1994.	— — — — — O* — — — — — — — — — — O* — — — — —
Ancient Murrelet <i>Synthliboramphus antiquus</i> 021.0 ANMU Accidental Rare Life Form	(AS)		Large lakes.	Feeds mainly on aquatic and planktonic invertebrates. Records of birds captured on land and photographed (L8: B. Raynes and S. Weiss, September 1981; L17: L. Serdiuk, November 1972). L15: S. Patla, 3 October 2003. L23: J. Lutz, 15 October 2002.	— — — — — O — — — — — O* — — — — — O — — — — — O* — — — — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Heermann's Gull <i>Larus heermanni</i> 057.0 HEEG Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on small fish, aquatic invertebrates, some insects. L19: O.K. Scott, 26 September 1984; V. Herold, 16 October 1993.	— — — — — — — — — — — — — — — O* — — — — — — — — — —
Mew Gull <i>Larus canus</i> 055.0 MEGU Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on a variety of dead and living animal and plant material. L1: T. McEneaney, 24 October 1991. L8: M. Windsor, 17 May 2011. L9: one record, 28 August 1893, Auk 36:276 (1919). L20: S. Dinsmore, 15 December 2001.	O* — — — — — O* h — — — — — — — — — — O* — — — — — — — — — —
Ring-billed Gull <i>Larus delawarensis</i> 054.0 RBGU Summer resident Common Life Form 3		NG	Large lakes, scavenges in most open habitats below 8,000 feet.	Nest is in the open or is concealed among rocks close to water. Feeds on a variety of dead and living animal and plant material. L27: breeding record in McCafferty (1930).	B O O O b O O O O B O O O O O O O O B O O O O O O O O O
California Gull <i>Larus californicus</i> 053.0 CAGU Summer resident Common Life Form 3		NG	Large lakes, scavenges in most open habitats below 8,000 feet.	Nests on sticks and dried weeds on the ground close to water, usually on an island or peninsula. Feeds on insects, small mammals, birds, bird eggs.	B O O O O O O O O B O O O — O O O O B O O O O O O O B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Herring Gull <i>Larus argentatus</i> 051.0 HERG Migrant Rare Life Form		NG	Large lakes, scavenges in most open habitats below 8,000 feet.	Feeds on a variety of foods from garbage to berries. L1: T. McEaney and D. Reinhart, 1 September 1992. L10: J. Lawrence, 1 April 1994.	O* _ _ _ O O O _ O O* _ O O O _ O _ _ O _ O O O O _ O B O
Thayer's Gull <i>Larus thayeri</i> 043.1 THGU Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on a variety of foods from garbage to berries. L8: S. Patla, 20 November 2009. L19: J. Lawrence, 4 November 2004.	_ _ _ _ O* _ _ O* _ _ _ _ _ _ _ _ _ _ O* _ _ _ _ _ _ _ O* _
Iceland Gull <i>Larus glaucooides</i> 043.0 ICGU Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds primarily on fish; also carrion, occasionally bird eggs, and nestlings. L27: J. Maley, 16 March 2013.	_ O* _
Lesser Black-backed Gull <i>Larus fuscus</i> 050.0 LBBG Accidental Rare Life Form	(AS except L19 & L27)		Aquatic areas.	Feeds on a variety of fish, insects, aquatic invertebrates, birds, bird eggs, small mammals; also berries, seeds, seaweed.	_ _ _ _ _ _ O* _ _ _ _ _ _ _ _ _ _ _ _ _ O* _ _ _ _ _ _ O* O* _

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Glaucous-winged Gull <i>Larus glaucescens</i> 044.0 GWGU Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on aquatic invertebrates, fish, garbage. L19: J. Lawrence, 5 May 2002.	— — — — — — — — — — — — — — — O* — — — — — — — — — — — — — —
Glaucous Gull <i>Larus hyperboreus</i> 042.0 GLGU Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on a variety of fish, marine invertebrates, carrion.	— — — — — O* — O* — — — — — O* — — — — — — — — — O* — — — — — — — — — — — — — — O* O*
Great Black-backed Gull <i>Larus marinus</i> 047.0 GBBG Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on a variety of fish, birds, bird eggs, small mammals, carrion, berries, grain, garbage. L27: D. McDonald, 14 April 2001; S. Billerman, 19 March 2013. L23: A. Moscicki, 2 January 2014.	— O* — — — — O* — — — —
Least Tern <i>Sternula antillarum</i> 074.0 LETE Accidental Rare Life Form	(AS)	End. NTMB	Large rivers, lakes.	Feeds on small fish, aquatic invertebrates. L21: McCreary (1939) reported observations near Torrington.	— — — — — O* — O* — — — — — O* — — — — — — — — — — — — — — — O* — — — — — — — — — — O* O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Caspian Tern <i>Hydroprogne caspia</i> 064.0 CATE Summer resident Uncommon Life Form 3		NG NSS3	Marshes, aquatic areas.	Nests on rocks or sand, often exposed and close to water, usually on an island or peninsula. Feeds on small fish, some aquatic invertebrates. L1: Skinner (1925) reported first breeding record from Yellowstone Lake.	<u>B</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>B</u> <u>O</u>
Black Tern <i>Chlidonias niger</i> 077.0 BLTE Summer resident Uncommon Life Form 3		NG NSS3 NTMB	Marshes, aquatic areas.	Nests on a floating mat of dead vegetation, often on a muskrat house. Feeds on insects, aquatic invertebrates, small fish.	<u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>b</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>O</u>
Common Tern <i>Sterna hirundo</i> 070.0 COTE Migrant Uncommon Life Form 3	(FL)	NG NTMB	Marshes, aquatic areas.	Feeds on small fish, some insects, aquatic invertebrates. Greatest numbers during migration.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u>
Arctic Tern <i>Sterna paradisaea</i> 071.0 ARTE Accidental Rare Life Form	(AS)	NTMB	Marshes, aquatic areas.	Feeds on fish, aquatic invertebrates, insects. L1: T. McEaney, 14 October 1997. L19: D. Faulkner, 18 September 2002.	<u>O</u> * <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> * <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Forster's Tern <i>Sterna forsteri</i> 069.0 FOTE Summer resident Common Life Form 3		NG NSS3	Marshes, aquatic areas.	Nests on the ground close to water or on a floating mat in emergents. Feeds on small fish, some insects, aquatic invertebrates. Greatest number during migration.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u>
Red-throated Loon <i>Gavia stellata</i> 011.0 RTLO Migrant Rare Life Form	(AS)	NG	Large lakes.	Feeds mostly on fish. L1: McCreary (1939). L5: M. Rozy, 10 November 1981; H. Downing and others, 24 October 1986, 15 May 1988. L10: R. Scott and B. Scott, 25 October and 15 November 2007. L27: S. Dinsmore, 24 May 2001.	<u>H</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u>
Pacific Loon <i>Gavia pacifica</i> 010.0 PALO Migrant Rare Life Form	(FL)	NG	Large lakes.	Feeds mostly on fish.	<u>O*</u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>O*</u> <u>O</u> <u>O*</u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O*</u> <u> </u> <u>O*</u> <u>O*</u> <u> </u>
Common Loon <i>Gavia immer</i> 007.0 COLO Summer resident Uncommon Life Form 3		NG NSS1	Lakes above 6,000 feet. Lower elevations during migration.	Nests next to water on a muskrat house, frequently on an island. Feeds on fish, aquatic invertebrates.	<u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Yellow-billed Loon <i>Gavia adamsii</i> 008.0 YBLO Accidental Rare Life Form	(AS)		Large lakes.	Feeds on fish.	— O* — — O* — O* — — — — — — — — — — — O* — — — — O* — —
Streaked Shearwater <i>Calonectris leucomelas</i> 088.1 STRS Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on fish, aquatic invertebrates, carrion. L27: D. Faulkner, M. Patz, F. Mauer, and D. McDonald, 13 June 2006.	— O* —
Wood Stork <i>Mycteria americana</i> 188.0 WOST Accidental Rare Life Form	(AS)	Thr.	Marshes, rivers.	Feeds mainly on fish. L1: McCreary (1939), 16 July 1925 and 28 June 1930. L2: S. Ahalt and others, 20 and 21 October 1994.	h O* —
Lesser Frigatebird <i>Fregata ariel</i> 128.2 LEFR Accidental Rare Life Form	(AS)		Aquatic areas.	Feeds on fish, aquatic invertebrates. L3: D. Faulkner, S. Ahalt, and B. Robertson, 11 July 2003.	— — O* —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Double-crested Cormorant <i>Phalacrocorax auritus</i> 120.0 DCCO Summer resident Common Life Form 3		NG	Cottonwood-riparian, lakes, rivers.	Nests on the ground on an island or in a tree, usually deciduous. Feeds on fish, crayfish, frogs, salamanders.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u>
American White Pelican <i>Pelecanus erythrorhynchos</i> 125.0 AWPE Summer resident Common Life Form 3		NG NSS4	Rivers, lakes, ponds.	Usually nests in colonies on islands. Nests in a scrape in the dirt. Feeds mostly on nongame fish, salamanders, crayfish. Eggs observed at Ocean Lake, but there has been no recent nesting record.	<u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u>
Brown Pelican <i>Pelecanus occidentalis</i> 126.0 BRPE Accidental Rare Life Form	(AS)		Lakes.	Feeds on fish. L28: McCreary (1939).	— <u>h</u>
American Bittern <i>Botaurus lentiginosus</i> 190.0 AMBI Summer resident Uncommon Life Form 3	(FL)	NG NSS3	Marshes.	Nests on dry ground above the water or mud in emergent vegetation. Feeds on fish, aquatic invertebrates, small mammals, insects.	<u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Least Bittern <i>Ixobrychus exilis</i> 191.0 LEBI Accidental Rare Life Form	(AS)		Marshes.	Feeds on fish, aquatic invertebrates, insects. L15: R.A. Grove, 1 and 17 June 1988. L22, 28: McCreary (1939).	— — — — — — — — — — — — — — B* — — — — — — — h — — — — — — h
Great Blue Heron <i>Ardea herodias</i> 194.0 GBHE Summer resident Common Life Form 3 12		NG NSS4	Cottonwood-riparian, lakes, rivers.	Nests in a tree, usually deciduous, occasionally coniferous; rarely on the ground. Feeds on fish, aquatic invertebrates, frogs, small mammals. Some individuals remain throughout the winter.	B B B B B B B B B B b B B O B B B B B B B B B B B B B B
Great Egret <i>Ardea alba</i> 196.0 GREG Accidental Rare Life Form	(FL)		Wet-moist meadow grasslands, marshes, other aquatic areas.	Feeds on fish, frogs, aquatic invertebrates.	O+ O* — O O* — O* O* O O* — — — — — — O* — O* O O* O* O* — — O* O* O*
Snowy Egret <i>Egretta thula</i> 197.0 SNEG Summer resident Rare Life Form 3		NG NSS3	Marshes, lakes, rivers.	Nests in bulrushes or cattails, or in a shrub on an island. Feeds on aquatic invertebrates, fish, insects. L27: first breeding record, 17 July 1967 (Findholt 1984). L28: Wyoming Wildlife 8(10):9, and observations of nonbreeders.	O O O O O O O O O B — — — — B B O O B O O B O O O B B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Black-crowned Night-Heron <i>Nycticorax nycticorax</i> 202.0 BCNH Summer resident Uncommon Life Form 3		NG NSS3	Marshes, lakes.	Nests in a low shrub or on the ground on an island or in bulrushes or cattails. Feeds mainly on fish.	<u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u>
Yellow-crowned Night-Heron <i>Nyctanassa violacea</i> 203.0 YCNH Accidental Rare Life Form	(AS)		Marshes, lakes.	Feeds on aquatic invertebrates, especially crustaceans. L5: H. Downing, 27 July and 22 August 1976. L27: F. Walter, 7 April 1983.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u>
White Ibis <i>Eudocimus albus</i> 184.0 WHIB Accidental Rare Life Form	(AS)		Marshes, lakes.	Feeds on aquatic invertebrates. L5: H. Downing, 12 and 13 September 1976, may have been released or may have escaped from captivity. L28: D. Lutz, 20 May 1994.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u>
Glossy Ibis <i>Plegadis falcinellus</i> 186.0 GLIB Accidental Rare Life Form	(AS)		Marshes, wet-moist meadows, lakes.	Feeds on aquatic invertebrates.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
White-faced Ibis <i>Plegadis chihi</i> 187.0 WFIB Summer resident Uncommon Life Form 3		NG NSS3	Marshes, wet-moist meadows, lakes, irrigated meadows.	Nests in bulrushes or cattails, occasionally on the ground on an island. Feeds mostly on aquatic invertebrates, insects.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u>
Black Vulture <i>Coragyps atratus</i> 326.0 BLVU Accidental Rare Life Form	(AS)		Coniferous and deciduous forests with open areas.	Feeds on carrion, occasionally live prey. L28: G. Johnson, 10 April 2012.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u>
Turkey Vulture <i>Cathartes aura</i> 325.0 TUVU Summer resident Common Life Form 5		NG NTMB	All habitats below 8,000 feet with open areas.	Nests on a cliff, in a cave, or in a hollow stump. Feeds on carrion.	<u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u>
Osprey <i>Pandion haliaetus</i> 364.0 OSPR Summer resident Common Life Form 12		NG NTMB	Lakes and rivers associated with lodgepole pine, Douglas fir, Englemann spruce-subalpine fir, and cottonwood-riparian.	Nests on a treetop, pole, or cliff near water. Feeds mostly on fish.	<u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>—</u> <u>—</u> <u>B</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Sharp-shinned Hawk <i>Accipiter striatus</i> 332.0 SSHA Summer resident Common Life Form 11		F NTMB	Coniferous forests, aspen, woodland-chaparral, cottonwood-riparian. Forages in a variety of habitats.	Nests in a tree. Feeds mostly on birds. Some individuals remain during the winter.	<u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Cooper's Hawk <i>Accipiter cooperii</i> 333.0 COHA Summer resident Common Life Form 11		F NTMB	Coniferous forests, aspen, cottonwood-riparian, woodland-chaparral. Forages in a variety of habitats.	Nests in a tree. Feeds mostly on birds, small mammals. Some individuals remain during the winter, especially in towns.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u>
Northern Goshawk <i>Accipiter gentilis</i> 334.0 NOGO Resident Common Life Form 11		F NSSU NTMB	Coniferous forests, especially Douglas fir and lodgepole pine, aspen. Forages in a variety of habitats.	Nests in a tree. Feeds mostly on birds, small mammals.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Harris's Hawk <i>Parabuteo unicinctus</i> 335.0 HRSH Accidental Rare Life Form	(AS)		Deciduous forests, especially near water.	Feeds mostly on small mammals. L10: B. Hargis, S. Hargis, and J. Taylor, 25 November 2007.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Red-shouldered Hawk <i>Buteo lineatus</i> 339.0 RSHA Accidental Rare Life Form	(AS)		Coniferous and deciduous forests near water.	Feeds mostly on small mammals; also some reptiles and amphibians. L1: D. Smith, 2 September 2003.	O* _
Broad-winged Hawk <i>Buteo platypterus</i> 343.0 BWHA Summer resident Rare Life Form 11	(FL)	NG NTMB	Grasslands, agricultural areas, urban areas.	Nests in a tree. Feeds on small mammals, birds, reptiles, insects. L4: first breeding record, B. Oakleaf and M. Wells, 7 July 2006.	O# _ O* B* O* _ _ O* _ O* _ _ _ _ _ O* _ O* O O _ _ O O O O O*
Swainson's Hawk <i>Buteo swainsoni</i> 342.0 SWHA Summer resident Common Life Form 11		F NSSU NTMB	Most habitats below 9,000 feet with open areas for foraging.	Nests in a tree, occasionally on a cliff. Feeds mostly on small mammals.	B b b B B B O B B b O B B B B B B B B B B B B B B B B B
Red-tailed Hawk <i>Buteo jamaicensis</i> 337.0 RTHA Resident Common Life Form 4 11		F NTMB	Most habitats below 9,000 feet with open areas for foraging.	Nests in a tree or on a cliff. Feeds mostly on rodents. Winter populations are considerably smaller than during other seasons. Includes "Harlan's Hawk".	B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Rough-legged Hawk <i>Buteo lagopus</i> 347.0 RLHA Winter resident Common Life Form		NG	Basin-prairie shrublands, grasslands, agricultural areas.	Feeds mostly on small mammals.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Ferruginous Hawk <i>Buteo regalis</i> 348.0 FEHA Resident Common Life Form 4 5 11		F NSS4 NTMB	Basin-prairie shrublands; eastern great plains, great basin-foothills, and mountain-foothills grasslands; rock outcrops; cottonwood-riparian.	Nests on a rock outcrop, the ground, a bank, or in a tree. Feeds mostly on small mammals. Winter populations are considerably smaller than during other seasons.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Golden Eagle <i>Aquila chrysaetos</i> 349.0 GOEA Resident Common Life Form 4 12		NG NSS4 NTMB	Most habitats with open areas for foraging.	Nests in a tree or on a cliff. Feeds mostly on small mammals, rabbits.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Barn Owl <i>Tyto alba</i> 365.0 BNOW Summer resident Uncommon Life Form 4 14 17	(AS)	NG	Basin-prairie shrublands, grasslands, agricultural areas.	Nests in a building, cave, or crevice. Feeds on rodents; sometimes birds.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>B</u> <u>O</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O</u> <u>—</u> <u>O</u> <u>b</u> <u>O</u> <u>—</u> <u>B</u> <u>O</u> <u>B</u> <u>—</u> <u>B</u> <u>O</u> <u>—</u> <u>O</u> <u>O*</u> <u>B*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Snowy Owl <i>Bubo scandiacus</i> 376.0 SNOW Winter resident Rare Life Form	(AS)	NG	Basin-prairie sagebrush grasslands, grasslands, agricultural areas.	Feeds primarily on small mammals.	<u>O</u> <u>O*</u> <u>O</u> <u>O*</u> <u>O*</u> <u>O</u> <u>O*</u> <u>—</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u>O*</u> <u>O*</u> <u>—</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u>
Northern Hawk Owl <i>Surnia ulula</i> 377.0 NHOW Accidental Rare Life Form	(AS)		Coniferous and mixed coniferous/deciduous forests.	Feeds primarily on small mammals during the summer, shifts to more birds during the winter. L1: Wilson Bulletin 42:198. L3: J. McGough, 6 March 1978. L8: A. Pinter, 1979.	<u>h</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>?</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u>
Northern Pygmy-Owl <i>Glaucidium gnoma</i> 379.0 NOPO Resident Rare Life Form 14	(FL)	NG NSSU	Lodgepole pine, Douglas fir, Englemann spruce-subalpine fir, other coniferous forests; also aspen.	Nests in a cavity in a snag. Feeds on small mammals, insects, birds, some invertebrates.	<u>B</u> <u>O*</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>B</u> <u>B</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>B</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u> <u>—</u>
Burrowing Owl <i>Athene cunicularia</i> 378.0 BUOW Summer resident Uncommon Life Form 15		NG NSSU NTMB	Grasslands, basin-prairie shrublands, agricultural areas.	Nests in a mammal burrow, especially that of a prairie dog. Feeds on insects, rodents, lizards, birds.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Barred Owl <i>Strix varia</i> 368.0 BDOW Accidental Rare Life Form	(AS)		Dense coniferous and mixed coniferous/deciduous forests.	Feeds primarily on small rodents; also birds, other small vertebrates. L7: McCreary (1939). L8: first accepted record, E. Bowman, 21 September 1982; first suspected breeding record, S. Derousseau, 31 May 2009. L14: D. Shorma, 10-11 March 2006.	— — — — — — — — — — b* —
Great Gray Owl <i>Strix nebulosa</i> 370.0 GGOW Resident Rare Life Form 12		NG NSSU	Lodgepole pine, Douglas fir, Englemann spruce-subalpine fir, other coniferous forests, aspen, mountain-foothills grasslands.	Nests in an abandoned hawk or eagle nest. Feeds primarily on voles; will also take birds, other small mammals.	B O* — — — — — — — — — — B O* — — — — — — — — — — B b B* — — — — — — — — — — O O — — — — — — — — — —
Long-eared Owl <i>Asio otus</i> 366.0 LEOW Resident Common Life Form 7 11		NG NTMB	Most habitats below 8,000 feet, especially cottonwood-riparian.	Nests in an abandoned crow, hawk, or squirrel nest. Feeds on small rodents, some birds.	B O B B B O O B O B O O B B B O O B b O B B B B B O B O
Short-eared Owl <i>Asio flammeus</i> 367.0 SEOW Resident Common Life Form 5		NG NSS4 NTMB	Basin-prairie shrublands, grasslands, marshes, irrigated native meadows, below 7,000 feet.	Nests on the ground, often concealed by low vegetation. Feeds primarily on rodents; will also take small birds, insects.	B O O B B B O B O B O O B b B b O O B O O b B b B O B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Boreal Owl <i>Aegolius funereus</i> 371.0 BOOW Resident Uncommon Life Form 14	(FL)	NG NSS3	Mature coniferous and mixed coniferous/deciduous forests with scattered openings, especially spruce-fir, lodgepole pine, and aspen.	Nests in an existing cavity in a tree. Feeds on small mammals, birds. L21: winter observation (McCreary 1939).	b* O* _ _ B* _ _ _ B O* _ _ _ _ _ O* _ O* _ _ _ h O _ _ O B* _ _
Northern Saw-whet Owl <i>Aegolius acadicus</i> 372.0 NSWO Resident Uncommon Life Form 14	(FL)	NG	Coniferous and mixed aspen/coniferous forests, aspen, cottonwood-riparian.	Nests in an existing cavity in a tree. Feeds on small mammals, birds.	B O* O* O B b O* B B O* _ O O O B O B O O b O b b* _ O* O B O
Belted Kingfisher <i>Megasceryle alcyon</i> 390.0 BEKI Resident Common Life Form 16		NG NTMB	Aquatic areas, cottonwood-riparian.	Nests in a horizontal burrow in a bank near water. Feeds primarily on fish, occasionally aquatic invertebrates, amphibians, reptiles, insects.	B B B B B B B B B B b B O B B B B b b B b B B B B b B
Lewis's Woodpecker <i>Melanerpes lewis</i> 408.0 LEWO Summer resident Uncommon Life Form 13		NG NSSU NTMB	Ponderosa pine savannah, pine-juniper, other coniferous forests, aspen, cottonwood-riparian, below 8,500 feet.	Nests in a cavity in a dead or live tree or in a pole. Feeds on insects, nuts, berries. L10: W. Major, 8 May 2013.	B B B B B O B B O O b O O B B b O _ B B B b b O O B B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Red-headed Woodpecker <i>Melanerpes erythrocephalus</i> 406.0 RHWO Summer resident Uncommon Life Form 13	(FL)	NG NSS4	Cottonwood-riparian, ponderosa pine savannah, aspen.	Nests in a cavity in a barkless dead tree or a stub on a live tree. Feeds on a variety of plant and animal material including insects, bird eggs, nestlings, mice, corn, berries, seeds.	<u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B*</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u> </u> <u>O*</u> <u> </u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>O*</u> <u>O</u> <u> </u> <u>O</u> <u>O*</u> <u>B</u>
Acorn Woodpecker <i>Melanerpes formicivorus</i> 407.0 ACWO Accidental Rare Life Form	(AS)		Oak and mixed oak/coniferous forests.	Feeds mostly on insects; also acorns, fruit, tree sap. L8: T. Tempest and C. McGinley, 6 June 1975; T. Williams, 14 June 1975. L24: F. Luke, 15 July 1989.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u>
Red-bellied Woodpecker <i>Melanerpes carolinus</i> 409.0 RBWO Accidental Rare Life Form	(AS)		Deciduous forests, also coniferous forests.	Feeds on insects, nuts, fruit, seeds. L20: D. Lutz and J. Bowen, 14 January 1993. L21: S. Dinsmore, 13 December 2002. L28: J. Cairo, 7 May 1992; D. Faulkner, 21-26 August 2006.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u>O*</u>
Williamson's Sapsucker <i>Sphyrapicus thyroideus</i> 404.0 WISA Summer resident Uncommon Life Form 13		NG NSS3 NTMB	Coniferous forests, especially those that have burned. Also aspen.	Nests in a cavity in an aspen, pine, or fir. Feeds on insects, tree sap.	<u>B</u> <u>O</u> <u> </u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>b</u> <u>O</u> <u>B</u> <u> </u> <u> </u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
White-headed Woodpecker <i>Picoides albolarvatus</i> 399.0 WHWO Accidental Rare Life Form	(AS)		Coniferous forests from 4,000 to 9,000 feet.	Feeds on insects, conifer seeds. L8: McCreary (1939). Several reports since 1952 lack details, including L1 where D. Tessen has four records, 1952-1971.	O _ _ _ _ _ h _ _ _ _ _ _ _ _ _ _ _ _ _ _ _
American Three-toed Woodpecker <i>Picoides dorsalis</i> 401.0 ATTW Resident Uncommon Life Form 13		NG	Lodgepole pine, Douglas fir, Englemann spruce-subalpine fir, especially those forests that have burned.	Nests in a cavity in a coniferous or deciduous snag, especially near a burn. Feeds on insects, especially wood boring beetles, tree sap.	B O O B O _ _ B B O B _ _ O B b b O b O _ b B O B O B _
Black-backed Woodpecker <i>Picoides arcticus</i> 400.0 BBWO Resident Rare Life Form 13	(FL)	NG NSSU	Lodgepole pine, Douglas fir, Englemann spruce-subalpine fir, especially those forests that have burned.	Nests in a cavity in a conifer. Feeds on insects, mostly larvae of wood boring insects. L7: S. Ritter and T. Ryder, 9 June 1987. L8: R. Wallen, 31 May 1993; K. Duffy and P. Matheny, 12 July 1993; R. Steenberg and others, 6 July 1996. L17: R. Scott and B. Scott, 19 July 1993.	B _ _ _ _ O B* B* _ _ _ _ O B O _ O* O _ _ _ _ _ _ _
Northern Flicker <i>Colaptes auratus</i> 412.2 NOFL Resident Common Life Form 13		NG	Most habitats as long as trees or poles are present, especially deciduous forests.	Nests in a cavity, preferably in a snag, but will use a pole, post, house, bank, or haystack. Feeds on insects, especially ants. Includes Red-shafted and Yellow-shafted races.	B B B B B B B B B B b B B B B B B B B B B B B B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Pileated Woodpecker <i>Dryocopus pileatus</i> 405.0 PIWO Accidental Rare Life Form	(AS)		Deciduous and coniferous forests, open woodlands.	Feeds primarily on insects, some fruit, acorns, nuts, tree sap. L1: B. Wood, December 1961 and 1963. L2: J. Holmes and B. Holmes, 21 February 2000. L8: C. Schrier, 23 April 1980; V. Herold, 7 July 1995. L17: P. Halstead, 25 May 2006.	B+ O* O* O* — — — — — — — — — — — — — — — —
Crested Caracara <i>Caracara cheriway</i> 362.0 CRCA Accidental Rare Life Form	(AS)		Basin-prairie shrublands, sagebrush-grasslands.	Feeds mostly on carrion. L1: J. Peaco, 8-12 September 1984. May have been a released bird.	O* —
American Kestrel <i>Falco sparverius</i> 360.0 AMKE Summer resident Common Life Form 4 14		F NSS4 NTMB	All habitats, especially below 8,500 feet.	Nests in a tree cavity, hole in a cliff, building, or magpie nest. Feeds on insects, small vertebrates, small mammals, birds. Some individuals remain throughout the winter.	B B
Merlin <i>Falco columbarius</i> 357.0 MERL Resident Uncommon Life Form 11		F NSSU NTMB	Most habitats below 8,500 feet.	Nests in an old nest, usually magpie, in a coniferous or deciduous tree. Feeds mostly on birds; also small mammals, insects.	O B O B B B B b O B O B B O B B B O B B O B B b O — B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b																												
Gyrfalcon <i>Falco rusticolus</i> 354.0 GYRF Winter resident Rare Life Form	(AS)	F	Basin-prairie shrublands, grasslands, agricultural areas.	Feeds mostly on birds; also small mammals.	<table border="0"> <tr><td>—</td><td>—</td><td>O</td><td>O*</td><td>O*</td><td>O</td><td>O</td></tr> <tr><td>O</td><td>—</td><td>O</td><td>—</td><td>O*</td><td>O</td><td>—</td></tr> <tr><td>—</td><td>O</td><td>O*</td><td>—</td><td>O*</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>O</td><td>O</td><td>—</td></tr> </table>	—	—	O	O*	O*	O	O	O	—	O	—	O*	O	—	—	O	O*	—	O*	—	—	—	—	—	—	O	O	—
—	—	O	O*	O*	O	O																											
O	—	O	—	O*	O	—																											
—	O	O*	—	O*	—	—																											
—	—	—	—	O	O	—																											
Peregrine Falcon <i>Falco peregrinus</i> 356.0 PEFA Resident Uncommon Life Form 4	(FL)	NG NSS3 NTMB	Cliffs in most habitats.	Nests on a ledge or in a hole on a tall cliff. Feeds on birds. Removed from the federal Endangered Species List in 1999.	<table border="0"> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>O</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>O*</td><td>O</td><td>O</td><td>—</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>O</td><td>B*</td><td>O*</td><td>H</td></tr> <tr><td>O</td><td>H</td><td>H</td><td>O</td><td>O</td><td>H</td><td>O</td></tr> </table>	B	B	B	B	B	O	B	B	B	B	O*	O	O	—	B	B	B	O	B*	O*	H	O	H	H	O	O	H	O
B	B	B	B	B	O	B																											
B	B	B	O*	O	O	—																											
B	B	B	O	B*	O*	H																											
O	H	H	O	O	H	O																											
Prairie Falcon <i>Falco mexicanus</i> 355.0 PRFA Resident Common Life Form 4		F NTMB	Cliffs in all habitats with open areas.	Nests in a hole or on a ledge on a cliff or rock outcrop. Feeds on birds, small mammals, insects, lizards.	<table border="0"> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> </table>	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B	B
B	B	B	B	B	B	B																											
B	B	B	B	B	B	B																											
B	B	B	B	B	B	B																											
B	B	B	B	B	B	B																											
Olive-sided Flycatcher <i>Contopus cooperi</i> 459.0 OSFL Summer resident Common Life Form 10		NG NTMB	Coniferous forests from 8,000 feet to timberline, aspen-riparian.	Nests often high in a conifer on a horizontal branch. Feeds exclusively on insects that can be caught in the air.	<table border="0"> <tr><td>B</td><td>b</td><td>O</td><td>B</td><td>B</td><td>—</td><td>O</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>—</td><td>B</td><td>—</td><td>—</td></tr> <tr><td>B</td><td>b</td><td>b</td><td>O</td><td>b</td><td>b</td><td>—</td></tr> <tr><td>b</td><td>b</td><td>b</td><td>B</td><td>B</td><td>B</td><td>O</td></tr> </table>	B	b	O	B	B	—	O	B	B	B	—	B	—	—	B	b	b	O	b	b	—	b	b	b	B	B	B	O
B	b	O	B	B	—	O																											
B	B	B	—	B	—	—																											
B	b	b	O	b	b	—																											
b	b	b	B	B	B	O																											

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Western Wood-Pewee <i>Contopus sordidulus</i> 462.0 WEWP Summer resident Common Life Form 11		NG NTMB	All habitats with trees.	Nests usually on a horizontal limb far from the trunk. Feeds almost entirely on insects; also a few berries.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Eastern Wood-Pewee <i>Contopus virens</i> 461.0 EAWP Accidental Rare Life Form	(AS)	NTMB	Deciduous and mixed deciduous/coniferous forests.	Feeds almost entirely on insects; also a few berries. L19: O.K. Scott, 9 July 1975. L10: S. Mlodinow, 28 June 2014.	-- -- -- -- -- -- -- -- -- O* -- -- -- -- -- -- -- -- O* -- -- -- -- -- -- -- -- --
Willow Flycatcher <i>Empidonax traillii</i> 466.0 WIFL Summer resident Common Life Form 8		NG NSS3 NTMB	Riparian shrub including willow, hawthorn, water birch, alder; below 9,000 feet.	Nests in an upright or slanting fork in a shrub. Feeds primarily on insects, occasionally berries.	<u>B</u> <u>B</u> <u> </u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u> </u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u>
Least Flycatcher <i>Empidonax minimus</i> 467.0 LEFL Summer resident Common Life Form 8 11	(FL)	NG NTMB	Deciduous forests with a variety of open habitats.	Nests on a horizontal limb of a tree, occasionally in a conifer. Feeds on insects, occasionally berries, seeds. L8: G. Rankin and W. Argabrite, 25 June 1994. L26: CJ Grimes, 13 June 2014.	<u> </u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O*</u> <u>b</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u> </u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>b</u> <u>O*</u> <u> </u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Hammond's Flycatcher <i>Empidonax hammondi</i> 468.0 HAFL Summer resident Uncommon Life Form 10	(FL)	NG NTMB	Douglas fir, Englemann spruce-subalpine fir, and other coniferous forests with aspen; willow riparian; above 6,500 feet.	Nests on a horizontal limb of a tall conifer, occasionally in a deciduous tree. Feeds exclusively on insects. Difficult to distinguish from Dusky Flycatcher; habitat differences are not always reliable. L4: G. Nutting, 28 June 2001. L8: S. Fitton, 19 June 1989; P. Dickson and D. Wile, 6 June and 5 July 1995.	<u>B</u> <u>O</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>B*</u> <u>B</u> <u>B</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u> </u> <u>B</u> <u>O</u> <u>O</u> <u>O</u>
Gray Flycatcher <i>Empidonax wrightii</i> 469.1 GRFL Summer resident Common Life Form 8	(FL)	NG NTMB	Pine-juniper, woodland-chaparral, basin-prairie and mountain-foothills shrublands.	Nests in the crotch of a juniper or sagebrush, or near the base of a thorny shrub. Feeds exclusively on insects. L4: E. Stone, 10 June 1998; B. Righter, 27 June 1998.	<u> </u> <u> </u> <u> </u> <u>b*</u> <u> </u> <u> </u> <u>?</u> <u>O</u> <u> </u> <u>b</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u> </u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u> </u> <u> </u> <u> </u>
Dusky Flycatcher <i>Empidonax oberholseri</i> 469.0 DUFL Summer resident Common Life Form 8		NG NTMB	Ponderosa pine savannah, pine-juniper, aspen, cottonwood-riparian, woodland-chaparral, riparian shrub.	Nests in the crotch of a juniper or sagebrush, or near the base of a thorny shrub. Feeds exclusively on insects. Difficult to distinguish from Hammond's Flycatcher. Habitat differences: Dusky prefers dry, open forests; Hammond's prefers wet, mountain forests.	<u>B</u> <u>b</u> <u> </u> <u>B</u> <u>B</u> <u> </u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u>
Cordilleran Flycatcher <i>Empidonax occidentalis</i> 464.0 COFL Summer resident Common Life Form 4 17		NG NTMB	Moist areas of coniferous forests, aspen-riparian, aspen-conifer.	Nests in a wide variety of situations from streambank to cave, cliff ledge, or cavity in a small tree. Feeds almost entirely on insects; also some berries, seeds.	<u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Eastern Phoebe <i>Sayornis phoebe</i> 456.0 EAPH Summer resident Peripheral Abundance unknown Life Form 6 17	(AS)	NG NTMB	Cottonwood-riparian, grasslands, agricultural areas, urban areas.	Nested originally in a cliff or bank, now mostly in a human-built structure. Feeds primarily on insects, occasionally small fish, frogs, berries, seeds. Nesting may occur elsewhere in eastern Wyoming.	— — — O* O — B* — — — O* O — — O — — — — O — O — — — — — O* O
Say's Phoebe <i>Sayornis saya</i> 457.0 SAPH Summer resident Common Life Form 4 17		NG NTMB	Basin-prairie shrublands, grasslands.	Nests in a cliff or bank, occasionally under an eave or bridge. Feeds almost exclusively on insects, occasionally some berries.	O O B B B B B b B b B B B B B B B B b B b B B B B B B
Vermilion Flycatcher <i>Pyrocephalus rubinus</i> 471.0 VEFL Accidental Rare Life Form	(AS)	NTMB	Riparian shrub, basin-prairie shrublands.	Feeds almost exclusively on insects. L8: S. MacDonald, 5 July 1986. L15: R. Steenberg, 22 May 1993. L23: P. Spivak, 31 May 1995.	— — — — — — O* — — — — — O* — — — — — — — O* — — — —
Ash-throated Flycatcher <i>Myiarchus cinerascens</i> 454.0 ATFL Summer resident Peripheral Abundance unknown Life Form 14	(FL)	NG NSS3 NTMB	Pine-juniper, juniper, basin-prairie shrublands.	Nests in a natural cavity, old woodpecker cavity, or hole in a fence post. Feeds mostly on insects, occasionally small fruit. L23: R. Steenberg, 29 June 1997. L24: includes cavity repair by adults; fledged young (S. Fitton and F. Broerman).	O — — — O* — — — O* O O — — — O — — — — O* — O — O* O B* B b O — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Great Crested Flycatcher <i>Myiarchus crinitus</i> 452.0 GCFL Accidental Rare Life Form	(AS)	NTMB	Deciduous forest edge.	Feeds primarily on insects, occasionally small lizards, some fruit. L20: McCreary (1939). L21: V. Herold and others, 8 June 1995. L28: Don Jones and others, 3 September 2012.	— <u>O</u> — — — — — <u>O</u> — — — — — — — — — — h <u>O*</u> — — — — — <u>O*</u>
Cassin's Kingbird <i>Tyrannus vociferans</i> 448.0 CAKI Summer resident Uncommon Life Form 11	(FL)	NG NTMB	Ponderosa pine savannah, pine-juniper, cottonwood-riparian, cottonwood-dryland, woodland-chaparral, basin-prairie and mountain-foothills shrublands.	Nests on a horizontal branch near the trunk of a tree. Feeds on insects, berries. L11: J. Allen, 12 May 1992. L19: L. Rognstad, 30 July 1992. L26: H. Haley, 26 May 2000.	<u>O</u> — — — — <u>O</u> b b — — — — <u>O</u> <u>O*</u> — — B — — — — B <u>O</u> <u>O*</u> <u>O</u> B — — — — <u>O</u> — — <u>O*</u> B b
Western Kingbird <i>Tyrannus verticalis</i> 447.0 WEKI Summer resident Common Life Form 11 17		NG NTMB	Basin-prairie shrublands, grasslands, agricultural areas, deciduous forest edge, riparian shrub.	Nests on a horizontal branch near or against the trunk of a tree. Feeds on insects, berries.	<u>B</u> b <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> b <u>B</u> b <u>B</u> b <u>B</u> b <u>B</u> b <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> b <u>B</u> <u>B</u>
Eastern Kingbird <i>Tyrannus tyrannus</i> 444.0 EAKI Summer resident Common Life Form 8 11 17		NG NTMB	Basin-prairie shrublands, grasslands, agricultural areas, deciduous forest edge, riparian shrub.	Nests on a horizontal limb of an isolated tree, usually at mid-height, occasionally on a fence post or stump. Feeds primarily on insects, some fruit.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> b <u>B</u> b <u>B</u> b <u>B</u> <u>B</u> <u>B</u> b <u>B</u> <u>O</u> <u>B</u> <u>O</u> b b <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Scissor-tailed Flycatcher <i>Tyrannus forficatus</i> 443.0 STFL Accidental Rare Life Form	(AS)	NTMB	Basin-prairie shrublands.	Feeds almost entirely on insects, occasionally a few berries.	<u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u>
Loggerhead Shrike <i>Lanius ludovicianus</i> 622.0 LOSH Summer resident Common Life Form 7		NG NSS4 NTMB	Open habitats with short vegetation, including hay fields, pastures, mowed roadsides, open woodlands, sagebrush. Woodland-chapparral, basin-prairie and mountain-foothills shrublands.	Nest is usually hidden below the crown in the crotch or on a large branch of a deciduous tree or shrub. Feeds on insects, small vertebrates, carrion.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u>
Northern Shrike <i>Lanius excubitor</i> 621.0 NSHR Winter resident Common Life Form		NG	Most open habitats below 7,000 feet, especially roadsides.	Feeds on small mammals, insects.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
White-eyed Vireo <i>Vireo griseus</i> 631.0 WEVI Accidental Rare Life Form	(AS)	NTMB	Riparian shrub, thickets, undergrowth.	Feeds primarily on insects, also berries. L27: D. Young, 16 November 1994. L28: J. Eddy, 17 May 2004.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Gray Vireo <i>Vireo vicinior</i> 634.0 GRVI Summer resident Rare Life Form 7	(AS)	NG NSSU	Pine-juniper, woodland-chaparral.	Nests in a shrub; nest rim is suspended from a forked twig. Feeds on insects. At the northern extent of its range. L23: D. McDonald, 29 May 2006. L24: C. Dexter, 16-19 June 2005.	— — — — — — — — — — — — — — — — O* O* — — — — —
Yellow-throated Vireo <i>Vireo flavifrons</i> 628.0 YTVI Accidental Rare Life Form	(AS)	NTMB	Deciduous and mixed deciduous/coniferous forests.	Feeds on insects.	— — — — — — O* — — — — — — O* — — — — — — — — — —
Cassin's Vireo <i>Vireo cassinii</i> 629.2 CAVI Migrant Rare Life Form	(AS)	NG NTMB	Deciduous and mixed deciduous/coniferous forests.	Feeds almost entirely on insects.	— — — — — — O* O* — — — — — — — — — — O* — O* — — — — —
Blue-headed Vireo <i>Vireo solitarius</i> 629.0 BHVI Migrant Rare Life Form	(AS)	NG NTMB	Deciduous and mixed deciduous/coniferous forests.	Feeds almost entirely on insects. L23: R. Steenberg and J. Steenberg, 15 September 1999.	— — — — — — — — — — — — — — — — O* — — — — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Plumbeous Vireo <i>Vireo plumbeus</i> 629.1 PLVI Summer resident Uncommon Life Form 11		NG NTMB	Douglas fir, ponderosa pine savannah, ponderosa pine-Douglas fir, other or mixed coniferous forests, aspen, cottonwood-riparian.	Nests in a conifer, occasionally in an oak. Feeds almost entirely on insects. Locally common.	<u>O</u> <u>b</u> <u> </u> <u>B</u> <u>B</u> <u> </u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u>
Philadelphia Vireo <i>Vireo philadelphicus</i> 626.0 PHVI Migrant Rare Life Form	(AS)	NG NTMB	Deciduous forests, urban areas.	Feeds mostly on insects, occasionally berries. L4, 5: H. Downing has compiled numerous observations. L19: L. Rognstad, 19 September 1982; G. Lawrence, 13 September 1992. L28: D. Faulkner, 20 May 2006.	<u> </u> <u> </u> <u> </u> <u>O</u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>
Warbling Vireo <i>Vireo gilvus</i> 627.0 WAVI Summer resident Common Life Form 11		NG NTMB	Deciduous and coniferous forests, urban areas.	Nest is usually high in a deciduous tree, occasionally low in a tree or shrub. Feeds mostly on insects, occasionally berries.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Red-eyed Vireo <i>Vireo olivaceus</i> 624.0 REVI Summer resident Uncommon Life Form 11		NG NSS4 NTMB	Deciduous and coniferous forests, urban areas.	Nests in a deciduous tree or shrub. Feeds mainly on insects, occasionally fruit.	<u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u> </u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>b</u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FI) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Gray Jay <i>Perisoreus canadensis</i> 484.0 GRAJ Resident Common Life Form 10		NG	Coniferous forests.	Nests on a horizontal branch near the trunk or in the crotch, usually in a conifer. Feeds primarily on insects, fruit, carrion.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u> </u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u>
Pinyon Jay <i>Gymnorhinus cyanocephalus</i> 492.0 PIJA Resident Uncommon Life Form 11		NG	Ponderosa pine savannah, pine-juniper, woodland-chaparral, mountain-foothills shrublands.	Nests in a juniper or pine, occasionally an oak. Feeds on conifer seeds, fruit, insects, eggs, nestlings. Locally common.	<u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>O</u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u>
Steller's Jay <i>Cyanocitta stelleri</i> 478.0 STJA Resident Common Life Form 11		NG	Coniferous forests, juniper-mountain mahogany, mountain mahogany, urban areas.	Nests on a horizontal branch or in the crotch of a conifer. Feeds on insects, invertebrates, eggs, nestlings, seeds, fruit.	<u>B</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u>
Blue Jay <i>Cyanocitta cristata</i> 477.0 BLJA Resident Uncommon Life Form 11		NG	Ponderosa pine-Douglas fir, other or mixed coniferous forests with cottonwood-riparian, riparian shrub, Gambel oak.	Nests on a horizontal branch or in the crotch of a conifer, occasionally on a deciduous tree. Feeds on insects, invertebrates, eggs, nestlings, carrion, seeds, nuts, fruit. Common in the Black Hills and Cheyenne; uncommon elsewhere, but is expanding its range.	<u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>B</u> <u>O</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Woodhouse's Scrub-Jay <i>Aphelocoma woodhouseii</i> 481.0 WOSJ Resident Peripheral Uncommon Life Form 11	(FL)	NG NSS3	Pine-juniper, woodland-chaparral.	Nests usually in a small conifer. Feeds on insects, invertebrates, small vertebrates, eggs, nestlings, nuts, seeds, fruit. L21: K. Kranik, 27 October 2002. L27: H. Whitman and V. Whitman, 13 September 1993. L28: D. Adams, 7 January 2008.	— — — — O — — O* — — — — — — — — — — — O — — O* O B B b O O* O*
Clark's Nutcracker <i>Nucifraga columbiana</i> 491.0 CLNU Resident Common Life Form 10		NG NSS4	Coniferous forests, aspen, cliffs in canyons or mountains, juniper-sagebrush, ponderosa pine-juniper.	Nests on a horizontal limb of a mature conifer. Feeds on conifer seeds, fruit, insects, small vertebrates, eggs, nestlings.	B B B B B O b B B b B B O O B B B B B B O B b b b B B B
Black-billed Magpie <i>Pica hudsonia</i> 475.0 BBMA Resident Abundant Life Form 7 11		NG	All habitats below 8,000 feet.	Nest is large and conspicuous in a small tree or shrub. Feeds on insects, carrion, invertebrates, small vertebrates, fruit, seeds.	B B B B B B B B B B B B B b B B B B B B B B B B B B B B
American Crow <i>Corvus brachyrhynchos</i> 488.0 AMCR Resident Common Life Form 11		NG	Most habitats below 9,000 feet.	Nests in a conifer or a deciduous tree or shrub. Feeds on insects, invertebrates, small vertebrates, carrion, eggs, nestlings, seeds, fruit.	B b B O B O b B O B b b O b b b B B b b B B O O B B B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Common Raven <i>Corvus corax</i> 486.0 CORA Resident Common Life Form 4 11		NG	All habitats.	Nests on a cliff or a human-built structure, occasionally in a deciduous tree. Feeds primarily on carrion; also small vertebrates, eggs, nestlings, insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u>
Horned Lark <i>Eremophila alpestris</i> 474.0 HOLA Resident Abundant Life Form 5		NG NTMB	Basin-prairie and mountain-foothills shrublands, grasslands including alpine.	Nests on the ground in a shallow depression. Feeds on seeds, insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Purple Martin <i>Progne subis</i> 611.0 PUMA Summer resident Rare Life Form 14 17	(AS)	NG NSSU NTMB	Habitats near aquatic areas below 7,000 feet, especially in urban areas.	Nests in a tree cavity, cliff, niche, or other cavity, often in a bird house. Feeds on insects, occasionally feeds on the ground taking ants and other insects. L11: H. Downing, 7 August 1978. L25: D. Faulkner and R. Levad, 21 July 2004; L25: CJ Grimes, 21 June 2014.	<u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>H</u> <u> </u> <u>O</u> <u> </u> <u>B*</u> <u>h</u> <u>O</u> <u>O</u>
Tree Swallow <i>Tachycineta bicolor</i> 614.0 TRES Summer resident Common Life Form 14 17		NG NTMB	Aerial feeder over most habitats below 8,500 feet, especially aspen and cottonwood-riparian; also lodgepole pine, Douglas fir, ponderosa pine savannah, ponderosa pine-Douglas fir.	Nests in a tree cavity or other cavity. Feeds on insects, occasionally berries.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Violet-green Swallow <i>Tachycineta thalassina</i> 615.0 VGSW Summer resident Common Life Form 4 14 17		NG NTMB	Aerial feeder over most habitats within aquatic areas below 8,500 feet.	Nests in a tree cavity or other cavity. Feeds entirely on insects.	<u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Northern Rough-winged Swallow <i>Stelgidopteryx serripennis</i> 617.0 NRWS Summer resident Common Life Form 16		NG NTMB	Adjacent to aquatic areas. Forages over a variety of habitats below 8,000 feet.	Nests at the end of a burrow in a cliff, or other cavity or niche. Feeds entirely on insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Bank Swallow <i>Riparia riparia</i> 616.0 BANS Summer resident Common Life Form 16		NG NTMB	Adjacent to aquatic areas. Forages over a variety of habitats below 8,000 feet.	Nests at the end of a burrow in a bank or cliff. Feeds almost entirely on insects.	<u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u>
Cliff Swallow <i>Petrochelidon pyrrhonota</i> 612.0 CLSW Summer resident Common Life Form 4 17		NG NTMB	All habitats with aquatic areas below 9,000 feet.	Nest is plastered on the underside of a bridge or culvert, on a cliff, or on a wall under an eave. Feeds almost entirely on insects, occasionally gorges on berries.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Barn Swallow <i>Hirundo rustica</i> 613.0 BARS Summer resident Common Life Form 4 17		NG NTMB	All habitats with aquatic areas below 8,000 feet.	Nest is usually plastered on a ledge or a wall of a building, occasionally in a cave or under a bridge. Feeds primarily on insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Black-capped Chickadee <i>Poecile atricapillus</i> 735.0 BCCH Resident Common Life Form 14		NG	Coniferous and deciduous forests, riparian shrub, urban areas, mostly below 8,500 feet.	Nests in a natural or woodpecker cavity in a tree snag, rarely in a conifer. Feeds on insects, conifer seeds, fruit.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Mountain Chickadee <i>Poecile gambeli</i> 738.0 MOCH Resident Common Life Form 14		NG	Coniferous forests, aspen, juniper-sagebrush. Common up to timberline, lower habitats during the winter.	Nests in a natural or woodpecker cavity in a tree or snag. Feeds on insects, conifer seeds, spiders and their eggs.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Juniper Titmouse <i>Baeolophus ridgwayi</i> 733.1 JUTI Resident Uncommon Life Form 14	(FL)	NG NSS3	Pine-juniper, woodland-chaparral, mountain-foothills shrublands, juniper-sagebrush, urban areas.	Nests in a natural or woodpecker cavity in a tree or snag. Feeds on insects, fruit, seeds.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Bushtit <i>Psaltriparus minimus</i> 743.0 BUSH Summer resident Peripheral Uncommon Life Form 8	(FL)	NG NSS3	Pine-juniper, woodland-chaparral, juniper-sagebrush, mountain-foothills shrublands, riparian shrub.	Nest is a gourd-shaped hanging pocket in a small tree or shrub. Feeds on insects, seeds, fruit.	— — — — O — — — — — — — — — — — O — O* — — O B b — O O O
Red-breasted Nuthatch <i>Sitta canadensis</i> 728.0 RBNU Resident Common Life Form 13		NG	Coniferous forests, aspen, cottonwood-riparian. A variety of lowland habitats during the winter.	Nests in a cavity excavated in a rotten branch or stump, occasionally in a deserted woodpecker cavity. Feeds primarily on insects; takes many conifer seeds in the winter.	B B O B B O B B B B O O B b b b B b B b b b b B B B B b
White-breasted Nuthatch <i>Sitta carolinensis</i> 727.0 WBNU Resident Common Life Form 13		NG	Coniferous forests, aspen, cottonwood-riparian. A variety of lowland habitats during the winter.	Nests in a natural cavity or deserted woodpecker cavity in a tree or snag. Feeds primarily on insects.	B B — B B O B B B B O O O B b O B — B b b O b O O B b b
Pygmy Nuthatch <i>Sitta pygmaea</i> 730.0 PYNU Resident Uncommon Life Form 13		NG NSS3	Coniferous forests, ponderosa pine. Lower habitats during the winter.	Nest is often excavated in a pine or post, occasionally uses a deserted woodpecker cavity. Feeds on insects, conifer seeds.	O# O — B O O b O O O O O — — b — — b B B O b b — — O B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Brown Creeper <i>Certhia americana</i> 726.0 BRCR Resident Common Life Form 14		NG NTMB	Coniferous forests. Lower habitats during the winter.	Nest is a hammock-like cup, usually beneath loose bark, rarely in a cavity. Feeds primarily on insects, some nuts, seeds.	<u>B</u> <u>B</u> <u> </u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u> </u> <u>B</u> <u>b</u> <u>B</u> <u>O</u>
Rock Wren <i>Salpinctes obsoletus</i> 715.0 ROWR Summer resident Common Life Form 4		NG NTMB	Rock outcrops/rock piles in pine-juniper, woodland-chaparral, basin-prairie and mountain-foothills shrublands, grasslands.	Nests in a hole or crevice, often under or around rocks. Feeds primarily on insects.	<u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Canyon Wren <i>Catherpes mexicanus</i> 717.0 CANW Resident Uncommon Life Form 4		NG NSS4	Cliffs, canyons, rock outcrops, and rock piles in grasslands, woodland-chaparral, pine-juniper, oak, and coniferous forests. Often found near water.	Nests in a crevice or cave on a bank or cliff. Feeds mostly on insects.	<u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u> </u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u> </u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u> </u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u>
House Wren <i>Troglodytes aedon</i> 721.0 HOWR Summer resident Common Life Form 14		NG NTMB	Aspen, cottonwood-riparian, other deciduous forests.	Nests in a natural cavity, occasionally in the nest of another bird. Feeds on insects, invertebrates.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Pacific Wren <i>Troglodytes pacificus</i> 722.1 PAWR Migrant Rare Life Form	(AS)	NG	Coniferous forests.	Feeds on insects, invertebrates, spiders. AOU split Winter Wren and Pacific Wren in 2011. May breed in northwestern Wyoming. L8: singing male, S. Findholt, 24 June 1981.	O* O* _ O* O* _ O* b* O* O* _ _ _ _ b* _ _ _ _ _ _ _ _ _ _ _ _
Winter Wren <i>Troglodytes hiemalis</i> 722.0 WIWR Migrant Rare Life Form	(AS)	NG	Coniferous forests.	Feeds mostly on insects; rarely juniper berries. L19: McCreary (1939) and recent observations during migration. AOU split Winter Wren and Pacific Wren in 2011.	_ _ _ _ O* _ _ _ _ _ _ _ _ _ _ _ _ _ _ H O* _ _ _ _ _ _ O O _ _
Sedge Wren <i>Cistothorus platensis</i> 724.0 SEWR Accidental Rare Life Form	(AS)	NTMB	Wet-moist meadow grasslands, sedge marshes.	Feeds primarily on insects. L18: J. Ward, 16-17 October 1994. L27: D. Finch, 1 June 1982. L28: collected by F. Bond, 14 April 1889 (McCreary 1939).	_ O* _ _ _ _ _ _ _ _ _ _ O* h
Marsh Wren <i>Cistothorus palustris</i> 725.0 MAWR Summer resident Common Life Form 3		NG NTMB	Marshes. Drier habitats during migration.	Nest is attached to reeds. Feeds on insects, snails. Abundant in some areas.	B O B O O O _ B O B B B _ _ B B B B O O O B B B b B B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Carolina Wren <i>Thryothorus ludovicianus</i> 718.0 CARW Accidental Rare Life Form	(AS)		Deciduous forests with heavy undergrowth.	Feeds on insects, invertebrates, small vertebrates, some seeds. L19: O.K. Scott, 8 May 1977.	— O* — — — — — — — — — — — —
Bewick's Wren <i>Thryomanes bewickii</i> 719.0 BEWR Summer resident Peripheral Uncommon Life Form 14	(FL)	NG NSS4	Pine-juniper, woodland-chaparral, mountain-foothills shrublands.	Nests in a natural cavity in a tree; also amid roots of an upturned tree. Feeds primarily on insects.	— — — — — O — — — — — — O — — — — O — — — — — — — — — O O — — — — b B B B B O O
Blue-gray Gnatcatcher <i>Poliioptila caerulea</i> 751.0 BGGN Summer resident Uncommon Life Form 11		NG NSS4 NTMB	Pine-juniper, cottonwood-riparian, juniper, mountain-foothills shrublands.	Nest is a saddle on a horizontal limb or in a fork, usually in a deciduous tree. Feeds on insects.	O# O — — — — O — — — — O O b O — — — — O — — O O B O B O B B b O O O
American Dipper <i>Cinclus mexicanus</i> 701.0 AMDI Resident Common Life Form 3		NG	Swift mountain streams.	Nests on a cliff face, behind a waterfall, or on a midstream rock. Feeds on aquatic invertebrates, fish, small aquatic plants.	B B O B B — — — — B B O O B — — — — B B B — — B B B O b — — b B B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Golden-crowned Kinglet <i>Regulus satrapa</i> 748.0 GCKI Resident Uncommon Life Form 10		NG NTMB	Coniferous forests, aspen-conifer.	Nest is hung from branches near the trunk of a conifer. Feeds on insects, tree sap, fruit, some seeds.	<u>B</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u>
Ruby-crowned Kinglet <i>Regulus calendula</i> 749.0 RCKI Summer resident Common Life Form 10		NG NTMB	Coniferous forests, juniper. Lower habitats during migration, including urban areas.	Nest is hung from a limb of a conifer. Feeds on insects, tree sap, berries, some seeds.	<u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Eastern Bluebird <i>Sialia sialis</i> 766.0 EABL Summer resident Uncommon Life Form 14 17	(FL)	NG NTMB	Cottonwood-riparian, eastern great plains grasslands, agricultural areas.	Often nests in a woodpecker cavity, usually in a snag. Feeds on insects, fruit, some invertebrates.	<u> </u> <u>O</u> <u> </u> <u>O*</u> <u>O*</u> <u> </u> <u>B*</u> <u>O*</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>O*</u> <u>O</u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>B</u> <u>B</u>
Western Bluebird <i>Sialia mexicana</i> 767.0 WEBL Summer resident Uncommon Life Form 14	(AS)	NG NTMB	Pine-juniper, juniper woodlands, associated with edges.	Often nests in a woodpecker cavity in a snag. Feeds on insects, fruit, some invertebrates.	<u>O+</u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>O*</u> <u>O*</u> <u>O*</u> <u>O</u> <u>O</u> <u> </u> <u>B</u> <u>O*</u> <u>O</u> <u>O</u> <u>O*</u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>b</u> <u>O</u> <u> </u> <u>B</u> <u>O*</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Mountain Bluebird <i>Sialia currucoides</i> 768.0 MOBL Summer resident Common Life Form 14 17		NG NTMB	Most habitats with nesting cavities and open areas for foraging.	Nests usually in a woodpecker cavity in a snag. Feeds primarily on insects; also fruit.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Townsend's Solitaire <i>Myadestes townsendi</i> 754.0 TOSO Resident Common Life Form 4 6		NG NTMB	Coniferous forests, aspen.	Nests often amid tree roots or other shelter on the ground. Feeds on insects, fruit, worms.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u>
Veery <i>Catharus fuscescens</i> 756.0 VEER Summer resident Uncommon Life Form 5		NG NTMB	Aspen, cottonwood-riparian, coniferous forests, below 9,000 feet.	Nests on the ground or in a shrub. Feeds on insects, some fruit.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u>
Gray-cheeked Thrush <i>Catharus minimus</i> 757.0 GCTH Migrant Uncommon Life Form	(AS)	NG NTMB	Coniferous forests.	Feeds on insects, berries. L28: W. Schreier, 15 May 1993.	<u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u>O*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Swainson's Thrush <i>Catharus ustulatus</i> 758.0 SWTH Summer resident Common Life Form 7		NG NTMB	Coniferous forests, aspen, cottonwood-riparian, riparian shrub, juniper.	Nests in a shrub or low in a coniferous tree. Feeds on insects, berries.	<u>B</u> <u>B</u> <u>—</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u>
Hermit Thrush <i>Catharus guttatus</i> 759.0 HETH Summer resident Common Life Form 5 7		NG NTMB	Coniferous and mixed aspen/coniferous forests, juniper.	Nests on the ground or low in a coniferous tree. Feeds on insects, berries.	<u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>—</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Wood Thrush <i>Hylocichla mustelina</i> 755.0 WOTH Migrant Rare Life Form	(AS)	NG NTMB	Deciduous or mixed deciduous/coniferous forests.	Feeds on insects, berries.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u>
American Robin <i>Turdus migratorius</i> 761.0 AMRO Resident Abundant Life Form 11		NG NTMB	All habitats with trees or shrubs present.	Nests in a deciduous or coniferous tree, or a building. Feeds on insects, earthworms, fruit. Winter populations and distribution are considerably smaller than during other seasons.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Varied Thrush <i>Ixoreus naevius</i> 763.0 VATH Migrant Rare Life Form	(AS)	NG	Found in moist coniferous and deciduous forests.	Feeds on insects, berries.	<u>O+</u> <u>O*</u> <u> </u> <u> </u> <u>O*</u> <u>O*</u> <u>O*</u> <u>O*</u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u>O*</u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O*</u>
Gray Catbird <i>Dumetella carolinensis</i> 704.0 GRCA Summer resident Common Life Form 7		NG NTMB	Riparian shrub below 8,000 feet, urban areas.	Nests often in a dense thicket. Feeds on insects, berries, other fruit.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u>
Brown Thrasher <i>Toxostoma rufum</i> 705.0 BRTH Summer resident Common Life Form 5 7		NG	Riparian shrub, woodland-chaparral, mountain mahogany, bitterbrush, serviceberry, chokecherry, cottonwood-riparian.	Nests in or beneath a low shrub. Feeds on insects, invertebrates, small vertebrates, berries, fruit, nuts.	<u> </u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u> </u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u>
Sage Thrasher <i>Oreoscoptes montanus</i> 702.0 SATH Summer resident Common Life Form 5 7		NG NSS4 NTMB	Basin-prairie and mountain-foothills shrublands.	Nest is concealed in or beneath a sagebrush shrub. Feeds on insects, some fruit.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Northern Mockingbird <i>Mimus polyglottos</i> 703.0 NOMO Summer resident Peripheral Uncommon Life Form 8		NG NTMB	Cottonwood-riparian, riparian shrub, basin-prairie shrublands, urban areas.	Nests in a conifer or a deciduous shrub. Feeds on insects, fruit, some invertebrates, a few small vertebrates.	_ _ O _ _ O O O _ _ O O O O O B O _ O O _ _ O O b _ b O b O b B
European Starling <i>Sturnus vulgaris</i> 493.0 EUST Resident Abundant Life Form 14 17		PD	Cottonwood-riparian, agricultural areas, urban areas.	Nests in a cavity. Feeds on insects, fruit, seeds. Non-native species brought into the U.S. from Europe. Is aggressive toward native songbirds. One of four bird species that can be legally killed without a license or permit.	B B B B B B B B B B B b B b B b B B B B b B B b B B B B
House Sparrow <i>Passer domesticus</i> 688.2 HOSP Resident Abundant Life Form 14 17		PD	Agricultural areas, urban areas.	Nests in an artificial or natural cavity. Feeds on seeds, insects, fruit. Non-native species brought into the U.S. from Europe. Is aggressive toward native songbirds. One of four bird species that can be legally killed without a license or permit.	B B B B B B B B b B b B B b B b B B B B b B B B B B B B
American Pipit <i>Anthus rubescens</i> 697.0 AMPI Summer resident Common Life Form 5		NG NSS4 NTMB	Alpine grasslands, alpine moss-lichen-forb. Lower habitats during migration.	Nest is sunk in the ground and partially overhung by a rock or vegetation. Feeds on insects, aquatic invertebrates, seeds, some fruit.	B B O B O O O b B O O O O O b O b _ _ O O O O O O O B O O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b																												
Sprague's Pipit <i>Anthus spragueii</i> 700.0 SPPI Migrant Rare Life Form	(AS)	NG NTMB	Eastern great plains grasslands.	Feeds primarily on insects, grass seeds, forb seeds. Nesting should be expected in northern Wyoming. L5: H. Downing, 7 May 1982. L19: J. Herold and V. Herold, 27 September 1994. L28: J. Lawrence, 16 May 2008.	<table border="0"> <tr><td>—</td><td>○</td><td>—</td><td>○</td><td>○*</td><td>○</td><td>—</td></tr> <tr><td>○</td><td>○</td><td>—</td><td>○</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>○*</td><td>—</td><td>○</td></tr> <tr><td>—</td><td>○</td><td>—</td><td>—</td><td>—</td><td>○</td><td>○*</td></tr> </table>	—	○	—	○	○*	○	—	○	○	—	○	—	—	—	—	—	—	—	○*	—	○	—	○	—	—	—	○	○*
—	○	—	○	○*	○	—																											
○	○	—	○	—	—	—																											
—	—	—	—	○*	—	○																											
—	○	—	—	—	○	○*																											
Brambling <i>Fringilla montifringilla</i> 514.1 BRAM Accidental Rare Life Form	(AS)		Habitat requirements unknown; this is an accidental species from Eurasia.	Feeds on seeds. European species. L5: H. Downing, 18 and 23 November and 1 December 1985. L9: M. Back, 10-26 November 1985. L26: F. Bergquist, 21-25 November 2014.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>○*</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>○*</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>○*</td><td>—</td><td>—</td></tr> </table>	—	—	—	—	○*	—	—	—	○*	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	○*	—	—
—	—	—	—	○*	—	—																											
—	○*	—	—	—	—	—																											
—	—	—	—	—	—	—																											
—	—	—	—	○*	—	—																											
Gray-crowned Rosy-Finch <i>Leucosticte tephrocotis</i> 524.0 GCRF Resident Common Life Form 4		NG	Alpine grasslands, alpine moss-lichen-forb, barren ground, fallow agricultural areas. A variety of habitats during the winter.	Nests on the ground or on a cliff. Feeds on seeds, insects.	<table border="0"> <tr><td>B</td><td>B</td><td>○</td><td>B</td><td>b</td><td>○</td><td>○</td></tr> <tr><td>b</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td></tr> <tr><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td></tr> <tr><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td></tr> </table>	B	B	○	B	b	○	○	b	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○	○
B	B	○	B	b	○	○																											
b	○	○	○	○	○	○																											
○	○	○	○	○	○	○																											
○	○	○	○	○	○	○																											
Black Rosy-Finch <i>Leucosticte atrata</i> 525.0 BLRF Resident Uncommon Life Form 4		NG NSSU	Alpine grasslands, alpine moss-lichen-forb, barren ground, fallow agricultural areas. A variety of habitats during the winter.	Nests on the ground or on a cliff. Feeds on seeds, insects.	<table border="0"> <tr><td>B</td><td>B</td><td>—</td><td>b</td><td>○</td><td>—</td><td>○</td></tr> <tr><td>B</td><td>B</td><td>○</td><td>○</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>b</td><td>b</td><td>○</td><td>—</td><td>○</td><td>○</td><td>—</td></tr> <tr><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td><td>○</td></tr> </table>	B	B	—	b	○	—	○	B	B	○	○	—	—	—	b	b	○	—	○	○	—	○	○	○	○	○	○	○
B	B	—	b	○	—	○																											
B	B	○	○	—	—	—																											
b	b	○	—	○	○	—																											
○	○	○	○	○	○	○																											

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Brown-capped Rosy-Finch <i>Leucosticte australis</i> 526.0 BCRF Resident Uncommon Life Form 4	(FL)	NG NSSU	Alpine grasslands, alpine moss-lichen-forb, barren ground, fallow agricultural areas. A variety of habitats during the winter.	Nests on the ground or on a cliff. Feeds on seeds, insects. L26: D. McDonald and T. Parchman, 18 July 2005.	— — — O — — — — O — — — — — — O* — — — — — — O O O B* O — —
Pine Grosbeak <i>Pinicola enucleator</i> 515.0 PIGR Resident Uncommon Life Form 9		NG	Coniferous forests, especially Douglas fir and Englemann spruce-subalpine fir, deciduous forests.	Usually nests in a conifer, occasionally in a shrub. Feeds on seeds, buds, fruit, some insects.	B B — B b — O b b — — — — O b O b — O b O b O — O B B O
House Finch <i>Haemorhous mexicanus</i> 519.0 HOFI Resident Common Life Form 8 11 17		NG	Coniferous and deciduous forests below 8,000 feet, woodland-chaparral, urban areas.	Nests in a variety of locations, usually in a tree, shrub, or on a building. Feeds on seeds, fruit, buds, tree sap.	O B b B B O O b O B b — O O b b B B B b b B B B B b B B
Purple Finch <i>Haemorhous purpureus</i> 517.0 PUF1 Winter resident Uncommon Life Form	(AS)	NG NTMB	Urban areas.	Feeds primarily on seeds; also insects, fruit. Most observations are at feeders.	— O — O O* O O* O* — — O — — — — — O — O* O — — O — — O O —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Cassin's Finch <i>Haemorhous cassinii</i> 518.0 CAFI Resident Common Life Form 10		NG NTMB	Coniferous forests up to timberline, including burns. Lower habitats during the winter, especially urban areas.	Nests in a conifer; nest is usually placed near the end of a large limb. Feeds on buds, berries, conifer seeds.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u>
Red Crossbill <i>Loxia curvirostra</i> 521.0 RECR Resident Common Life Form 10		NG NSS4	Coniferous forests. Lower habitats during the winter including urban areas.	Nest is far from the trunk on a horizontal branch of a conifer. Feeds primarily on conifer seeds; also buds, forb seeds, berries, a few insects.	<u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u>
White-winged Crossbill <i>Loxia leucoptera</i> 522.0 WWCR Resident Uncommon Life Form 10	(FL)	NG	Coniferous forests.	Nests in a conifer; nest is usually far from the trunk on a horizontal branch. Feeds primarily on conifer seeds; also other seeds, a few berries, insects. L1, 4, 8, 17, 19, 26: summer observations. L15: S. Patla, B. Oakleaf, and T. Laurion , 7 July 2000. L19: S. Fitton, 25 June 1981.	<u>b</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>B</u> <u>b</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u>O</u> <u> </u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u>
Common Redpoll <i>Acanthis flammea</i> 528.0 CORE Winter resident Uncommon Life Form		NG	Basin-prairie and mountain-foothills shrublands, grasslands, at feeders in urban areas.	Feeds primarily on seeds of deciduous and coniferous shrubs; also insects when available.	<u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Hoary Redpoll <i>Acanthis hornemanni</i> 527.0 HORE Winter resident Rare Life Form	(AS)	NG	Basin-prairie and mountain-foothills shrublands, grasslands, at feeders in urban areas.	Feeds primarily on seeds of deciduous shrubs, forbs, and grasses. L5: H. Downing, 7-9 November and 16 December 1984.	— — — — — — — — — — O —
Pine Siskin <i>Spinus pinus</i> 533.0 PISI Resident Common Life Form 10		NG NTMB	Coniferous forests, aspen, cottonwood-riparian. Lower habitats during the winter including urban areas.	Usually nests in a conifer; nest is placed on a horizontal limb far out from the trunk. Feeds on seeds, insects.	B B B B B O B B B B B O b O B b B b B b B b b b B B B B
Lesser Goldfinch <i>Spinus psaltria</i> 530.0 LEGO Migrant Uncommon Life Form	(FL)	NG NTMB	Cottonwood-riparian, riparian shrub, urban areas.	Feeds on seeds, insects, berries, floral buds.	O* O O O O* — — — O* O* O* — — O* — — — — — O* O O* O* O* — — — O O* O*
Lawrence's Goldfinch <i>Spinus lawrencei</i> 531.0 LAGO Accidental Rare Life Form	(AS)	NTMB	Cottonwood-riparian, riparian shrub, woodland chaparral, pine-juniper, arid weedy areas near water.	Feeds on seeds, insects. L21: D. Felley, 17 June 1999.	— —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
American Goldfinch <i>Spinus tristis</i> 529.0 AMGO Resident Common Life Form 11		NG NTMB	Cottonwood-riparian and riparian shrub below 8,000 feet, urban areas.	Nests in a fork of a branch in a shrub or tree. Feeds on seeds of deciduous trees, forbs, especially composites, grasses; also floral buds, berries.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u>
European Goldfinch <i>Carduelis carduelis</i> 526.1 EUGO Accidental Controlled Life Form			Not native to North America; many attempts to introduce this species have failed. Brought to North America for private collections; occasionally escapes from captivity.	Feeds on seeds. L19: G. Chase, 6 June 2004, probably released or escaped from captivity.	— — — — — — — — — — — — — — — — — — — <u>O*</u> — — — — — — — — — —
Evening Grosbeak <i>Coccothraustes vespertinus</i> 514.0 EVGR Resident Common Life Form 10		NG	Coniferous and deciduous forests, urban areas.	Nest is usually far out on a horizontal limb of a conifer. Feeds primarily on seeds, especially pinyon nuts and juniper berries; also some insects.	<u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>O</u>
Bohemian Waxwing <i>Bombycilla garrulus</i> 618.0 BOWA Winter resident Common Life Form		NG	Pine-juniper, cottonwood-riparian, woodland-chaparral, urban areas.	Feeds on insects, berries, fruit. Locally abundant in some winters.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Cedar Waxwing <i>Bombycilla cedrorum</i> 619.0 CEDW Resident Uncommon Life Form 11		NG NTMB	Pine-juniper, cottonwood-riparian, juniper, urban areas.	Nests usually in a conifer. Feeds on fruit, berries, flowers, tree sap, insects.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u>
Lapland Longspur <i>Calcarius lapponicus</i> 536.0 LALO Winter resident Uncommon Life Form		NG	Eastern great plains and great basin-foothills grasslands, agricultural areas.	Feeds on insects, seeds. L8: S. Williams and B. Raynes, 1 January-25 March 1999. L16: R. Steenberg, 21 January 1995. L21: K. Kranik, December 2001.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u> <u>—</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u>
Chestnut-collared Longspur <i>Calcarius ornatus</i> 538.0 CCLO Summer resident Uncommon Life Form 5	(FL)	NG NSS4 NTMB	Eastern great plains and great basin-foothills grasslands, basin-prairie shrublands, agricultural areas.	Nests on the ground in a shallow depression, usually concealed by a tuft of grass. Feeds on insects, seeds.	<u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>—</u> <u>b</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>b</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>O</u> <u>b</u> <u>B</u>
Smith's Longspur <i>Calcarius pictus</i> 537.0 SMLO Accidental Rare Life Form	(AS)		Grasslands.	Feeds on insects, seeds. L7: D. Svingen, 26 September 2007. L19: V. Herold and J. Herold, 2 October 1994.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
McCown's Longspur <i>Rhynchophanes mccownii</i> 539.0 MCLO Summer resident Common Life Form 5		NG NSS4 NTMB	Eastern great plains and great basin-foothills grasslands, basin-prairie shrublands, agricultural areas.	Nests on the ground in a shallow, natural or scraped depression. Feeds on seeds, insects.	<u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u> </u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u> </u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u>
Snow Bunting <i>Plectrophenax nivalis</i> 534.0 SNBU Winter resident Uncommon Life Form		NG	Basin-prairie and mountain-foothills shrublands, grasslands.	Feeds primarily on insects, seeds. Common during some winters.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u> </u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u>
Ovenbird <i>Seiurus aurocapilla</i> 674.0 OVEN Summer resident Uncommon Life Form 5		NG NTMB	Aspen, cottonwood-riparian.	Nests on the leaf-covered forest floor. Feeds mostly on insects.	<u>O</u> <u>O*</u> <u> </u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>B</u> <u>O</u> <u> </u> <u>O</u> <u> </u> <u>b</u> <u>b</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u>O</u> <u>O</u>
Worm-eating Warbler <i>Helmitheros vermivorum</i> 639.0 WEWA Accidental Rare Life Form	(AS)	NTMB	Deciduous forests with heavy undergrowth associated with slopes.	Feeds entirely on insects.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Northern Waterthrush <i>Parkesia noveboracensis</i> 675.0 NOWA Migrant Uncommon Life Form		NG NTMB	Aspen, cottonwood-riparian, willow.	Feeds on aquatic invertebrates, insects. L1, 5, 8, 27: summer records.	<u>0</u> <u>0</u> <u>—</u> <u>0</u> <u>0</u> <u>0</u> <u>—</u> <u>b</u> <u>0</u> <u>0</u> <u>—</u> <u>—</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>0</u> <u>—</u> <u>0</u> <u>0</u> <u>0</u> <u>—</u> <u>0</u> <u>—</u> <u>b</u> <u>0</u>
Golden-winged Warbler <i>Vermivora chrysoptera</i> 642.0 GWWA Accidental Rare Life Form	(AS)	NTMB	Overgrown native meadows.	Feeds primarily on insects.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u>
Blue-winged Warbler <i>Vermivora cyanoptera</i> 641.0 BWWA Accidental Rare Life Form	(AS)	NTMB	Second growth woodlands, mountain-foothills shrublands, marsh-swamp wetlands, riparian shrub.	Feeds on insects, spiders. L15: J. Larson, 28 May 1995. L19: J. Gaither, B. South, and J. Herold, 23 July 1994.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u>
Black-and-white Warbler <i>Mniotilta varia</i> 636.0 BAWW Migrant Uncommon Life Form	(FL)	NG NTMB	Deciduous forests.	Feeds almost entirely on insects.	<u>0*</u> <u>—</u> <u>—</u> <u>0</u> <u>0*</u> <u>0</u> <u>0*</u> <u>0*</u> <u>0</u> <u>—</u> <u>0</u> <u>—</u> <u>—</u> <u>—</u> <u>0*</u> <u>0</u> <u>—</u> <u>—</u> <u>0</u> <u>0</u> <u>0</u> <u>—</u> <u>0*</u> <u>—</u> <u>0</u> <u>—</u> <u>0</u> <u>0</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Prothonotary Warbler <i>Protonotaria citrea</i> 637.0 PROW Accidental Rare Life Form	(AS)	NTMB	Deciduous or coniferous forests associated with marsh-swamp wetlands.	Feeds on insects, snails. L23: M. Adams, 2 September 1986; F. Luke, 2 September 1989; V. Herren and A. Dahl, 16 October 2000. L27: McCreary (1939).	<u>O</u> _ _ _ _ _ _ _ _ _ _ _ _ _ _ O* _ _ _ O* _ _ _ h _
Tennessee Warbler <i>Oreothlypis peregrina</i> 647.0 TEWA Migrant Uncommon Life Form	(FL)	NG NTMB	Deciduous forests, urban areas.	Feeds primarily on insects, some fruit.	<u>O*</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ <u>O</u> <u>O*</u> _ <u>O</u> <u>O</u> _ <u>O*</u> _ <u>O</u> _ <u>O</u> _ <u>O</u> _ <u>O</u> <u>O*</u> <u>O</u> <u>O</u>
Orange-crowned Warbler <i>Oreothlypis celata</i> 646.0 OCWA Summer resident Uncommon Life Form 6		NG NTMB	Aspen, cottonwood-riparian.	Nest is usually hidden on the ground. Feeds on insects, fruit, nectar, tree sap.	<u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u>
Nashville Warbler <i>Oreothlypis ruficapilla</i> 645.0 NAWA Migrant Rare Life Form	(FL)	NG NTMB	Deciduous forests, juniper, urban areas.	Feeds entirely on insects.	<u>O+</u> _ _ <u>O*</u> <u>O*</u> _ <u>O</u> <u>b*</u> _ <u>O*</u> _ _ _ _ <u>O</u> _ _ <u>O</u> <u>O*</u> _ _ _ <u>O*</u> <u>O</u> _ _ _ <u>O</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Virginia's Warbler <i>Oreothlypis virginiae</i> 644.0 VIWA Summer resident Uncommon Life Form 6	(FL)	NG NSSU NTMB	Pinyon-juniper, woodland-chaparral.	Nests on the ground, usually hidden by vegetation. Feeds entirely on insects.	<u> </u> <u> </u> <u> </u> <u>B*</u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b*</u> <u> </u> <u>O*</u> <u>b</u> <u>b</u> <u>O*</u> <u>O</u> <u> </u> <u>O</u> <u>B*</u> <u>b</u> <u>O</u> <u> </u> <u>O</u> <u>b</u>
Connecticut Warbler <i>Oporornis agilis</i> 678.0 CONW Accidental Rare Life Form	(AS)	NTMB	Englemann spruce forests associated with swamps.	Feeds mainly on insects. L8: T. Gorman, 20 September 2001. L15: F. Luke, 28 May 1988. L28: S. Dinsmore, 18 May 2002.	<u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u>
MacGillivray's Warbler <i>Geothlypis tolmiei</i> 680.0 MGWA Summer resident Common Life Form 8		NG NSS4 NTMB	Aspen, cottonwood-riparian, coniferous forests, riparian shrub, logged areas, burned areas, downed timber and windfalls, below 9,000 feet.	Nests close to the ground in dense shrubs. Feeds mostly on insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u> </u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u>
Mourning Warbler <i>Geothlypis philadelphia</i> 679.0 MOWA Accidental Rare Life Form	(AS)	NTMB	Dense shrublands in open deciduous forests.	Feeds mainly on insects. L15: F. Luke, 27 May 1989; R. Steenberg, 27 May 2001. L28: P. Lehman, 27 May 2004.	<u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O*</u> <u>O*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Kentucky Warbler <i>Geothlypis formosus</i> 677.0 KEWA Accidental Rare Life Form	(AS)	NTMB	Deciduous forests with heavy undergrowth.	Feeds mostly on insects. L2: C. Neal, 27 May 2000. L19: L. Rognstad, 15 May 1981. L28: S. Streeper, 30 April 1987; F. Lebsack, 17 May 1989.	— O* — — — — O* — — — — — — — — — — — — O* — — — — — — — — — O*
Common Yellowthroat <i>Geothlypis trichas</i> 681.0 COYE Summer resident Common Life Form 3		NG NSS4 NTMB	Willow and marshes below 8,000 feet.	Nests on the ground. Feeds mostly on insects.	B B B B B b B B b B b b B b B O B O B B B b b b b B B b
Hooded Warbler <i>Setophaga citrina</i> 684.0 HOWA Accidental Rare Life Form	(AS)	NTMB	Heavy undergrowth in moist deciduous forests.	Feeds mostly on insects. L6: R. Cox, K. Missett, and V. Hays, 13 June 1999. L15: J. Larson and B. Boevers, 19-20 May 1990; R. Steenberg, S. Seppio, and R. Dorn, 21 and 24 May 1989; R. Steenberg, J. Steenberg, and D. Kerr, 22 May 2009. L16: A. King, 5 June 2004.	O+ — — — — O* — — — — — — O — O* O* — — — O — — — O* — — — — O O*
American Redstart <i>Setophaga ruticilla</i> 687.0 AMRE Summer resident Uncommon Life Form 9		NG NTMB	Cottonwood-riparian, riparian shrub.	Usually nests in a fork of a low deciduous tree or shrub. Feeds almost entirely on insects. Locally common.	B b — — B B O B B b O B b O B O O b O B B O b b O b B O O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Cape May Warbler <i>Setophaga tigrina</i> 650.0 CMWA Accidental Rare Life Form	(AS)	NTMB	Englemann spruce-subalpine fir forests.	Feeds primarily on insects, occasionally fruit, nectar. L1: 16 June 1975, American Birds 30:104. L15: F. Luke, 26 May 1990.	<u>0</u> _ _ _ _ ? _ _ _ _ _ _ _ _ _ _ _ <u>0*</u> _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _
Northern Parula <i>Setophaga americana</i> 648.0 NOPA Migrant Rare Life Form	(FL)	NG NTMB	Cottonwood-riparian.	Feeds mostly on insects.	_ _ _ _ _ <u>0*</u> <u>0*</u> _ _ <u>0*</u> _ _ _ _ _ _ _ _ <u>0*</u> <u>0*</u> <u>0*</u> _ _ <u>0</u> _ <u>0*</u> <u>0</u> <u>0*</u> _ _ _ <u>0*</u> <u>0</u> <u>0*</u>
Magnolia Warbler <i>Setophaga magnolia</i> 657.0 MAWA Migrant Rare Life Form	(FL)	NG NTMB	Coniferous forests.	Feeds almost entirely on insects.	_ _ _ _ <u>0</u> <u>0*</u> <u>0*</u> <u>0*</u> _ <u>0</u> _ _ _ _ _ _ <u>0</u> <u>0*</u> _ _ _ <u>0*</u> <u>0</u> <u>0*</u> _ <u>0*</u> _ _ _ _ <u>0</u> <u>0*</u>
Bay-breasted Warbler <i>Setophaga castanea</i> 660.0 BBWA Migrant Rare Life Form	(AS)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds mostly on insects; also a few berries. L15: F. Luke, M. Adams, and R. Dean, 2 June 1990; F. Luke and others, 23-26 May 1991. L19: L. Rognstad and B. Stratton, 24 May 1990; A. Hines and W. Hines, 25 September 1993. L22: C. Kesselheim, 13 October 1985.	_ _ _ _ _ _ _ _ <u>0</u> _ _ _ _ _ _ _ _ <u>0*</u> _ _ _ _ <u>0*</u> _ _ <u>0*</u> _ _ _ _ _ _ <u>0*</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Blackburnian Warbler <i>Setophaga fusca</i> 662.0 BLBW Migrant Rare Life Form	(AS)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds primarily on insects, occasionally some berries.	O* _ _ O* O _ _ O* _ O _ _ _ O* O* _ _ O* _ _ _ O* _ _ _ O O*
Yellow Warbler <i>Setophaga petechia</i> 652.0 YEWA Summer resident Abundant Life Form 8		NG NTMB	Riparian shrub below 8,000 feet, urban areas.	Nests in a small deciduous tree or shrub. Feeds mainly on insects, some berries.	B B B B B B B B B B b b B b B B B b B B B B B B B B B B
Chestnut-sided Warbler <i>Setophaga pensylvanica</i> 659.0 CSWA Migrant Rare Life Form	(FL)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds primarily on insects, occasionally berries.	_ O* _ O O O* O* O* _ _ _ _ O _ O* _ O* _ O* _ O* _ O* _ _ _ O O
Blackpoll Warbler <i>Setophaga striata</i> 661.0 BLPW Migrant Uncommon Life Form	(FL)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds primarily on insects, some berries, seeds.	O O* _ O* O* O O* _ _ _ _ _ O O* _ O _ O O O _ O _ _ O O O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b																												
Black-throated Blue Warbler <i>Setophaga caeruleascens</i> 654.0 BTBW Migrant Rare Life Form	(FL)	NG NTMB	Deciduous and mixed deciduous/ coniferous forests with heavy undergrowth.	Feeds primarily on insects; also seeds, other vegetation.	<table border="0"> <tr><td>—</td><td>O</td><td>—</td><td>—</td><td>O*</td><td>O*</td><td>O*</td></tr> <tr><td>O</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>O</td><td>—</td></tr> <tr><td>O*</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>O*</td><td>—</td><td>—</td><td>O*</td><td>O</td><td>O*</td></tr> </table>	—	O	—	—	O*	O*	O*	O	O*	—	—	—	O	—	O*	—	—	—	O*	—	—	—	O*	—	—	O*	O	O*
—	O	—	—	O*	O*	O*																											
O	O*	—	—	—	O	—																											
O*	—	—	—	O*	—	—																											
—	O*	—	—	O*	O	O*																											
Palm Warbler <i>Setophaga palmarum</i> 672.0 PAWA Migrant Rare Life Form	(AS)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds almost entirely on insects. L5: J. Ward, 8 May 1996. L19: V. Herold and J. Herold, 24 August 1993. L28: F. Lebsack and others, 29 April and 1-4 May 1989.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td></tr> <tr><td>O</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>O</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>O*</td></tr> </table>	—	—	—	—	O*	—	—	O	—	—	—	—	—	—	—	—	—	—	O*	—	O	—	—	—	—	—	O*	O*
—	—	—	—	O*	—	—																											
O	—	—	—	—	—	—																											
—	—	—	—	O*	—	O																											
—	—	—	—	—	O*	O*																											
Pine Warbler <i>Setophaga pinus</i> 671.0 PIWA Accidental Rare Life Form	(AS)		Mostly coniferous forests; also mixed coniferous/deciduous forests.	Feeds mainly on insects, occasionally some seeds, fruit. L6: V. Hays and F. Grams, 6- 19 December 1988. L19: O.K. Scott, 1985. L20: American Birds 31:207.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>O</td><td>O</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td></tr> </table>	—	—	—	—	—	O*	—	—	—	—	—	—	—	—	—	—	—	—	O	O	—	—	—	—	—	—	O*	—
—	—	—	—	—	O*	—																											
—	—	—	—	—	—	—																											
—	—	—	—	O	O	—																											
—	—	—	—	—	O*	—																											
Yellow-rumped Warbler <i>Setophaga coronata</i> 655.6 YRWA Summer resident Common Life Form 11		NG NTMB	Coniferous forests, aspen, cottonwood-riparian.	Nests on a horizontal branch of a conifer. Feeds primarily on insects, occasionally some fruit, berries.	<table border="0"> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>b</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>O</td><td>B</td><td>B</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>b</td><td>O</td></tr> <tr><td>B</td><td>B</td><td>B</td><td>B</td><td>b</td><td>B</td><td>O</td></tr> </table>	B	B	B	B	B	b	B	B	B	B	B	O	B	B	B	B	B	B	B	b	O	B	B	B	B	b	B	O
B	B	B	B	B	b	B																											
B	B	B	B	O	B	B																											
B	B	B	B	B	b	O																											
B	B	B	B	b	B	O																											

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Yellow-throated Warbler <i>Setophaga dominica</i> 663.0 YTWA Accidental Rare Life Form	(AS)	NTMB	Coniferous/oak forests.	Feeds primarily on insects.	<p>— — — — — O*</p> <p>— — — — — — — —</p> <p>— — — — — O* O*</p> <p>— — O* — — — — O*</p>
Prairie Warbler <i>Setophaga discolor</i> 673.0 PRAW Accidental Rare Life Form	(AS)	NTMB	Woodland-chaparral, mountain-foothills shrublands, mountain-foothills grasslands, mixed coniferous/ deciduous forest edge.	Feeds on insects. L28: R. Dorn and J. Dorn, 18 May 1996.	<p>— — — — — — — —</p> <p>— — — — — — — —</p> <p>— — — — — — — —</p> <p>— — — — — — — — O*</p>
Black-throated Gray Warbler <i>Setophaga nigrescens</i> 665.0 BTYW Summer resident Uncommon Life Form 10	(FL)	NG NSS4 NTMB	Pine-juniper, woodland-chaparral, mountain-foothills shrublands.	Nests far out on a horizontal branch, usually in a conifer. Feeds almost entirely on insects.	<p>— — O — — b* — — — —</p> <p>O* O* B — — — — —</p> <p>O O O* b b — — — —</p> <p>b B b b — — O* O</p>
Townsend's Warbler <i>Setophaga townsendi</i> 668.0 TOWA Summer resident Uncommon Life Form 10		NG NTMB	Douglas fir, Englemann spruce-subalpine fir, other or mixed coniferous forests, urban areas.	Nests usually far out on a horizontal limb of a fir. Feeds primarily on insects. Expected to be found nesting in northwestern Wyoming. L22: McCreary (1939).	<p>O O — — O O O O</p> <p>O O O — — — — —</p> <p>O O O O O O O O</p> <p>H O O O O O O O</p>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Red-faced Warbler <i>Cardellina rubrifrons</i> 690.0 RFWA Accidental Rare Life Form	(AS)	NTMB	Coniferous forests, oak, aspen, mostly between 6,400 and 8,000 feet.	Feeds mainly on insects. L8: D. Browne, 29 May 2008. L23: F. Luke, 29 April-3 May 1989.	— — — — — — — — O* — — — — — — — — — — — — — — — — — O* — — — — — — — —
Yellow-breasted Chat <i>Icteria virens</i> 683.0 YBCH Summer resident Common Life Form 3 8		NG NTMB	Riparian shrub and marshes below 7,000 feet.	Nests beneath or in a low shrub; nest is usually large but well concealed. Feeds on insects, berries.	O O B B B O B O O B O b — B O O B O B b B b B — b — b O
Green-tailed Towhee <i>Pipilo chlorurus</i> 590.0 GTTO Summer resident Common Life Form 8		NG NTMB	Mixed coniferous forests, woodland- chaparral, juniper-sagebrush, basin- prairie and mountain-foothills shrublands, riparian shrub.	Nests in a shrub or on the ground. Feeds on insects, seeds, berries.	B B b B b O B B B B B B b O B B B B B B B b B B B B B B
Spotted Towhee <i>Pipilo maculatus</i> 588.0 SPTO Summer resident Common Life Form 5 7		NG NTMB	Pine-juniper, aspen, woodland- chaparral, juniper-sagebrush, basin- prairie and mountain-foothills shrublands, riparian shrub.	Nests often in a scratched depression on the ground. Feeds on insects, seeds, fruit.	B b b B B b B O O B O b O B O O B B b b B O B — b b B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Canyon Towhee <i>Melospiza fusca</i> 591.0 CANT Accidental Rare Life Form	(AS)		Woodland-chaparral, riparian shrub, basin-prairie shrublands, urban areas.	Feeds on seeds, insects, berries. L8: B. Raynes, 3 June 1979.	— — — — — — — — O —
Cassin's Sparrow <i>Peucaea cassinii</i> 578.0 CASP Accidental Rare Life Form	(AS)	NTMB	Eastern great plains and great basin-foothills grasslands, sagebrush-grasslands.	Feeds on insects. Species has been expanding its breeding range. All sightings should be reported, except confirmed breeding in the Torrington area (L21). L12: C. Faanes, 8 June 1978, American Birds 32:1190. L14: A. Knue, 1 July 2008. L21: W. Howe, 28 June 1990.	— — — — — — — — — — — — — O* — O* —
American Tree Sparrow <i>Spizelloides arborea</i> 559.0 ATSP Winter resident Uncommon Life Form		NG	Cottonwood-riparian, riparian shrub, agricultural areas including shelterbelts, roadsides, urban areas.	Feeds on seeds, buds, berries in the winter.	O O O O O O O O O O O O O O O O O O O — O O O O O O O O O O O O
Chipping Sparrow <i>Spizella passerina</i> 560.0 CHSP Summer resident Common Life Form 7		NG NTMB	Coniferous forests, aspen, cottonwood-riparian, woodland-chaparral, up to 10,000 feet.	Nests in a tree. Feeds on insects, seeds.	B B B B B b B B B B B b b B B B B b B b B b B B B B B B b B B B

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Clay-colored Sparrow <i>Spizella pallida</i> 561.0 CCSP Summer resident Uncommon Life Form 7		NG NTMB	Ponderosa pine savannah, pine-juniper, aspen, cottonwood-riparian, mountain-foothills shrublands, sagebrush-grasslands, shelterbelts.	Nests in a shrub or on the ground. Feeds on insects, seeds. L6: H. Downing, 10 May 1996. L6: R.T. Cox and K. Missett, 9 May 1999. L16: D. Thomas, 22 June 2013.	<u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>O*</u> <u>b</u> <u>B</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u>
Brewer's Sparrow <i>Spizella breweri</i> 562.0 BRSP Summer resident Common Life Form 7		NG NSS4 NTMB	Basin-prairie and mountain-foothills shrublands, especially sagebrush, woodland-chaparral.	Nests in a shrub. Feeds on insects, seeds.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Field Sparrow <i>Spizella pusilla</i> 563.0 FISP Summer resident Abundance unknown Life Form 6 8	(AS)	NG	Old fields, shrubs, edge of deciduous forests.	Nests on the ground or in a small tree. Feeds on insects, seeds. Nesting pairs should be present in eastern Wyoming. L5: H. Corbett, 23 March 2000. L19: J. Lawrence, 8 October 2008. L7: J. Adams, 8-14 May 2014.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u>
Vesper Sparrow <i>Pooecetes gramineus</i> 540.0 VESP Summer resident Common Life Form 5		NG NTMB	Basin-prairie and mountain-foothills shrublands, grasslands, agricultural areas.	Nests in an excavated depression on the ground. Feeds half on insects, half on grass and forb seeds.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Lark Sparrow <i>Chondestes grammacus</i> 552.0 LASP Summer resident Common Life Form 6		NG NTMB	Pine-juniper, woodland-chaparral, basin-prairie and mountain-foothills shrublands, grasslands, agricultural areas.	Nests in a hollow depression on the ground. Feeds on seeds, insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u>
Black-throated Sparrow <i>Amphispiza bilineata</i> 573.0 BTSP Summer resident Abundance unknown Life Form 7	(AS)	NG NTMB	Basin-prairie shrublands, especially in rocky areas.	Nests in a shrub or cactus. Feeds on insects, seeds, some new shoots of grasses and forbs. Documentation of breeding should be sought in southwestern Wyoming. L21, 26: McCreary (1939).	<u>O*</u> <u>O*</u> — — — — — <u>O</u> — — — — — — — — — — <u>h</u> — <u>O*</u> <u>O*</u> — <u>h</u> <u>O</u> <u>O*</u>
Sagebrush Sparrow <i>Artemisospiza nevadensis</i> 574.3 SBSP Summer resident Common Life Form 7		NG NSS4 NTMB	Basin-prairie and mountain-foothills shrublands.	Usually nests in or under a sagebrush. Feeds on insects, seeds.	<u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> — <u>O</u> <u>b</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u>
Lark Bunting <i>Calamospiza melanocorys</i> 605.0 LARB Summer resident Abundant Life Form 5		NG NTMB	Basin-prairie and mountain-foothills shrublands, eastern great plains and great basin-foothills grasslands, agricultural areas.	Nests on the ground, usually with the rim of the nest flush with the ground. Feeds on insects, especially grasshoppers, seeds.	<u>O</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Savannah Sparrow <i>Passerculus sandwichensis</i> 542.0 SAVS Summer resident Common Life Form 5		NG NTMB	Willow, wet-moist meadow grasslands, eastern great plains grasslands, marshes, irrigated native meadows.	Nests in a natural or excavated depression on the ground. Feeds on insects, seeds, occasionally snails.	<u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u>
Grasshopper Sparrow <i>Ammodramus savannarum</i> 546.0 GRSP Summer resident Common Life Form 5		NG NSS4 NTMB	Basin-prairie shrublands, eastern great plains grasslands, wet-moist meadow grasslands, agricultural areas.	Nest is sunk in a slight depression on the ground. Feeds on insects, seeds.	<u>O</u> <u>O</u> <u>O</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>B</u> <u>—</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>—</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>—</u> <u>O</u> <u>—</u> <u>O</u> <u>b</u>
Baird's Sparrow <i>Ammodramus bairdii</i> 545.0 BAIS Summer resident Uncommon Life Form 5	(AS)	NG NSS4 NTMB	Shortgrass eastern great plains grasslands.	Nests in a natural or scratched depression on the ground. Feeds on seeds, insects.	<u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u> <u>—</u> <u>O*</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>b*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>b*</u> <u>b*</u>
Le Conte's Sparrow <i>Ammodramus leconteii</i> 548.0 LCSP Migrant Rare Life Form	(AS)	NG	Wet-moist meadow grasslands, marshes.	Feeds on insects, seeds. L4, 5: H. Downing and P. Hall recorded the species several times. L19: J. Lawrence and G. Lawrence, 3 October 1993.	<u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Nelson's Sparrow <i>Ammodramus nelsoni</i> 549.1 NESP Accidental Rare Life Form	(AS)		Marshes, wet-moist meadow grasslands.	Feeds on insects. L8: J. Aukland, 30 June 1999. L19: K.C. Roberts and B. Roberts, 7 October 1993.	— — — — — — — — O* — — — — — — — — — — — — — O* — — — — — — — — — — — —
Fox Sparrow <i>Passerella iliaca</i> 585.0 FOSP Resident Common Life Form 7		NG NTMB	Native riparian shrub with adjacent coniferous forest or woodland-chaparral. Also burned coniferous and logged/thinned forests. Lower habitats during migration and winter.	Usually nests on the ground or in a shrub. Feeds on insects, seeds, berries.	B b O b b — O B O O O — O — B b O — O — O B B O b b O O
Song Sparrow <i>Melospiza melodia</i> 581.0 SOSP Resident Common Life Form 7		NG NTMB	Cottonwood-riparian, native riparian shrub, marshes.	Usually nests on the ground beneath a grass tuft, occasionally in a shrub. Feeds on insects, seeds, some fruit.	B O B B B b B B b B B b b O b b B b B b B B B B b B B b
Lincoln's Sparrow <i>Melospiza lincolnii</i> 583.0 LISP Summer resident Common Life Form 3		NG NTMB	Riparian shrub or marshes, above 7,000 feet. Lower during migration.	Nests in a grass tussock or in a depression on the ground. Feeds on insects, seeds.	B B B B B O O B B B B B O O b b B B b b O b b O B B B O

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Swamp Sparrow <i>Melospiza georgiana</i> 584.0 SWSP Migrant Rare Life Form	(FL)	NG NTMB	Marshes.	Feeds on seeds, insects.	<u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>O*</u> <u>O</u> <u>O*</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>O*</u>
White-throated Sparrow <i>Zonotrichia albicollis</i> 558.0 WTSP Migrant Uncommon Life Form		NG NTMB	Riparian shrub, urban areas.	Feeds on insects, seeds, fruit. Occasionally winters in Wyoming. L5: H. Downing, 23 December 1979, 8-9 October 1991, 30 October 1992. L10: A. Hines and W. Hines, 13 February 1993. L11: J. Allen, 12 April 1992.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u> <u>O*</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u>
Harris's Sparrow <i>Zonotrichia querula</i> 553.0 HASP Winter resident Uncommon Life Form		NG	Deciduous forests, agricultural areas, urban areas.	Feeds on insects, seeds, berries.	<u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u>
White-crowned Sparrow <i>Zonotrichia leucophrys</i> 554.0 WCSP Summer resident Common Life Form 7		NG NTMB	Willow, hawthorn, wild plum, dogwood, water/bog birch, alder.	Nests in a shrub or on the ground. Feeds on insects, seeds, berries.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Golden-crowned Sparrow <i>Zonotrichia atricapilla</i> 557.0 GCSP Accidental Rare Life Form	(AS)		Riparian shrub, cottonwood-riparian.	Feeds on insects, seeds, berries.	— — — — — O* — — — O* — — — — — — — — — — O* — — O* — — — — — — — — —
Dark-eyed Junco <i>Junco hyemalis</i> 567.7 DEJU Resident Common Life Form 5		NG NTMB	Coniferous and deciduous forests. Urban areas during the winter.	Nests on the ground in a shallow depression with overhead protection. Feeds on insects, seeds. Includes White-winged, Slate-colored, Oregon, Pink-sided, and Gray-headed races.	B B O B B O B B B B B O B B B b B B B B O b B B B B B O
Hepatic Tanager <i>Piranga flava</i> 609.0 HETA Accidental Rare Life Form	(AS)	NTMB	Open coniferous and mixed coniferous/oak forests, pine-juniper.	Feeds primarily on insects, occasionally some fruit. L19: O.K. Scott, F. Layton, and L. Layton. L27: American Birds 31:1030 and K. Diem, May 1972. L28: American Birds 35:210.	— O — — — — — — — — — — — — O O
Summer Tanager <i>Piranga rubra</i> 610.0 SUTA Migrant Rare Life Form	(FL)	NG NTMB	Cottonwood-riparian, urban areas.	Feeds on insects, especially bees and wasps, occasionally fruit. L22: F. Luke and others, 26 May 1991. L26: W. Haas, 13 May 1983. L28: J. Cairo, 16 May 1994.	— O — — — — — — — O — — O* — — — — — O — — — — O* — — — O* — — — — O* — — O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Scarlet Tanager <i>Piranga olivacea</i> 608.0 SCTA Accidental Rare Life Form	(AS)	NTMB	Deciduous and mixed deciduous/ coniferous forests.	Feeds on insects, fruits. L2: J. Cicco, 1 October 1999. L26: F. Bergquist, 13 October 2007. L28: J. Cairo, 14 May 1994.	— O* — O* — O*
Western Tanager <i>Piranga ludoviciana</i> 607.0 WETA Summer resident Common Life Form 10		NG NTMB	Coniferous and deciduous forests.	Usually nests in a conifer, in a fork or on a horizontal branch, well out from the trunk. Feeds on insects, fruit.	B B O B B O B B b B B b b O B b B O b b B B b b b B B B
Northern Cardinal <i>Cardinalis cardinalis</i> 593.0 NOCA Migrant Rare Life Form	(AS)	NG	Deciduous forests, urban areas.	Feeds on insects, fruit, seeds.	— — — O O* — O* — O* — — O* — O* — O O* — O* O* O* — — — — — — O
Yellow Grosbeak <i>Pheucticus chrysopleus</i> 594.1 YEGR Accidental Rare Life Form	(AS)		Deciduous forests, woodland- chaparral edge.	Feeds on insects, fruit, seeds. Native to western Mexico, straggler to Arizona in the summer. L8: R. Vaughn, 19 May 2002.	— — — — — — — O* —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting
observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Rose-breasted Grosbeak <i>Pheucticus ludovicianus</i> 595.0 RBGR Summer resident Uncommon Life Form 11	(FL)	NG NTMB	Deciduous forests, urban areas.	Nests in a shrub. Feeds on insects, fruit, seeds, buds. Most observations are of migrants. L2: B. Newell, 9 June 2004. L7: J.R. Flier, 17 May 1989.	<u>O</u> <u>O*</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O*</u> <u>O</u> <u>O</u> <u>O*</u> <u>O*</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>—</u> <u>B</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>O</u> <u>O*</u> <u>O*</u>
Black-headed Grosbeak <i>Pheucticus melanocephalus</i> 596.0 BHGR Summer resident Common Life Form 11		NG NTMB	Aspen and riparian woodlands below 8,000 feet.	Nests in a deciduous tree or shrub; nest is bulky and loosely built. Feeds on insects, seeds, fruit.	<u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u>
Blue Grosbeak <i>Passerina caerulea</i> 597.0 BLGR Summer resident Rare Life Form 11		NG NSS4 NTMB	Cottonwood-riparian, rural developments, urban areas.	Nests in a shrub or tree, occasionally in a vine tangle. Feeds on insects, seeds, some fruit, occasionally snails.	<u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>B*</u> <u>B*</u> <u>O</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>O</u> <u>—</u> <u>B</u> <u>O</u> <u>B</u> <u>—</u> <u>B</u> <u>—</u> <u>B</u> <u>O</u> <u>O</u> <u>O</u> <u>—</u> <u>B</u>
Lazuli Bunting <i>Passerina amoena</i> 599.0 LAZB Summer resident Common Life Form 8		NG NTMB	Pine-juniper, woodland-chaparral, mountain-foothills shrublands with aspen, cottonwood-riparian.	Nests in a shrub or vine tangle. Feeds on insects, seeds.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>O</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b																												
Indigo Bunting <i>Passerina cyanea</i> 598.0 INBU Summer resident Uncommon Life Form 7	(FL)	NG NTMB	Willow, mixed riparian shrub.	Nests in a shrub, tree, or vine tangle. Feeds on insects, seeds, fruit. A few records include hybrids with Lazuli Bunting.	<table style="border: none; width: 100%; text-align: center;"> <tr><td>—</td><td>O*</td><td>O</td><td>B</td><td>O*</td><td>—</td><td>B</td></tr> <tr><td>O</td><td>O</td><td>O*</td><td>O*</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>O</td><td>—</td><td>O*</td><td>O</td><td>O</td></tr> <tr><td>—</td><td>O*</td><td>—</td><td>O</td><td>O</td><td>O*</td><td>O</td></tr> </table>	—	O*	O	B	O*	—	B	O	O	O*	O*	—	—	—	—	—	O	—	O*	O	O	—	O*	—	O	O	O*	O
—	O*	O	B	O*	—	B																											
O	O	O*	O*	—	—	—																											
—	—	O	—	O*	O	O																											
—	O*	—	O	O	O*	O																											
Painted Bunting <i>Passerina ciris</i> 601.0 PABU Accidental Rare Life Form	(AS)	NTMB	Deciduous forest edge, riparian shrub.	Feeds primarily on seeds, some insects. L7: E. Miller, 14-18 May 1991. L9: R. Moldenhauer, 11 October 2003. L19: G. Lawrence, 8 May 2000. L28: M. Hanesworth, 23-26 May 1951, 4 June 1975, American Birds 29:1010.	<table style="border: none; width: 100%; text-align: center;"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td></tr> <tr><td>—</td><td>O*</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O*</td></tr> </table>	—	—	—	—	—	—	O*	—	O*	—	—	—	—	—	—	—	—	—	O*	—	—	—	—	—	—	—	—	O*
—	—	—	—	—	—	O*																											
—	O*	—	—	—	—	—																											
—	—	—	—	O*	—	—																											
—	—	—	—	—	—	O*																											
Dickcissel <i>Spiza americana</i> 604.0 DICK Summer resident Uncommon Life Form 5	(FL)	NG NSSU NTMB	Eastern great plains grasslands.	Nests on the ground in a clump of grass; nest is bulky. Feeds on insects, seeds. L2: M. Klein, 14-17 May 2002. L23: A. Halvorson, 20 April 1994.	<table style="border: none; width: 100%; text-align: center;"> <tr><td>—</td><td>O*</td><td>O</td><td>O</td><td>B</td><td>O</td><td>B</td></tr> <tr><td>O</td><td>—</td><td>O</td><td>—</td><td>—</td><td>O</td><td>O</td></tr> <tr><td>—</td><td>—</td><td>O</td><td>O</td><td>O</td><td>O</td><td>B</td></tr> <tr><td>—</td><td>O*</td><td>—</td><td>—</td><td>O</td><td>O</td><td>O</td></tr> </table>	—	O*	O	O	B	O	B	O	—	O	—	—	O	O	—	—	O	O	O	O	B	—	O*	—	—	O	O	O
—	O*	O	O	B	O	B																											
O	—	O	—	—	O	O																											
—	—	O	O	O	O	B																											
—	O*	—	—	O	O	O																											
Bobolink <i>Dolichonyx oryzivorus</i> 494.0 BOBO Summer resident Uncommon Life Form 5	(FL)	NG NSS4 NTMB	Basin-prairie shrublands, eastern great plains grasslands, great basin-foothills grasslands, alfalfa, irrigated native and introduced meadows.	Nests on the ground in a natural or scraped depression with a dense cover of forbs. Feeds primarily on insects, seeds.	<table style="border: none; width: 100%; text-align: center;"> <tr><td>O</td><td>B</td><td>—</td><td>B</td><td>B</td><td>O</td><td>b</td></tr> <tr><td>B</td><td>O</td><td>b</td><td>O</td><td>O</td><td>O</td><td>—</td></tr> <tr><td>b</td><td>O</td><td>B</td><td>O</td><td>O</td><td>O</td><td>O</td></tr> <tr><td>O</td><td>b</td><td>O</td><td>b</td><td>O</td><td>O</td><td>b</td></tr> </table>	O	B	—	B	B	O	b	B	O	b	O	O	O	—	b	O	B	O	O	O	O	O	b	O	b	O	O	b
O	B	—	B	B	O	b																											
B	O	b	O	O	O	—																											
b	O	B	O	O	O	O																											
O	b	O	b	O	O	b																											

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Red-winged Blackbird <i>Agelaius phoeniceus</i> 498.0 RWBL Summer resident Abundant Life Form 3		NG NTMB	Riparian shrub, marshes, agricultural areas.	Nests near or over water, usually in emergent vegetation. Feeds on insects, seeds. A few birds may remain during the winter.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Eastern Meadowlark <i>Sturnella magna</i> 501.0 EAME Accidental Rare Life Form	(AS)		Grasslands, agricultural areas.	Feeds on insects, seeds. L28: C. Michelson, 16 May 2008.	— O*
Western Meadowlark <i>Sturnella neglecta</i> 501.1 WEME Summer resident Abundant Life Form 5		NG NTMB	Basin-prairie and mountain-foothills shrublands and grasslands below 8,000 feet, agricultural areas.	Nests on the ground in a natural or scraped depression. Feeds on insects, seeds.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Yellow-headed Blackbird <i>Xanthocephalus xanthocephalus</i> 497.0 YHBL Summer resident Common Life Form 5		NG NTMB	Marshes.	Nests over water in emergent vegetation. Feeds on insects, seeds.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Rusty Blackbird <i>Euphagus carolinus</i> 509.0 RUBL Migrant Uncommon Life Form	(AS)	NG	Cottonwood-riparian, agricultural areas, urban areas.	Feeds on insects, seeds, small vertebrates, some invertebrates.	O+ _ _ O O* O* O* O* _ _ O _ O _ O* _ O* O* _ O _ O* _ _ O _ O
Brewer's Blackbird <i>Euphagus cyanocephalus</i> 510.0 BRBL Summer resident Abundant Life Form 3 6 7 11		NG NTMB	Deciduous forests, basin-prairie and mountain-foothills shrublands, riparian shrub, grasslands, marshes, agricultural areas, urban areas.	Nests in a variety of locations; typically in a tree or shrub or just above the ground. Feeds on insects, seeds, fruit, some invertebrates. A few birds remain during the winter.	B B B B B b B B B B B b B b B B B B B B B B B B B B B B
Common Grackle <i>Quiscalus quiscula</i> 511.0 COGR Summer resident Common Life Form 3 5 7 11 14		NG	Cottonwood-riparian, basin-prairie shrublands, willow/mixed riparian shrub, grasslands, marshes, agricultural areas, urban areas.	Nests in a deciduous tree or shrub, often near water. Feeds on insects, small vertebrates, some invertebrates, fruit, seeds.	O B B B B b B O B B O b B O B B B O B B B B B O B B B B
Great-tailed Grackle <i>Quiscalus mexicanus</i> 512.0 GTGR Accidental Rare Life Form	(FL)		Grasslands with scattered trees, agricultural areas, riparian shrub, marsh-swamp wetlands, urban areas.	Feeds on insects, lizards, aquatic invertebrates, aquatic vertebrates, fruit, grain, other foods.	_ _ _ _ _ O* _ _ _ _ _ _ _ _ O* O* _ _ O* _ _ _ O* O* O*

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Brown-headed Cowbird <i>Molothrus ater</i> 495.0 BHCO Summer resident Common Life Form 5 7 11 14		NG NTMB	Cottonwood-riparian, juniper, basin-prairie and mountain-foothills shrublands, riparian shrub, grasslands, agricultural areas, urban areas.	Does not build its own nest or raise its own young; lays eggs in nests of other birds. Feeds on insects, seeds. Can be locally detrimental to songbird populations.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Orchard Oriole <i>Icterus spurius</i> 506.0 OROR Summer resident Uncommon Life Form 9	(FL)	NG NTMB	Cottonwood-riparian, rural developments, urban areas.	Nests in a deciduous tree; nest is usually suspended from a forked terminal twig. Feeds on insects, fruit. L5: first breeding record H. Downing, J. Collins, and N. Collins, 8 July 1987.	<u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>B*</u> <u>B</u> <u>O*</u> <u>O</u> <u>O</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>b</u> <u>O</u> <u>B</u> <u>—</u> <u>O</u> <u>—</u> <u>—</u> <u>—</u> <u>B</u> <u>B</u>
Bullock's Oriole <i>Icterus bullockii</i> 508.0 BUOR Summer resident Common Life Form 9		NG NTMB	Cottonwood-riparian, cottonwood-dryland, rural developments, urban areas.	Nests in a deciduous tree; nest is usually hung from a drooping branch. Feeds on insects, fruit, nectar.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Baltimore Oriole <i>Icterus galbula</i> 507.0 BAOR Accidental Rare Life Form	(AS)	NTMB	Cottonwood-riparian, cottonwood-dryland, rural developments, urban areas.	Feeds on insects, fruit, nectar.	<u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O*</u> <u>—</u> <u>O*</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>—</u> <u>O*</u> <u>O*</u> <u>O*</u> <u>—</u>

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
(FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
= accepted by the Yellowstone Bird Observation Committee.
+ = accepted by both the WBRC and YBOC.

NAME AND STATUS	WBRC^a	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^b
Scott's Oriole <i>Icterus parisorum</i> 504.0 SCOR Summer resident Rare Life Form 10	(AS)	NG NSSU NTMB	Arid habitats dominated by juniper-sagebrush or yucca.	Nests in a juniper, toward the end of a branch. Feeds on insects, fruit, nectar. L17: E. Crane, 26 May 2004. L23, 24, 25: S. Findholt and S. Fitton, 1982, Western Birds 14:109-110.	— — — — — — — — — — — — — — O* — — — — — B b B — — —

^a (AS) = report all sightings to the Wyoming Bird Records Committee.
 (FL) = report the first sighting in each latilong and all nesting observations to the Wyoming Bird Records Committee.

^b * = accepted by the Wyoming Bird Records Committee.
 # = accepted by the Yellowstone Bird Observation Committee.
 + = accepted by both the WBRC and YBOC.

Mammals

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Virginia Opossum <i>Didelphis virginiana</i> 800.0 DIVI Accidental Rare Life Form	NG	Cottonwood-riparian, cattail wetlands, agricultural areas, roadside/railroad banks, urban areas.	Nests on the ground, possibly under a log or boulder. Feeds on fruit, berries, carrion, bird eggs, frogs, crayfish, worms, insects. Common east of Wyoming and in the northwest U.S.	— — <u>O</u> — — — — — — — — <u>O</u> — — <u>O</u> — — — — <u>O</u> — — <u>B</u> <u>B</u> <u>O</u> <u>O</u> — — — — <u>O</u> <u>O</u>
Masked Shrew <i>Sorex cinereus</i> 801.0 SOCI Resident Common Life Form 5 15	NG	Damp coniferous forests, aspen, cottonwood-riparian, mountain-foothills shrublands, riparian shrub, eastern great plains and moist meadow grasslands, mixed alpine meadows, marsh-swamp wetlands, rock outcrops, talus fields.	Nests on the ground or in a burrow. Feeds on insects, beetles, moths, flies, insect larvae, crickets, spiders, small vertebrates, only rarely on vegetable matter.	<u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> — — — — <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> — — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — —
Hayden's Shrew <i>Sorex haydeni</i> 801.1 SOHA Resident Rare Life Form 5 15	NG NSS4	Riparian shrub, mixed grass-like meadows, marsh-swamp wetlands, coniferous forests. May be restricted to the Black Hills region.	Nests under a log, rock, or in a crevice. Feeds on beetles, grasshoppers, spiders, earthworms, insect larvae, small vertebrates. Access to open water is important. Taxonomically similar to the Masked Shrew, although presently considered distinct.	— — — — <u>b</u> — — <u>B</u> —
American Pygmy Shrew <i>Sorex hoyi</i> 806.0 SOHO Resident Rare Life Form 15	NG NSS2	Coniferous forests, mountain-foothills grasslands, mixed grass/forb meadows, bogs, wet areas.	Nests in old, decaying logs and in the roots of tree stumps. Feeds on a variety of arthropods, worms, insects. L26: specimens collected by Brown (1967) in the Medicine Bow Mountains are a disjunct population separate from the boreal range to the north.	— <u>B</u> — —

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Merriam's Shrew <i>Sorex merriami</i> 805.0 SOME Resident Rare Life Form 15	NG	Basin-prairie and mountain-foothills shrublands, eastern great plains grasslands, rural developments, barren areas, below 7,550 feet.	Nests in a burrow. Feeds on spiders, adult and larval beetles, crickets, larval moths, butterflies, wasps, caterpillars in warmer months. Widespread species in the western U.S.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Dusky Shrew <i>Sorex monticolus</i> 807.0 SOMO Resident Common Life Form 5 15	NG	Englemann spruce-subalpine fir, lodgepole pine, Douglas fir, aspen, cottonwood-riparian, sagebrush-grasslands, willow, grasslands, mixed grass/forb meadows, rock outcrops, talus fields.	Nests on the ground or in a burrow. Feeds on insects, spiders, earthworms, some plants. Forages in ground litter and under vegetation, rocks, logs. Uses echolocation to locate prey.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Dwarf Shrew <i>Sorex nanus</i> 803.0 SONA Resident Rare Life Form 15	NG NSS3	Coniferous forests, aspen, mountain-foothills shrublands, alpine grasslands, mixed alpine meadows, rock outcrops, talus fields.	Nests in a burrow. Feeds on carrion of small mammals, spiders, soft-bodied insects.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Western Water Shrew <i>Sorex navigator</i> 804.0 SONV Resident Common Life Form 3 16	NG	Coniferous forests, riparian shrub, alpine grasslands, marsh-swamp wetlands, near standing water.	Nests in a stream bank; nest is lined with moss or grass. Feeds on beetles, insects, flies, caddisflies, mayflies, snails, leeches, planarians, tadpoles, small fish.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Preble's Shrew <i>Sorex preblei</i> 804.1 SOPR Resident Rare Life Form 15	NG NSS3	Marsh grass, creeks and bogs bordered by willow or riparian shrub, occasionally wetter areas of open conifer stands, in association with mountain-foothills grasslands.	Nests in a burrow. Food habits are unknown. Most likely feeds on animal material, insects. L1: the first Wyoming specimen collected in Lamar Valley, Yellowstone National Park at 6,000 feet. Recent specimens from Snake River Canyon.	B O _ _ _ _ _ O _ _ _ _ _ _ _ _ _ _ _ _ _ h _ _ _ _ _ _
Vagrant Shrew <i>Sorex vagrans</i> 802.0 SOVA Resident Rare Life Form 5 15	NG	Riparian shrub, moist meadow grasslands, bogs and other riparian or marsh habitats with moist soil. Prefers leaf litter, rotting logs. Also found in coniferous forests, sagebrush-grasslands, mixed shrublands.	Nests on the ground or in a burrow. Feeds on insects, spiders, earthworms, some plants. Forages in ground litter and under vegetation, rocks, logs. Uses echolocation to locate prey.	h O B B B _ B B B b _ _ _ b b b _ _ b _ h b O _ _ b h
Eastern Mole <i>Scalopus aquaticus</i> 810.0 SCAQ Resident Abundance unknown Life Form 15	NG	Eastern great plains grasslands, primarily small grain agricultural areas, urban areas.	Nests in its underground tunnel system; nest is lined with dry leaves. Feeds on insects, soil invertebrates, some plant material. Mainly distributed east of Wyoming.	_ _ _ _ b _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ B _ _ _ _ _ _ B
California Myotis <i>Myotis californicus</i> 815.1 MYCA Peripheral Rare Life Form 4 17	NG	Pine-juniper, woodland-chaparral, basin-prairie shrublands, grasslands, cliffs, rock outcrops.	Roosts in any type of shelter including mines, caves. May hibernate only briefly, possibly in Wyoming. Feeds on flying insects including beetles, moths. L3: recent specimens from Bighorn Canyon and Powell areas.	_ _ O _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ a a _ O a a a a _

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Western Small-footed Myotis <i>Myotis ciliolabrum</i> 816.0 MYCI Resident Uncommon Life Form 4 17	NG NSS4	Pine-juniper; basin-prairie shrublands; sagebrush-grasslands; eastern great plains, great basin-foothills, and mountain-foothills grasslands; cliffs; rock outcrops.	Often roosts and has nurseries in buildings. Hibernates in caves, mines, probably in Wyoming. Feeds on flying insects, especially beetles. Extremely sensitive to human disturbance during hibernation and in nurseries.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Long-eared Myotis <i>Myotis evotis</i> 818.0 MYEV Resident Uncommon Life Form 4 14 17	NG NSS4	Coniferous forests, especially ponderosa pine and juniper; cottonwood-riparian; basin-prairie shrublands; sagebrush-grasslands.	Roosts in caves, buildings, mine tunnels. May hibernate in Wyoming. Feeds primarily on beetles, moths. Extremely sensitive to human disturbance during hibernation.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Northern Long-eared Myotis <i>Myotis septentrionalis</i> 819.0 MYSE Residency unknown Peripheral Rare Life Form 4 14 17	NG NSS3 Thr.	Ponderosa pine, mixed coniferous/deciduous forests, often dense forests.	Roosts in caves, tree cavities. May hibernate in caves, mines in Wyoming. Feeds on flying insects on hillsides and near ridgetops. Extremely sensitive to human disturbance during hibernation. Classified as a federally Threatened species in 2015.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Little Brown Myotis <i>Myotis lucifugus</i> 815.0 MYLU Resident Common Life Form 4 14 17	NG NSS4	Coniferous and deciduous forests, sagebrush-grasslands, riparian shrub, urban areas, close to water.	Day roosts in humid caves, buildings; frequently night roosts in buildings. May hibernate in caves, mines in Wyoming. Feeds over water on aquatic insects. Extremely sensitive to human disturbance during hibernation.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Hoary Bat <i>Lasiurus cinereus</i> 822.0 LACI Summer resident Uncommon Life Form 11	NG	Coniferous and deciduous forests, aspen, basin-prairie shrublands, eastern great plains grasslands, sagebrush-grasslands, mountain foothills shrublands, riparian shrub, urban areas.	Roosts in deciduous trees; rarely uses buildings. Feeds primarily on moths; also beetles, flies, grasshoppers, wasps. Probably migrates to a southern climate for the winter.	_ B B B B O B O O O O B h B B O B B B B B O O O O B O B
Silver-haired Bat <i>Lasionycteris noctivagans</i> 820.0 LANO Summer resident Uncommon Life Form 4 14 17	NG	Coniferous and deciduous forests, often near water.	Roosts in tree foliage, hollow trees, mines, caves, houses, under loose bark. Most likely feeds on moths, insects, carab beetles, flies, caddisflies, occasionally spiders.	_ B B B B O B B B O O B _ B B _ B O B B B O O a O B O O
American Perimyotis (Tri-colored Bat) <i>Perimyotis subflavus</i> 820.1 PESU Accidental Rare Life Form 4 14 17	NG	Open wooded areas and along woodland borders.	Roosts in caves, mines, rock crevices, tree foliage, buildings. Hibernates in caves, mines, buildings. Feeds on leafhoppers, beetles, flies, moths, ants.	_ O _ _ _ _ _ _ _
Big Brown Bat <i>Eptesicus fuscus</i> 825.0 EPFU Resident Common Life Form 4 14 17	NG	Coniferous and deciduous forests, basin-prairie and mountain-foothills shrublands, grasslands, urban areas.	Roosts and hibernates in buildings, occasionally in rock crevices, caves, hollow trees, crevices in bark. Feeds primarily on beetles; also other insects, very few moths. Forages over open meadows, city streets, corrals, farmsteads.	_ O B B O O B B O B O O h B B O B B B B B O O O B B O O

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Spotted Bat <i>Euderma maculatum</i> 824.0 EUMA Summer resident Rare Life Form 4 17	NG NSS4	Associated with a variety of habitat types over their range, but known only from juniper shrublands and desert sagebrush-grasslands in Wyoming. Cliffs over perennial water are an important habitat component.	Roosts in rock crevices. Feeds primarily on moths, occasionally grasshoppers, beetles. Maternity roosts are extremely sensitive to human disturbance. L3: two specimens recently collected from Big Horn County. Very rare species in North America.	_ _ O b O _ _ _ _ _ _ _ O B O _ _ _ _ _ _ _ _ _ _ _ _ _ _ O a _ _ _ _ _
Townsend's Big-eared Bat <i>Corynorhinus townsendii</i> 823.0 COTO Resident Uncommon Life Form 4 17	NG NSS3	Deciduous forests, dry coniferous forests, basin-prairie and mountain-foothills shrublands, desert grasslands, juniper.	Day roosts in caves, mines, rock outcrops; night roosts in buildings. Hibernates in caves, probably in Wyoming. Feeds primarily on moths, occasionally flies, beetles. Extremely sensitive to human disturbance during hibernation.	h B B B _ _ a O O B O O _ O _ O O O B O B _ O _ a a B _
Pallid Bat <i>Antrozous pallidus</i> 827.0 ANPA Summer resident Rare Life Form 4 11 17	NG NSS3	Sagebrush-grasslands, cliffs, rock outcrops, cottonwood-riparian, eastern great plains grasslands.	Day roosts in small crevices; night roosts in open buildings, rock overhangs, porches. Feeds on a variety of terrestrial vertebrates and invertebrates. Extremely sensitive to human disturbance during roosting.	_ a B B _ _ a a a B B _ _ _ a _ O a O a B _ O _ O a O _
Brazilian Free-tailed Bat <i>Tadarida brasiliensis</i> 828.0 TABR Accidental Rare Life Form 4	NG	Deciduous forests, woodland-chaparral, riparian shrub, caves, cliffs.	Roosts in large colonies in caves. Feeds primarily on small moths. Has a higher probability of transmitting rabies to humans than any other species due to its colonial nature.	_ _ O _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ O _ _ _ _ _ _ _ _ _ _ _ _ _ O

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Big Free-tailed Bat <i>Nyctinomops macrotis</i> 829.0 NYMA Accidental Rare Life Form 4	NG	Lowland and highland habitats. Prefers cliffs, crevices, occasionally buildings.	Roosts primarily in crevices of rocky cliffs; has been reported roosting in buildings. Feeds primarily on small moths; also other insects.	— — — — — — — — <u>O</u> —
American Pika <i>Ochotona princeps</i> 830.0 OCPR Resident Common Life Form 4	NG NSS2	Talus fields and outcrops of shattered rock near grass or forb meadows in alpine grasslands.	Nests between rocks or in burrows in talus slopes. Feeds on grasses, sedges, forbs, conifer twigs, aspen, lichens. Classified as a protected species in Wyoming.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> — — — — <u>B</u> <u>B</u> <u>b</u> — — — — — — <u>B</u> <u>B</u> <u>B</u> — — — — — — <u>b</u> <u>b</u> — — <u>B</u> <u>B</u> <u>b</u> — —
Pygmy Rabbit <i>Brachylagus idahoensis</i> 837.0 BRID Resident Rare Life Form 5	NG NSS3	Dense, tall stands of big sagebrush, usually along intermittent streams or riparian areas in sagebrush-grasslands.	Nests on the ground, most likely under a sagebrush, or in a burrow. Ninety-nine percent of the diet is sagebrush; also feeds on some grasses in mid- to late summer. Overgrazing by livestock may impact habitat values.	— — — — — — — — — — — — — — — — <u>b</u> <u>b</u> <u>b</u> — — — — — — <u>B</u> <u>B</u> <u>b</u> <u>b</u> — — — —
Desert Cottontail <i>Sylvilagus audubonii</i> 833.0 SYAU Resident Common Life Form 5	SG	Cottonwood-riparian, cottonwood-dryland, juniper, basin-prairie shrublands, sagebrush-grasslands, eastern great plains and great basin-foothills grasslands.	Nests on the ground; nest is lined with fur. Feeds on grasses, herbaceous vegetation, occasionally buds or bark from shrubs. Difficult to distinguish from other <i>Sylvilagus</i> species.	<u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — <u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Eastern Cottontail <i>Sylvilagus floridanus</i> 834.0 SYFL Resident Uncommon Life Form 5	SG	Heavy shrub cover in cottonwood-riparian, cottonwood-dryland, eastern great plains grasslands.	Nests on the ground. Feeds on virtually all green vegetation; also buds and bark of woody vegetation in the winter. May be more widespread than indicated; difficult to distinguish from other <i>Sylvilagus</i> species.	-- B -- -- -- -- -- h h
Mountain Cottontail <i>Sylvilagus nutallii</i> 835.0 SYNU Resident Common Life Form 5	SG	Heavy shrub cover in deciduous forests, juniper, woodlands.	Nests on the ground; nest is fur-lined and covered. Feeds on green grasses; also buds and bark of sagebrush, juniper, and other woody shrubs in the winter. Difficult to distinguish from other <i>Sylvilagus</i> species.	-- h h -- h B h B h h B -- b -- -- h B h h h h B h h O h B --
Snowshoe Hare <i>Lepus americanus</i> 836.0 LEAM Resident Common Life Form 5	SG	Coniferous forests, aspen, willow.	Nests in a shallow, fur-lined depression in a dense thicket, frequently in willows. Feeds on grasses, herbs, willows in the summer; bark and twigs of deciduous trees and shrubs, occasionally buds and bark of conifers in the winter.	B B -- B B -- -- B B -- b -- -- -- B B B h -- b -- B b -- B B B --
Black-tailed Jackrabbit <i>Lepus californicus</i> 832.0 LECA Resident Common Life Form 5	PD	Eastern shortgrass prairies, open greasewood and sagebrush grasslands, moist meadow grasslands, agricultural areas, roadside/railroad banks, unreclaimed mine areas.	Nests in a simple depression; nest is sometimes fur-lined. Feeds on virtually all green plants, grasses and small forbs are preferred; any exposed vegetation is eaten in the winter.	-- -- b b -- B B -- -- -- -- -- B B b -- -- -- B B B B -- -- -- b B B

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
White-tailed Jackrabbit <i>Lepus townsendii</i> 831.0 LETO Resident Common Life Form 5	PD	Occurs in most open habitats from shrub grasslands, to openings in foothill conifer stands and montane coniferous forests, to alpine tundra. Sagebrush-grasslands are most commonly used.	Nests in dense vegetation; nest is shallow, lined with fur and dry leaves, well hidden. Feeds on virtually all green, succulent vegetation in the summer; buds and bark of exposed vegetation in the winter.	— <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Yellow-pine Chipmunk <i>Tamias amoenus</i> 841.0 TAAM Resident Uncommon Life Form 15	NG NSS4	Coniferous forests, especially lodgepole pine; mixed aspen; juniper; occasionally sagebrush-grasslands, willow, and mountain-foothills grasslands near forested habitats.	Nests in a burrow. Feeds mostly on seeds, nuts, buds, roots, bulbs, fungi; also insects such as larvae, grasshoppers, beetles.	<u>B</u> <u>b</u> — — — — — <u>B</u> <u>b</u> <u>B</u> — — — — — <u>b</u> <u>B</u> — — — — — <u>B</u> — — — — —
Cliff Chipmunk <i>Tamias dorsalis</i> 842.0 TADO Resident Rare Life Form 15	NG NSS3	Steep, rocky hillsides; rock outcrops; talus fields; in pine-juniper and juniper.	Nests deep in a crevice of a cliff. Feeds on a wide variety of flowers, stems, seeds. In Wyoming, probably restricted to rock outcrops along Flaming Gorge Reservoir.	— <u>B</u> <u>B</u> — — — — —
Least Chipmunk <i>Tamias minimus</i> 840.0 TAMI Resident Abundant Life Form 15	NG	Occurs in most habitat types. Inhabits more biotic communities and has a greater altitudinal range than other chipmunk species whose range it overlaps.	Nests in a burrow. Feeds on green plant parts, roots, nuts, fruit, berries, fungi; also arthropods such as beetles, grasshoppers, caterpillars.	<u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Uinta Chipmunk <i>Tamias umbrinus</i> 843.0 TAUM Resident Uncommon Life Form 15	NG NSS4	Coniferous forest, aspen, rocky slopes, logged areas and clearings near forests. Hibernacula include logs, rock crevices, burrows, and trees.	Nests in a burrow. Feeds on a wide variety of seeds; also buds, tender green shoots when seeds are unavailable. Generally a tree dweller.	h h _ _ _ _ B b B _ _ _ B h b _ _ _ B _ _ _ B h _
Yellow-bellied Marmot <i>Marmota flaviventris</i> 844.0 MAFL Resident Common Life Form 15	NG	Rock outcrops; talus fields; roadside/railroad banks in coniferous forests, aspen, cottonwood-riparian, juniper, sagebrush-grasslands, mountain-foothills shrublands, riparian shrub, grasslands.	Nests in a burrow. Feeds on a wide variety of foods including grasses, flowers, forbs.	_ B _ B B B B B h B B B B B B B B B B B B B B B
Uinta Ground Squirrel <i>Urocitellus armatus</i> 846.0 URAA Resident Uncommon Life Form 15	NG	Open areas in coniferous forests, pine-juniper, aspen, cottonwood-riparian, juniper; sagebrush-grasslands; mountain-foothills grasslands; irrigated native meadows.	Nests in a burrow. Feeds on grasses, forbs, shrubs, mushrooms, insects; also carrion including roadkills of other Uinta ground squirrels.	O h _ _ _ _ B B B _ _ _ B B _ _ _ _ B B B _ _ B
Wyoming Ground Squirrel <i>Urocitellus elegans</i> 845.0 UREL Resident Common Life Form 15	NG	Aspen, juniper, basin-prairie and mountain-foothills shrublands, riparian shrub, grasslands, agricultural areas, rock outcrops, disturbed areas.	Nests in a burrow. Feeds mainly on plant material; also animal tissue, seed fragments, arthropods including grasshoppers, beetles, flies, bees.	b b _ _ _ _ B B B B B _ b B B B B B B B B B B B

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Golden-mantled Ground Squirrel <i>Callospermophilus lateralis</i> 849.0 CALT Resident Common Life Form 15	NG	Rock outcrops, rock slides, and open areas in coniferous forests; aspen; cottonwood-riparian; juniper; sagebrush-grasslands; mountain-foothills shrublands; riparian shrub; mountain-foothills grasslands.	Nests in a burrow. Feeds primarily on leaves, buds, seeds, nuts, roots, bulbs, fruit, mushrooms; also bird eggs, insects, carrion.	B B _ _ _ _ _ B B _ _ _ _ _ B B B b B h B B B B B B B B
Spotted Ground Squirrel <i>Xerospemophilus spilosoma</i> 847.0 XESP Resident Rare Life Form 15	NG NSS4	Sagebrush-grasslands, eastern great plains grasslands, disturbed areas, sand dunes. Frequents areas with deep, sandy soils.	Nests in a burrow. Feeds on seeds, green plant parts, especially forbs, occasionally carnivorous. Mainly distributed southeast of Wyoming.	_ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ _ B B _ _ _ _ _ B
Thirteen-lined Ground Squirrel <i>Ictidomys tridecemlineatus</i> 848.0 ICTR Resident Common Life Form 15	NG	Juniper, basin-prairie and mountain-foothills shrublands, grasslands, small grain agricultural areas, barren areas, roadside/railroad banks, mined areas, overgrazed areas.	Nests in a burrow. Feeds on grasses, seeds, insects, occasionally birds, young cottontails, lizards, snakes.	_ _ _ B B B B _ O b B B B B _ B b B B B B B B B B B B B
White-tailed Prairie Dog <i>Cynomys leucurus</i> 851.0 CYLE Resident Common Life Form 15	NG NSS4	Basin-prairie and mountain-foothills shrublands, sagebrush-grasslands, shortgrass and midgrass grasslands.	Nests in a burrow. Feeds on green plant parts, seeds, roots, flowers, insects, occasionally carrion.	_ B B h _ _ _ _ B

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION																												
Black-tailed Prairie Dog <i>Cynomys ludovicianus</i> 850.0 CYLU Resident Common Life Form 15	NG NSS4	Shortgrass and midgrass grasslands.	Nests in a burrow. Feeds on grasses, forbs, shrubs, occasionally insects.	<table border="0"> <tr><td>—</td><td>h</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>B</td><td>B</td></tr> </table>	—	h	B	B	B	B	B	—	—	—	B	B	B	B	—	—	—	—	B	B	B	—	—	—	—	—	B	B
—	h	B	B	B	B	B																										
—	—	—	B	B	B	B																										
—	—	—	—	B	B	B																										
—	—	—	—	—	B	B																										
Abert's Squirrel <i>Sciurus aberti</i> 855.0 SCAB Resident Rare Life Form 9	NG NSS4	Ponderosa pine savannah. Peripheral	Nests on a ponderosa pine branch; nest is made of twigs. Feeds primarily on ponderosa pine parts; also fungi, carrion, antlers, bones.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>O</td><td>B</td></tr> </table>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	O	B
—	—	—	—	—	—	—																										
—	—	—	—	—	—	—																										
—	—	—	—	—	—	—																										
—	—	—	—	—	O	B																										
Eastern Gray Squirrel <i>Sciurus carolinensis</i> 856.0 SCCA Resident Abundance unknown Life Form 9	SG	Deciduous forests in urban areas.	Nests in a leaf nest or cavity in a tree. Feeds primarily on nuts; also seeds, fruit, buds, leaves, flowers, insects, occasionally bird eggs and nestlings. Mainly distributed east of Wyoming.	<table border="0"> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>b</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> <tr><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td><td>—</td></tr> </table>	—	—	—	—	b	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—							
—	—	—	—	b	—	—																										
—	—	—	—	—	—	—																										
—	—	—	—	—	—	—																										
Eastern Fox Squirrel <i>Sciurus niger</i> 852.0 SCNI Resident Common Life Form 9	SG	Deciduous forests, woodland-chaparral, cottonwood-riparian, usually near agricultural lands; urban areas.	Nests in a leaf nest in a tree. Feeds on buds of ornamental trees, fruit, nuts, tree seeds, insects, mushrooms, corn.	<table border="0"> <tr><td>—</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>—</td><td>—</td><td>B</td><td>—</td><td>—</td><td>b</td><td>B</td></tr> <tr><td>—</td><td>—</td><td>B</td><td>—</td><td>B</td><td>B</td><td>B</td></tr> <tr><td>B</td><td>—</td><td>—</td><td>B</td><td>B</td><td>B</td><td>B</td></tr> </table>	—	B	B	B	B	B	B	—	—	B	—	—	b	B	—	—	B	—	B	B	B	B	—	—	B	B	B	B
—	B	B	B	B	B	B																										
—	—	B	—	—	b	B																										
—	—	B	—	B	B	B																										
B	—	—	B	B	B	B																										

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Red Squirrel <i>Tamiasciurus hudsonicus</i> 854.0 TAHU Resident Common Life Form 10	SG	Coniferous forests, aspen.	Nests in a leaf nest or cavity in a tree. Feeds on evergreen terminal buds and seeds; also fungi, insects, old bones, occasionally small birds.	B B h b B _ B B B B b _ _ _ B B B B B B _ B B _ B B B _
Northern Flying Squirrel <i>Glaucomys sabrinus</i> 853.0 GLSA Resident Uncommon Life Form 14	NG NSS4	Coniferous forests.	Nests in a cavity or leaf nest in a tree. Feeds primarily on lichens, fungi; also conifer cones, fruit, buds, arthropods, bird eggs, nestling birds.	B h _ _ _ _ B B B b _ _ _ b B B B _ _ _ _ _ B _ _ _ _ B
Wyoming Pocket Gopher <i>Thomomys clusius</i> 862.0 THCL Resident Uncommon Life Form 15	NG NSS2	Upland drier ridge tops, gravelly loose soils, greasewood.	Most likely nests in a maternal burrow, similar to the Northern Pocket Gopher. Feeds on roots and plant parts of forbs, grasses, herbs. Usually feeds underground in a shallow tunnel, pulling roots and plants underground.	_ B B _ _ _
Idaho Pocket Gopher <i>Thomomys idahoensis</i> 863.0 THID Resident Uncommon Life Form 15	NG NSS3	Shallow, stony soils in sagebrush, sagebrush-grasslands, and mountain meadows.	Most likely nests in a maternal burrow, similar to the Northern Pocket Gopher. Feeds on roots and plant parts of forbs, grasses, herbs. Usually feeds underground in a shallow tunnel, pulling roots and plants underground.	_ _ _ _ _ _ _ _ _ _ _ _ _ _ h h _ _ _ _ _ B _ _ _ _ _ _

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Northern Pocket Gopher <i>Thomomys talpoides</i> 860.0 THTA Resident Common Life Form 15	NG	Virtually all habitat types with loose soil and relatively few rocks and stones. Prefers deeper soils.	Nests in a maternal burrow. Feeds on roots and plant parts of forbs, grasses, herbs. Usually feeds underground in a shallow tunnel, pulling roots and plants underground.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>h</u> <u>B</u> <u>O</u> <u>b</u> <u>h</u> <u>b</u> <u>B</u> <u>B</u> <u>h</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u>
Sand Hills Pocket Gopher <i>Geomys lutescens</i> 861.0 GELU Resident Uncommon Life Form 15	NG NSS3	Sagebrush-grasslands, eastern great plains grasslands, agricultural areas. Prefers deep, sandy soils.	Nests in a burrow. Feeds on grasses, herbs, roots, underground plant parts.	— — — — — <u>h</u> — — — — — <u>B</u> <u>h</u> — — — — — <u>h</u> <u>b</u> <u>B</u> — — — — — <u>B</u>
Olive-backed Pocket Mouse <i>Perognathus fasciatus</i> 865.0 PEFS Resident Common Life Form 15	NG NSS4	Arid and semiarid upland habitat, including basin-prairie shrublands, sagebrush-grasslands, eastern great plains and great basin-foothills grasslands. May also be found in farmland and semi-wooded areas. Prefers sandy soils.	Nests in a burrow. Feeds on a variety of seeds, occasionally insects in season.	— <u>B</u> <u>B</u> — <u>b</u> <u>B</u> <u>b</u> — — <u>h</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> — — <u>B</u> <u>B</u> <u>B</u> <u>h</u> — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Plains Pocket Mouse <i>Perognathus flavescens</i> 893.0 PEFE Resident Peripheral Rare Life Form 15	NG NSSU	Sagebrush-grasslands, yucca-grasslands, grama grasslands, sand dunes, occasionally agricultural fields. Prefers sandy soils with sparse vegetation.	Nests in a maternal burrow. Feeds on a variety of seeds. Mainly distributed east of Wyoming.	— — — — <u>b</u> — — — — — — — — — — — — — <u>b</u> — <u>B</u> — — — — — <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Beaver <i>Castor canadensis</i> 875.0 CACA Resident Common Life Form 3 16	FB	Aquatic habitats adjacent to aspen, willow, or cottonwood stands.	Nests in a lodge in water or a stream bank. Feeds on a wide variety of leaves and twigs, inner bark of woody and aquatic plants; prefers aspen and willow. Harvested in all latilongs for its fur.	B B
Western Harvest Mouse <i>Reithrodontomys megalotis</i> 877.0 REME Resident Common Life Form 5 15	NG	Basin-prairie and mountain-foothills shrublands, riparian shrub, sagebrush-grasslands, eastern great plains grasslands often associated with yucca.	Nests in dense vegetation; nest is woven of plant fibers. Feeds on a wide variety of seeds, insects, fruit, leaves, basal parts of plants.	h b B b B b B h B B h h B B B h B B
Plains Harvest Mouse <i>Reithrodontomys montanus</i> 876.0 REMO Resident Peripheral Rare Life Form 5	NG NSS3	Eastern great plains grasslands, most frequently grama grasslands. Prefers well vegetated areas, occasionally found in weedy areas. Uses rock outcrops for shelter.	Nests in a rock crevice, burrow, dense thicket; nest is made of dried leaves, lined with finely chewed, dry plant fibers. Feeds primarily on seeds; also green shoots, new grass leaves, fruit, berries.	h b B B B B
Canyon Deermouse <i>Peromyscus crinitus</i> 878.0 PECR Resident Peripheral Rare Life Form 4	NG NSS3	Juniper, rock outcrops.	Nests in a loose accumulation of plant fiber in a rock crevice. Diet is unknown.	B

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
White-footed Deermouse <i>Peromyscus leucopus</i> 881.0 PELE Resident Rare Life Form 5 7 11 14 15 17	NG	Deciduous woodlands, cottonwood-riparian, occasionally grasslands, open areas near woodlands.	Nests in a hollow tree, in shrubs, a hole, burrow, or human-built structure; nest is made of plant fibers, lined with very finely chewed fibers. Feeds on green sprouts, insects, fruit, nuts, buds.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
North American Deermouse <i>Peromyscus maniculatus</i> 880.0 PEMA Resident Abundant Life Form 7 11 15 17	NG	Virtually all habitats. Usually the most abundant small mouse in any given area.	Nests under a log or in a rock crevice; nest is woven from plant fibers, lined with very finely chewed fibers. Feeds on seeds, fruit, nuts, fungi, insects, buds, green shoots, carrion of other small mammals.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Piñon Deermouse <i>Peromyscus truei</i> 879.0 PETR Resident Rare Life Form 4 5 11 15	NG NSS3	Pine-juniper, drier shrublands, rock outcrops.	Nests in a hollow juniper tree or rock crevice; nest is made of shredded juniper bark and grass. Feeds on seeds, mushrooms, leaves, insects; also juniper berries in the winter. Mainly distributed southwest of Wyoming.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>
Northern Grasshopper Mouse <i>Onychomys leucogaster</i> 882.0 ONLE Resident Common Life Form 15	NG	Basin-prairie and mountain-foothills shrublands, grasslands, fallow lands, rock outcrops, disturbed areas. Requires silty or sandy soils.	Nests in a shallow burrow. Feeds primarily on beetles, grasshoppers, spiders, larval moths, small rodents in the spring and summer; also forbs, grasses, seeds in the winter.	<u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Bushy-tailed Woodrat <i>Neotoma cinerea</i> 883.0 NECI Resident Common Life Form 4 5 7 11 17	NG	Most habitat types with rock outcrops. Frequently occupies abandoned buildings, caves, mines.	Nests in a vertical chimney or crack in a rock outcrop, or on a shelf or in an attic of an abandoned building; nest is made of fine, dry plant fibers centered in other large objects such as pine needles, sticks, etc. Feeds on foliage of most shrubs, except sagebrush, seeds, bark, mushrooms, fruit, insects.	<u>h</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Southern Red-backed Vole <i>Myodes gapperi</i> 884.0 MYGA Resident Common Life Form 5 15	NG	Coniferous forests, aspen, mixed deciduous forests containing downed timber and windfalls, willow/wet meadow associations.	Nests on the ground; nest is made of finely chewed, interwoven dry grass and stems. Feeds on leaves, nuts, seeds, berries, moss, lichens, ferns, fungi, arthropods.	<u>B</u> <u>B</u> <u>h</u> <u>B</u> <u>B</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>B</u> <u>—</u> <u>B</u> <u>h</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u>
Western Heather Vole <i>Phenacomys intermedius</i> 885.0 PHIN Resident Common Life Form 5 15	NG	Lodgepole pine, spruce-fir, aspen, sagebrush-grasslands, brushy riparian shrub, alpine meadows, occasionally boulder fields.	Nests in a burrow; nest is made of lichens and grass. Feeds on seeds, beargrass, huckleberries, fungi, willow and birch bark, leaves or buds of various shrubs.	<u>B</u> <u>h</u> <u>—</u> <u>B</u> <u>B</u> <u>—</u> <u>—</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>B</u> <u>h</u> <u>h</u> <u>—</u> <u>—</u> <u>—</u> <u>—</u> <u>B</u> <u>—</u> <u>—</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u>
Long-tailed Vole <i>Microtus longicaudus</i> 888.0 MILO Resident Common Life Form 5 15	NG	Mountain-foothills grasslands, alpine and wet meadows, marsh-swamp wetlands in or near coniferous forests, aspen, sagebrush-grasslands, mountain-foothills shrublands, riparian shrub.	Nests in a burrow; at times in a rock crevice or hollow log. Feeds on grasses, sedges, bulbs, a wide variety of herbaceous vegetation.	<u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>—</u> <u>h</u> <u>—</u> <u>—</u> <u>h</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>h</u> <u>h</u> <u>B</u> <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Montane Vole <i>Microtus montanus</i> 887.0 MIMO Resident Common Life Form 5 15	NG	Mixed grass-like meadows, sedge, marsh-swamp wetlands, irrigated native meadows in or near coniferous forests, aspen, sagebrush-grasslands, mountain-foothills and riparian grasslands, riparian shrub. Generally occurs at higher elevations than the similar meadow vole.	Nests in a burrow; nest is made of dried grass. Feeds on grasses, sedges, rushes, fungi. Can impact native hay pastures at high densities.	B B B B B _ _ B B B b _ _ _ B B B B b B _ B B B B B B h
Prairie Vole <i>Microtus ochrogaster</i> 890.0 MIOC Resident Common Life Form 5 15	NG	Basin-prairie shrublands, grasslands, eastern great plains grasslands, small grain agricultural areas.	Nests in a burrow or depression. Feeds on certain parts of green vegetation; also seeds, fruit, bark, tubers.	_ b B B B B b h _ b b b B B _ _ B _ B B B _ _ _ _ B B B
Meadow Vole <i>Microtus pennsylvanicus</i> 886.0 MIPE Resident Common Life Form 5 15	NG	Moist to wet meadows and grasslands in or near coniferous forests, aspen, basin-prairie shrublands, sagebrush-grasslands, mixed shrublands, riparian shrub, grasslands. Generally occurs at lower elevations than the similar montane vole.	Nests at the base of a stump or shrub; nest is made of woven dry grass and other plant fibers. Feeds on most available herbaceous vegetation including grasses, sedges.	B B h B B B B B B B _ _ h B B _ B _ _ _ b B B B _ _ B b
Water Vole <i>Microtus richardsoni</i> 889.0 MIRI Resident Rare Life Form 16	NG NSS3	Subalpine and alpine meadow watercourses with overhanging banks, occasionally willow, shrubby cinquefoil, dry alpine meadows, and mountain-foothills grasslands adjacent to streams.	Prefers to nest in a burrow dug into an overhanging stream bank, usually with both above-ground and underwater entrances. Feeds on a variety of grasses, sedges, seeds, inner bark of small woody plants, nutritious herbs.	B B _ B _ _ _ B B _ _ b _ _ h B b _ _ _ _ h _ _ _ _ _

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Sagebrush Vole <i>Lemmiscus curtatus</i> 891.0 LECU Resident Common Life Form 15	NG NSS4	Basin-prairie shrublands, sagebrush-grasslands, eastern great plains and great basin-foothills grasslands, agricultural areas.	Nests in a burrow. Feeds on grasses, seeds tender shoots, flower buds, leaf buds, leaves.	— — — <u>b</u> — — <u>B</u> — <u>b</u> <u>b</u> — — — <u>B</u> — <u>B</u> <u>B</u> — — <u>B</u> <u>B</u> <u>B</u> — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Common Muskrat <i>Ondatra zibethicus</i> 892.0 ONZI Resident Common Life Form 3 16	FB	Marsh-swamp wetlands, aquatic habitats.	Nests in the center of a conical heap of mud and vegetation, or deep in a burrow in a stream bank; nest is lined with fresh vegetation. Feeds on roots and basal parts of aquatic vegetation, emergent vegetation, crayfish, fish, mollusks. Harvested in all latilongs for its fur.	<u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Norway Rat <i>Rattus norvegicus</i> 894.2 RANO Resident Uncommon Life Form 15 17	NG	Agricultural areas, urban areas.	Nests in a burrow or human-built structure; nest is made of soft material including grass, leaves, paper. Feeds on carrion, grain, alfalfa pellets, greasy paper, small mammals, green plants, any other available food. Introduced species; is considered a pest.	— — — — <u>B</u> <u>B</u> — — — — — — — — — — — — <u>B</u> — — — — — — — <u>B</u> <u>B</u>
House Mouse <i>Mus musculus</i> 894.1 MUMU Resident Common Life Form 5 15 17	NG	Agricultural areas, rural developments, urban areas.	Nests on the ground, in a burrow, or in a human-built structure; nest is made of any soft material including paper, cloth, dried grass, upholstery of abandoned furniture. Feeds on anything edible. Introduced species.	— — <u>B</u> — <u>h</u> <u>B</u> <u>b</u> — — <u>B</u> <u>B</u> — <u>B</u> <u>B</u> — — <u>B</u> — <u>B</u> <u>b</u> <u>b</u> <u>h</u> <u>B</u> — — — <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Meadow Jumping Mouse <i>Zapus hudsonius</i> 895.0 ZAHU Resident Rare Life Form 3 15	NG NSS4	Marshy areas and moist meadow grasslands near streams in eastern grasslands, coniferous and deciduous forests, mixed shrublands, riparian shrub, at lower elevations.	Nests in a burrow. Feeds on selected buds, leaves, insects, grasses, fungi, berries, nuts. May be seriously impacted by heavy livestock grazing along riparian areas.	— — — — — <u>b</u> <u>B</u> —
Preble's Meadow Jumping Mouse <i>Zapus hudsonius preblei</i> 895.1 ZAHP Resident Rare Life Form 3 15	NG NSS3 Thr.	Marshy areas and moist meadow grasslands near streams in eastern grasslands, coniferous and deciduous forests, mixed shrublands, riparian shrub, below 8,000 feet.	Nests in a burrow. Feeds on selected buds, leaves, insects, grasses, fungi, berries, nuts. May be seriously impacted by heavy livestock grazing along riparian areas.	— — — — — — — — — — — — — — — — — — — <u>O</u> <u>B</u> — — — — — — <u>O</u> <u>B</u> <u>b</u> —
Western Jumping Mouse <i>Zapus princeps</i> 896.0 ZAPR Resident Uncommon Life Form 3 15	NG	Marshy areas, moist meadow grasslands, pond margins and streambanks in coniferous and deciduous forests, riparian shrub, willow, mountain-foothills grasslands, mixed grass-like meadows, at higher elevations and in the foothills.	Nests in a burrow; nest is made of finely shredded, clean, dry plant fibers. Feeds on a variety of seeds, especially in late summer, invertebrates.	<u>B</u> <u>B</u> <u>h</u> <u>B</u> <u>B</u> — — <u>B</u> <u>B</u> — <u>B</u> <u>B</u> — — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — <u>B</u> <u>B</u> — <u>B</u> <u>B</u> <u>B</u> <u>B</u>
North American Porcupine <i>Erethizon dorsatum</i> 900.0 ERDO Resident Common Life Form 6	PD	Most habitat types including coniferous and deciduous forests, vegetated riparian areas, sagebrush-grasslands, grasslands.	No nest is made. Feeds on the inner bark of trees, evergreen needles; also buds, leaves, small twigs, herbs in the summer.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — <u>B</u> <u>B</u> <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Coyote <i>Canis latrans</i> 901.0 CALA Resident Common Life Form 15	PD	Most habitat types including plains, deserts, and mountains with grass and shrubs, dense forest.	Whelps in a den. Feeds on almost anything edible including voles, ground squirrels, carrion; also plants, frogs, lizards, insects.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Gray Wolf <i>Canis lupus</i> 902.0 CALU Resident Uncommon Life Form 15	TG	Coniferous forests, mountain-foothills shrublands, mountain-foothills grasslands.	Whelps in a den. Feeds mainly on elk, deer, moose; also smaller mammals. Reintroduced into Yellowstone National Park in 1995 as an experimental, non-essential population. Removed from the federal Endangered Species List in 2008; delisting was revoked several months later. Delisted in 2011; relisted in 2014 due to a court decision.	<u>B</u> <u>B</u> <u>O</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>H</u> <u>H</u> <u>H</u> <u>O</u> <u>B</u> <u>B</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u> <u>H</u>
Swift Fox <i>Vulpes velox</i> 904.0 VUVE Resident Common Life Form 15	NG NSS4	Short-grass or mid-grass prairies, including eastern great plains grasslands and sagebrush-grasslands, occasionally agricultural areas, irrigated native meadows, roadside/railroad banks.	Uses underground dens year-round. Feeds on horned larks, jackrabbits, deer mice in the winter; ground squirrels in the spring; beetles, small mammals, grasshoppers in the summer and fall. Incidental harvest in eastern counties.	<u>—</u> <u>—</u> <u>—</u> <u>b</u> <u>O</u> <u>b</u> <u>b</u> <u>—</u> <u>—</u> <u>—</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>—</u> <u>O</u> <u>O</u> <u>O</u> <u>B</u> <u>b</u> <u>B</u> <u>O</u> <u>b</u> <u>—</u> <u>O</u> <u>b</u> <u>B</u> <u>B</u>
Red Fox <i>Vulpes vulpes</i> 903.0 VUVU Resident Common Life Form 15	PD	Most habitat types. Prefers a mixture of streamside communities, rolling agricultural areas, brush, pastures, open areas.	Whelps in a den. Feeds primarily on mice, insects, plant matter in the summer; rabbits in the winter. Harvested throughout Wyoming for its fur.	<u>—</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Common Gray Fox <i>Urocyon cinereoargenteus</i> 905.0 URCI Resident Rare Life Form 5 15	NG	Deciduous forests, cottonwood-riparian, basin-prairie shrublands, sagebrush-grasslands, riparian shrub, grasslands, agricultural areas, rock outcrops, roadside/railroad banks.	Whelps in a den. Feeds on rabbits, squirrels, mice, rats, weasels, pocket gophers, songbirds, crows, ducks, bird eggs, turtles, muskrats, domestic cats and dogs, insects, wheat, corn, grasses, sedges, nuts, elderberries, apples, grapes, pears, grain. Probably found mostly in eastern counties.	— — <u>b</u> — — — <u>B</u> <u>O</u> — <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> — — — <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> — — <u>b</u> <u>O</u> —
Black Bear <i>Ursus americanus</i> 940.0 URAM Resident Common Life Form 15	TG	Coniferous forests, aspen, riparian shrub, mountain-foothills grasslands.	Whelps in a den. Feeds mainly on grasses, forbs, buds, berries, roots; also some insects, beetles, carrion. Harvested in latilongs of occurrence. Has increased distribution in past decade.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> — <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u>
Grizzly Bear <i>Ursus arctos</i> 941.0 URAR Resident Uncommon Life Form 15	TG Thr.	Coniferous forests, mountain-foothills shrublands, riparian shrub, mountain-foothills grasslands, alpine tundra, alpine talus.	Whelps in a den. Feeds on a wide variety of foods including ants, moths, gophers, ground squirrels, deer, elk, bison; grasses and forbs in the spring and summer; pine nuts in the fall. Removed from the federal Threatened Species List in 2007; added to Threatened Species List in 2009 due to litigation. Proposed delisting in 2016.	<u>B</u> <u>B</u> <u>B</u> <u>H</u> — — <u>H</u> <u>B</u> <u>B</u> <u>B</u> — — — — <u>B</u> <u>B</u> <u>O</u> — <u>H</u> — <u>H</u> <u>H</u> — — <u>H</u> <u>H</u> <u>H</u> <u>H</u>
Ringtail <i>Bassariscus astutus</i> 906.0 BAAS Accidental Rare Life Form	NG NSSU	Most often associated with canyons and rock outcrops within mixed woodland and shrubland as well as dense riparian shrub. Dens in rock crevices, small caves, hollow trees, and downed logs. May also be found in agricultural and suburban areas.	Nests in a den in a rock crevice or burrow. Feeds primarily on insects, crickets, beetles, ants, lizards, pocket gophers, squirrels, cottontails, birds, fruit of prickly pear and persimmons, juniper berries.	— <u>O</u> <u>O</u> — — <u>b</u> <u>O</u> <u>b</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Northern Raccoon <i>Procyon lotor</i> 907.0 PRLO Resident Common Life Form 4 14	PD	Especially abundant in cottonwood-riparian areas and along perennial streams, but is found in most habitat types below 6,500 feet where water is available, including urban areas.	Nests in a den. Feeds on a wide variety of plants and animals, especially aquatic animals and insects.	<u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Pacific Marten <i>Martes caurina</i> 908.0 MACA Resident Uncommon Life Form 14	FB	Mature spruce-fir forests; also lodgepole pine, Douglas fir, occasionally cottonwood-riparian. Considered an old-growth conifer obligate.	Nests in a den in a hollow tree on the ground, or underground. Feeds on small mammals, especially red-backed voles, birds, insects, fruit, berries. Harvested in some latilongs for its fur.	<u>B</u> <u>B</u> <u> </u> <u>b</u> <u>b</u> <u> </u> <u> </u> <u>B</u> <u>B</u> <u>b</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>b</u> <u>h</u> <u>B</u> <u> </u> <u>b</u> <u> </u> <u> </u> <u> </u> <u>b</u> <u> </u> <u>b</u> <u>B</u> <u>O</u> <u> </u>
Fisher <i>Pekania pennanti</i> 909.0 PEPE Accidental Rare Life Form	NG	Coniferous forests, aspen, cottonwood-riparian. Prefers coniferous forests with a continuous closed canopy.	Nests in a hole, either naturally occurring or one made by another animal. Feeds on small to medium sized birds, snowshoe hares, red and flying squirrels, mice, voles, shrews, porcupines; also carrion, especially deer carcasses. Mainly distributed north of Wyoming. Classified as a protected species in Wyoming.	<u>b</u> <u>b</u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u> <u>O</u> <u>b</u> <u> </u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u>O</u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u> </u> <u>O</u> <u> </u> <u> </u>
Short-tailed Weasel (Ermine) <i>Mustela erminea</i> 910.0 MUER Resident Uncommon Life Form 15	FB	Found in most habitat types, especially coniferous forests, riparian shrub, alpine meadows, marsh edges.	Nests in a burrow. Feeds mainly on small mammals, especially voles; also songbirds, beetles, grasshoppers, frogs. Harvested in some latilongs for its fur.	<u> </u> <u>b</u> <u> </u> <u>b</u> <u>b</u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u> </u> <u>b</u> <u> </u> <u> </u> <u>b</u> <u>b</u> <u>B</u> <u> </u> <u>B</u> <u>b</u> <u>b</u> <u>b</u> <u>B</u> <u>b</u> <u>b</u> <u>B</u> <u>h</u> <u>b</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Long-tailed Weasel <i>Mustela frenata</i> 911.0 MUF Resident Common Life Form 15	FB	Found in most habitat types including open fields, willow, desert shrub, grasslands, riparian shrub. Typically found in rock outcrops near water. Often occupies open habitats in or near coniferous zones.	Nests in a burrow. Feeds mainly on shrews, mice, rats, squirrels, rabbits; also birds, bird eggs, insects, carrion. Harvested in some latilongs for its fur.	h b b B b b b b b b b b b B b b B B b b b b b b b b B
Black-footed Ferret <i>Mustela nigripes</i> 913.0 MUNG Resident Rare Life Form 15	NG NSS1 End.	Found in association with prairie dog colonies in basin-prairie shrublands, sagebrush-grasslands, eastern great plains and great basin-foothills grasslands.	Occupies prairie dog burrows. Feeds primarily on prairie dogs; also deer mice, pocket gophers, pocket mice, birds, ground squirrels. Classified as a federally endangered species; is a protected species in Wyoming. Is very rare or extinct throughout its range in N.A.	O O O h O h O O O O h h O O O O B h h h O h h B B h
Least Weasel <i>Mustela nivalis</i> 919.0 MUNV Resident Rare Life Form 15	NG NSSU	Meadows, riparian willow and cottonwood in basin sagebrush-grasslands.	Nests in a burrow. Feeds mainly on voles, mice, shrews; also birds, insects.	B b b b b b b
American Mink <i>Vison vison</i> 912.0 VIVI Resident Common Life Form 16	FB	Most habitat types near open water, especially riparian zones adjacent to watercourses and ponds.	Nests in a burrow. Feeds on mice, rats, rabbits, muskrats, frogs, fish, crayfish, birds, invertebrates, snakes. Harvested throughout Wyoming for its fur.	B B b b b B B B B O b b B b b B b b b b b b b B B

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Wolverine <i>Gulo gulo</i> 914.0 GUGU Resident Rare Life Form 5	NG NSS3	Coniferous forests, especially dense, continuous stands in remote areas.	Nests on the ground. Feeds on mule deer, elk, moose, rabbits, hares, porcupines, beaver, squirrels, chipmunks, marmots, mice, birds, gophers, berries in the summer, carrion in late winter/early spring. Mainly distributed north of Wyoming. Classified as a protected species in Wyoming.	b O _ _ O _ _ b b _ _ _ _ b b O _ _ _ O _ _ _ b _ O
American Badger <i>Taxidea taxus</i> 915.0 TATA Resident Common Life Form 15	FB	Most common in basin grasslands and basin sagebrush-grasslands. Occurs in most habitat types with relatively deep soils.	Dens in a burrow. Feeds on ground squirrels, prairie dogs, rabbits, hares, chipmunks, marmots, squirrels, gophers, mice, rats, coyotes, skunks. Harvested throughout Wyoming for its fur.	_ b B b b b b B B b b B B B b B B B B B B B B B B
Western Spotted Skunk <i>Spilogale gracilis</i> 916.1 SPGR Resident Uncommon Life Form 15	PD NSSU	Basin-prairie shrublands, greasewood, grasslands.	Dens in the burrow of another animal, or a hole, rock pile, or fallen log. Feeds on beetles, crickets, grubs, grasshoppers, worms, carrion, rodents, young rabbits, bird eggs, frogs, crayfish, lizards, some fruit.	_ B b _ _ _ b _ O b b _ _ _ _ h B B b _ b b _ _ b _ _
Eastern Spotted Skunk <i>Spilogale putorius</i> 916.0 SPPU Resident Uncommon Life Form 15	PD NSS3	Ponderosa pine, pine-juniper, cottonwood-riparian, irrigated native meadows, rural developments, rock outcrops, roadside/railroad banks, near streams and rivers.	Dens in a burrow made by another animal, under a building or brush pile, or in a rock crevice. Feeds on fruit, berries, corn, carrion, nuts, voles, mice, birds, bird eggs.	_ _ _ _ _ _ _ _ _ _ b _ h _ _ _ _ B b b _ _ _ O O O b

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Striped Skunk <i>Mephitis mephitis</i> 917.0 MEME Resident Common Life Form 15	PD	Most habitat types including mixed woodlands, brushy areas, open fields with broken wooded ravines, rock outcrops.	Dens in a burrow. Feeds primarily on grasshoppers, beetles, crickets, butterfly larvae, deer mice, voles, bird eggs, berries, fruit.	h _ B B B b B B B b B B B B B B B B B B B B b b b b B B
Northern River Otter <i>Lontra canadensis</i> 918.0 LOCA Resident Uncommon Life Form 16	NG NSS3	Lakes, streams, and aquatic habitats in aspen, cottonwood-riparian, riparian shrub, willow, most meadow grasslands, and marsh-swamp wetlands.	Dens in a natural shelter, often in a driftwood pile or den made by another animal, especially beaver. Feeds primarily on fish, some amphibians, insects, birds, mammals. Classified as a protected species in Wyoming.	B b O O O _ _ B B O _ B _ _ B B b b b _ _ O B _ _ O B _
Mountain Lion (Puma) <i>Puma concolor</i> 922.0 PUCO Resident Common Life Form 4	TG	Most habitat types where deer and other ungulate species are present. Prefers remote areas with dense cover and rocky, rugged terrain.	Dens in an inaccessible, remote area such as a cave, rock crevice, dense shrub, or under cover. Feeds primarily on deer, elk, small and medium-sized mammals. Is an obligate carnivore. Harvested as a trophy game animal in most Wyoming latilongs.	B B
Canada Lynx <i>Lynx canadensis</i> 920.0 LYCA Resident Rare Life Form 5	NG NSS1 Thr.	Dense coniferous forests, especially Englemann spruce-subalpine fir, at high elevations.	Dens on the ground. Feeds primarily on snowshoe hares; also mice, grouse, squirrels. Classified as a protected species in Wyoming.	h h _ h O _ _ B O _ _ _ _ _ O _ _ _ _ _ _ _ O _ _ _ O _

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Bobcat <i>Lynx rufus</i> 921.0 LYRU Resident Common Life Form 4	FB	Most habitat types with broken, brushy country or mountains. Often found in rocky areas with cliffs or along rocky rims.	Dens on a cliff or rim, or in a cave or talus field. Feeds on birds, rabbits, hares, marmots, porcupines, beavers, ungulates, squirrels, voles, mice, pocket gophers; also fish, amphibians, reptiles, insects. Harvested in some latilongs for its fur.	<u> </u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u>
Wapiti (Elk) <i>Cervus canadensis</i> 930.0 CECA Resident Common Life Form 5	BG	Coniferous forests, aspen, basin-prairie and mountain-foothills shrublands, great basin-foothills and mountain-foothills grasslands, agricultural areas.	No nest is made. Feeds on grasses and forbs in the winter, grasses in the spring, forbs in the summer, shrubs year-round. Harvested in latilongs of occurrence as a big game animal.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
Mule Deer (Black-tailed Deer) <i>Odocoileus hemionus</i> 932.0 ODHE Resident Common Life Form 5	BG	Most habitat types including desert, riparian areas, broken grasslands, shrublands, foothills, coniferous and deciduous forests.	No nest is made. Feeds on a wide variety of plants including forbs, grasses, sedges, shrubs, trees. Diet depends on habitat type, age, sex, season, and the condition of each animal. Harvested in all latilongs as a big game animal.	<u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u>
White-tailed Deer <i>Odocoileus virginianus</i> 933.0 ODVI Resident Common Life Form 5	BG	Deciduous forests, sagebrush-grasslands, mountain-foothills shrublands, agricultural areas, riparian shrub.	No nest is made. Feeds on bearberry, cottonwood, aspen, snowberry, willow, serviceberry, ponderosa pine, Douglas fir, crops. Is both a browser and a grazer. Harvested in some latilongs as a big game animal.	<u>b</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>B</u> <u>B</u> <u>b</u> <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u> </u> <u>b</u> <u>b</u> <u>B</u> <u>B</u>

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Moose <i>Alces americanus</i> 931.0 ALAM Resident Common Life Form 5	BG NSS4	Coniferous forests; aspen; cottonwood-riparian; mountain-foothills shrublands; willow; wild plum; water birch; bog birch; alder; shrubby cinquefoil; alpine, moist meadow, Kentucky bluegrass, and annual grasslands; mixed alpine and mixed grass-like meadows; agricultural areas.	No nest is made. Feeds on willows, antelope bitterbrush, Douglas fir, subalpine fir, white-bark pine, cottonwood, sedges, rushes, blue spruce. Harvested in some latilongs as a big game animal.	B B B B B _ _ B B B _ _ _ _ B B B B O _ _ B B _ B B B _
Pronghorn <i>Antilocapra americana</i> 935.0 ANAM Resident Common Life Form 5	BG	Basin-prairie and mountain-foothills shrublands, eastern great plains and great basin-foothills grasslands, sagebrush-grasslands.	No nest is made. Feeds on shrubby plants, especially sagebrush; also forbs in the spring and summer, some grasses. Harvested in all latilongs as a big game animal.	B B
Bison <i>Bos bison</i> 925.0 BOBI Resident Rare Life Form 5	BG	Sagebrush-grasslands, mountain-foothills grasslands, eastern great plains grasslands.	No nest is made. Feeds on sedges, grasses, some forbs and browse in the summer. Formerly a widespread species in North America. Historically, wild populations occurred throughout Wyoming. Domestic populations which currently occur statewide are not included.	B O h h h h h B h
Mountain Goat <i>Oreamnos americanus</i> 926.0 ORAM Resident Rare Life Form 4	BG	Coniferous forests, mountain-foothills shrublands, mountain-foothills and alpine grasslands, mixed alpine meadows, cliffs.	No nest is made. Feeds on sedges, grasses, rushes, forbs, evergreen shrubs, conifers. Introduced species. L2: harvested as a big game animal.	b B _ _ _ _ _ b _ _ _ _ _ _ _ _ O _ _ _ _ _ _ _ _ _ _ _

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION
Bighorn Sheep (Mountain Sheep) <i>Ovis canadensis</i> 927.0 OVCA Resident Common Life Form 4	BG NSS4	Coniferous forests; winterfat; mountain-foothills shrublands; great basin-foothills, mountain-foothills, and alpine grasslands; mixed alpine meadows; cliffs; rock outcrops.	No nest is made. Feeds on a variety of grasses, forbs, browse. Harvested in most latilongs of occurrence as a big game animal.	B B B B _ _ h B B B B h _ _ B B B B B B _ _ _ _ _ B B B

Amphibians & Reptiles

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Tiger Salamander <i>Ambystoma mavortium</i> 950.0 AMMA Resident Common Life Form 1 3 14	A	Most habitat types with standing or slow moving water nearby for breeding. Overwinters in soil, rodent burrows, cellars and other suitable moist habitat.	Deposits eggs in shallow standing or slow moving water, where larvae remain for 2-24 months. Metamorphosed salamanders leave the water and return only to breed and lay eggs. Neoteny may occur in Wyoming populations. Adults feed on earthworms, insects, small vertebrates. Larvae feed on aquatic invertebrates, other salamander larvae. Includes Blotched, Western, and Arizona subspecies.	B O O O# B# B# h* B O# O O O O h* O O B B O O O O B B O B B O
Plains Spadefoot <i>Spea bombifrons</i> 951.0 SPBO Resident Common Life Form 1 5 14 15	A NSS4	Grasslands and sagebrush communities below 6,000 feet, east of the Continental Divide. Excavates a deep burrow in the winter to avoid desiccation and spend periods of dormancy below the frost line. Prefers sandy and other loose soils for burrowing.	Deposits eggs in flooded areas and temporary pools formed by heavy rains. Returns to semiarid shrublands after egg laying. Eggs hatch in 2-3 days. Tadpoles complete metamorphosis in 36-40 days. Adults feed on moths, beetles, caterpillars, other small arthropods. Larvae feed on aquatic vegetation, small invertebrates, other spadefoot larvae.	— h* B ? h ? ? — — O O* B B O — — — — B O* B — — — — O B B
Great Basin Spadefoot <i>Spea intermontana</i> 951.1 SPIN Resident Common Life Form 1 5 14 15	A NSS4	Sagebrush communities mainly below 6,000 feet, west of the Continental Divide. Overwintering habits are probably similar to other spadefoot toads. Prefers sandy and other loose soils for burrowing.	Deposits eggs in springs or flooded areas formed by heavy rains; may also deposit eggs in permanent water sources. Breeding, egg laying, and feeding habits are probably similar to other spadefoot toads.	— — — — — — — — — — — — — B B B — — — h* B B O — — —
Western Toad (Boreal Toad) <i>Anaxyrus boreas</i> 951.2 ANBO Resident Common Life Form 1 3 14 15	A NSS1	Wet areas in foothills, montane, and subalpine zones from 6,000 to 11,000 feet.	Deposits eggs in ponds and small lakes. Tadpoles hatched below 10,000 feet metamorphose during their first summer; those above 10,000 feet may fail to transform or survive overwinter. Adults feed primarily on ants; also beetles, moths, other insects. Populations in the Medicine Bow and Sierra Madre Mountains are declining. Formerly Boreal Toad.	B O# — — — — B B — — — — B O O* — — h — O* — O O# B h* —

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Great Plains Toad <i>Anaxyrus cognatus</i> 951.3 ANCO Resident Uncommon Life Form 1 5 14 15	A NSSU	Plains grasslands, sandhills, agricultural areas, below 6,000 feet.	Deposits eggs in ponds and flooded areas formed by heavy rains. Eggs hatch in 2-3 days. Tadpoles begin metamorphosis at 45 days. Adults feed on moths, caterpillars, cutworms, flies, beetles, other small insects.	— —
Wyoming Toad <i>Anaxyrus baxteri</i> 951.5 ANBA Resident Rare Life Form 1 3 14 15	A NSS1 End.	Floodplains, ponds, and small seepage lakes in the shortgrass prairie of the Laramie basin.	Deposits eggs in ponds and small lakes. Egg laying and tadpole metamorphosis are similar to other toads in the genus Bufo. Adults feed on ants, beetles, other small insects. Primarily nocturnal and is an adept burrower in soft soil.	— —
Rocky Mountain Toad (Woodhouse's Toad) <i>Anaxyrus woodhousii woodhousii</i> 951.4 ANWW Resident Common Life Form 1 3 14 15	A	Floodplains containing permanent or ephemeral pools in the plains and foothills east of the Continental Divide below 6,000 feet.	Deposits eggs in floodplain containing permanent or ephemeral pools. Data are lacking, but egg laying and tadpole metamorphosis are probably similar to other toads in the genus Anaxyrus. Adults feed on ants, beetles, other small insects.	h* O* O O h* h h* —
American Bullfrog <i>Lithobates catesbeianus</i> 952.1 LICA Resident Uncommon Life Form 1 3	A	Permanent water below 6,000 feet on the eastern plains. Specimens collected at warm springs in western and northwestern Wyoming were probably introduced by humans.	Deposits eggs in permanent bodies of water. Tadpoles may attain a length of 3 inches or more before metamorphosis. Some tadpoles may overwinter and transform the second summer. Adults feed on birds, frogs, snakes, insects.	O — — — — — — — — — — O —

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Northern Leopard Frog <i>Lithobates pipiens</i> 952.2 LIPI Resident Common Life Form 1 3	A NSS4	Swampy cattail marshes, beaver ponds, streams, rivers, and lakes in the plains, foothills, and montane zones up to 9,000 feet.	Deposits eggs in shallow, non-flowing areas of permanent bodies of water, streams, and rivers or seasonally flooded areas near permanent pools, streams, and rivers. Eggs hatch in 4-15 days. Tadpoles metamorphose in 14-60 days. Adults feed on insects, invertebrates, tadpoles, snakes, fish.	_ h* h* O O# h* O O# O# O O# O O O O* O B B O O B O h* _ O O O# O
Columbia Spotted Frog <i>Rana luteiventris</i> 952.3 RALU Resident Common Life Form 1 3	A NSS3	Ponds, sloughs, and small streams in the foothills and montane zones. May avoid warm stagnant ponds with cattails.	Deposits eggs in small streams and beaver ponds in late May or June. Tadpoles metamorphose in late August or early September. Adults feed on insects, invertebrates, tadpoles, frogs.	B O# B _ _ _ B O# _ _ _ _ h# O _ _ _ _ _ O _ _ _ _ _
Wood Frog <i>Lithobates sylvaticus</i> 952.4 LISY Resident Rare Life Form 1 3 5	A NSS2	Beaver ponds, small lakes, slow moving streams, wet meadows, willow thickets, in the montane zone usually at or near 9,000 feet.	Deposits eggs on the north side of natural ponds or inactive beaver ponds. Eggs hatch in 4-20 days. Tadpoles metamorphose in 70-85 days. Tadpoles may overwinter and transform the following spring. Move from breeding locations to other moist habitats during summer months. Adults feed on insects, worms, spiders.	_ _ _ B _ B _ _
Boreal Chorus Frog <i>Pseudacris maculata</i> 952.0 PSMA Resident Common Life Form 1 3	A	Moist margins of marshes, ponds, small lakes, creeks, up to 12,000 feet.	Deposits eggs in rain pools, marshes, bog ponds, lakes, reservoirs, flooded areas, and other water sources lacking current. Tadpoles metamorphose in approximately 60 days. Adults feed on small invertebrates, insects.	B O O _ O h* O B O* ? O O ? O B B O O B O O B O h* O B B ?

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Eastern Spiny Softshell <i>Apalone spinifera spinifera</i> 953.0 APSS Resident Common Life Form 1 3	NG NSS4	Permanent lakes, ponds, and large streams in association with sandbars and other sandy aquatic substrates, below 6,000 feet.	Deposits eggs on sand or fine gravel near water. Forages in water on fish, frogs, tadpoles, crayfish, aquatic insects.	-- -- 0 0 0 0 0 -- -- B -- -- B -- -- -- -- 0 0 -- -- -- -- 0 0
Plains Box Turtle <i>Terrapene ornata ornata</i> 953.2 TEOO Resident Uncommon Life Form 5 14 15	NG NSSU	A terrestrial species; prefers grasslands, sandhills.	Deposits eggs in soft, well drained soil. Feeds on insects, invertebrates, vegetation. Common in the pet trade; specimens found outside of Goshen County in L10, 17, 25, 27 are released individuals collected from Nebraska or Kansas or released pets.	-- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- 0 -- -- -- -- -- --
Western Painted Turtle <i>Chrysemys picta bellii</i> 953.3 CHPB Resident Common Life Form 1 3	NG NSS4	Permanent ponds, reservoirs, marshes, slow moving streams, below 6,000 feet.	Deposits eggs in soft soil in a sunny area. Hatchlings probably overwinter in the nest and emerge in the spring. Forages in water on fish, live and dead plants, vertebrates, invertebrates.	-- -- -- h* h* 0 ? -- -- -- 0 ? ? -- -- -- 0 0* 0 -- -- -- -- h* h*
Snapping Turtle <i>Chelydra serpentina</i> 953.1 CHSE Resident Common Life Form 1 3	NG	Permanent lakes, ponds, large streams, below 6,000 feet.	Deposits eggs in a burrow near water or in a muskrat house. Feeds on vegetation, fish, snakes, amphibians, small mammals, birds (especially ducklings), invertebrates, carrion.	-- -- -- 0 0 0 0 -- -- -- -- 0 ? -- -- -- 0 0 -- -- -- -- 0

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Plateau Fence Lizard <i>Sceloporus tristichus</i> 954.4 SCTR Resident Uncommon Life Form 4 5	NG NSS4	Rock outcrops and canyon walls in sagebrush communities.	Lizards in Utah lay three clutches of eggs averaging 6.3 eggs per clutch. It is unknown whether this also occurs in Wyoming. Feeds on grasshoppers, crickets, leaf hoppers, flying ants, moths, other insects.	-- -- -- -- -- -- -- -- -- -- -- -- -- -- -- B O -- -- O O -- -- O# --
Prairie Lizard <i>Sceloporus consobrinus</i> 954.6 SCCO Resident Uncommon Life Form 5	NG NSSU	Sandy areas with low shrubs in grassland communities, or sandstone and limestone outcrops in scarp woodland communities.	Breeding and egg laying habits are probably similar to other members of the genus <i>Sceloporus</i> . May produce two clutches of eggs per year. Feeds on grasshoppers, small beetles, other insects.	-- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- O -- -- -- -- -- -- --
Northern Tree Lizard <i>Urosaurus ornatus wrighti</i> 954.8 UROW Resident Uncommon Life Form 4 5	NG NSS1	Cliffs and rocky canyon slopes in sagebrush desert communities.	Little is known about the breeding habits of this species. It is thought to produce two clutches of 3-5 eggs per year. Feeds on thrips, plant bugs, aphids, beetles, caddisflies, moths, butterflies, flies, wasps, ants, spiders.	-- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- -- O* -- -- -- --
Greater Short-horned Lizard <i>Phrynosoma hernandesi</i> 954.2 PHHE Resident Common Life Form 5	NG NSS4	Grasslands, sagebrush-grasslands.	Bears live young; litters range from 12-24. Feeds on ants, beetles, grasshoppers, other insects.	h* O* h* O O h* h* ? O O O h* h* B O B B O h* B O B O O B h* h*

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Great Basin Gophersnake <i>Pituophis catenifer deserticola</i> 955.6 PICD Resident Uncommon Life Form 5 14	NG NSS2	Sagebrush communities and desert habitats.	Breeding, feeding, and denning habits are probably very similar to the Bullsnake.	_ _ _ _ _ <u>O</u> _ _ _ _ _ _ _ <u>O</u> _ _ _ _ _ _ <u>O</u> <u>O</u> _ _ _ _ _
Bullsnake <i>Pituophis catenifer sayi</i> 955.5 PICS Resident Common Life Form 5 14	NG	Plains grasslands, sagebrush-grasslands, sandhills, riparian shrub, marshes, rocky canyons, mountain-foothills shrub, agricultural areas, urban areas.	Clutches contain 10-20 eggs. Feeds on ground squirrels, mice, wood rats, small cottontail rabbits, other rodents; also nestling birds, bird eggs, lizards. Overwinters in large aggregations in deep underground crevices. May use the same den as Eastern Yellow-bellied Racers, Pale Milksnakes, gartersnakes, and rattlesnakes.	h* h* <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ _ <u>B</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> _ _ _ _ <u>O</u> <u>O*</u> <u>O</u>
Plains Black-headed Snake <i>Tantilla nigriceps</i> 956.4 TANI Resident Common Life Form 5 14 15	NG NSSU	Plains, grasslands, scrub brush, and woodlands. Prefers habitats with loose soil, rock outcroppings, or other structures suitable for cover.	Has been found in Platte County. Likely active from April to September. Mating probably occurs in the spring, followed by egg laying in June or July. Females lay 1-3 eggs per clutch, on average. Primarily preys on insects, earthworms and other small invertebrates. Secretive species spends most of its time underground or under cover. Forages at night often following rains. Overwinters underground in the soil.	_ _ _ _ _ _ _ _ _ _ _ _ _ _ _ <u>O</u> _ _ _ _ _
Wandering Gartersnake <i>Thamnophis elegans vagrans</i> 955.8 THEV Resident Common Life Form 3 14	NG	Found in all habitat zones except alpine, usually near water.	Bears live young; litters range from 7-14. Feeds on fish, frogs, small mammals, earthworms, slugs, grasshoppers, other small invertebrates. May overwinter in crevices or small mammal burrows, often with other snake species.	<u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ <u>h*</u> <u>O*</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> <u>O</u> _ <u>O</u> <u>O</u> <u>B</u> <u>B</u> <u>O</u> <u>O</u> <u>h*</u> <u>O</u> <u>O</u> <u>h*</u> <u>O</u> <u>O</u> <u>O</u> <u>h*</u>

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Valley Gartersnake <i>Thamnophis sirtalis fitchi</i> 956.0 THSF Resident Common Life Form 3 5 14	NG NSSU	Plains, foothills, montane zones, usually near permanent water sources.	Breeding, feeding, and denning habits are probably similar to the Red-sided Gartersnake.	— — — — — — — — O — — — — — — — — B — — — — — — — — O — — — — — — — —
Red-sided Gartersnake <i>Thamnophis sirtalis parietalis</i> 955.9 THSP Resident Common Life Form 3 5 14	NG NSSU	Plains, foothills, montane zones, near permanent water sources.	Bears live young; litters range from 12-24. Feeds on fish, frogs, small mammals, invertebrates. Large aggregations of this species may share denning sites with other snake species.	— O O O O ? O — — — — — ? O — — — — — — h* — — — — — — O
Plains Gartersnake <i>Thamnophis radix</i> 955.7 THRA Resident Common Life Form 3 5 14	NG NSSU	Near small streams, sloughs, marshes, ponds. May be found in urban areas, dry grasslands, sandhills.	Bears live young; litters range from 9-21. Feeds on fish, frogs, small mammals, insects, earthworms. Overwinters in large aggregations, often with rattlesnakes, Bullsnares, Pale Milksnares, and Eastern Yellow-bellied Racers.	— — — — — O O O — — — — — ? ? — — — — — — O h* — — — — — — h*
Prairie Rattlesnake <i>Crotalus viridis</i> 955.0 CRVI Resident Common Life Form 5 14	NG NSS4	Plains, foothills, and scarp woodlands, especially near granite or limestone outcrops.	Bears live young; litters range from 4-21. Feeds on ground squirrels, prairie dogs, chipmunks, cottontail rabbits, other rodents, frogs, toads, lizards, snakes, birds. Overwinters in large aggregations in deep underground crevices, prairie dog burrows, or other abandoned mammal burrows.	— O O O O O O — h* O O O h* O — — B B B O O — O O O O O O

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

NAME AND STATUS	MGMT. STATUS	HABITAT	COMMENTS	DISTRIBUTION^a
Midget Faded Rattlesnake <i>Crotalus oreganus concolor</i> 955.1 CROC Resident Rare Life Form 5 14	NG NSS1	Rock outcrops in sagebrush desert communities. Range is restricted to the lower Green River Valley from the cities of Green River and Rock Springs south to the Utah-Wyoming state line.	Little is known about the breeding and food habits of this species, but they are probably similar to those of the Prairie Rattlesnake.	— <u>B</u> — — — — — —

^a * = University of Wyoming Museum
 # = Wyoming Natural Diversity Database

Index

Index of Common Names

Common Name	Scientific Name	Page No.
Abert's Squirrel	<i>Sciurus aberti</i>	143
Acorn Woodpecker	<i>Melanerpes formicivorus</i>	73
African Collared-Dove	<i>Streptopelia roseogrisea</i>	33
American Avocet	<i>Recurvirostra americana</i>	41
American Badger	<i>Taxidea taxus</i>	157
American Bittern	<i>Botaurus lentiginosus</i>	60
American Black Duck	<i>Anas rubripes</i>	21
American Bullfrog	<i>Lithobates catesbeianus</i>	164
American Coot	<i>Fulica americana</i>	40
American Crow	<i>Corvus brachyrhynchos</i>	86
American Dipper	<i>Cinclus mexicanus</i>	93
American Golden-Plover	<i>Pluvialis dominica</i>	42
American Goldfinch	<i>Spinus tristis</i>	103
American Kestrel	<i>Falco sparverius</i>	76
American Mink	<i>Vison vison</i>	156
American Perimyotis (Tri-colored Bat)	<i>Perimyotis subflavus</i>	136
American Pika	<i>Ochotona princeps</i>	138
American Pipit	<i>Anthus rubescens</i>	98
American Pygmy Shrew	<i>Sorex hoyi</i>	131
American Redstart	<i>Setophaga ruticilla</i>	109
American Robin	<i>Turdus migratorius</i>	96
American Three-toed Woodpecker	<i>Picoides dorsalis</i>	75
American Tree Sparrow	<i>Spizelloides arborea</i>	116
American White Pelican	<i>Pelecanus erythrorhynchos</i>	60
American Wigeon	<i>Anas americana</i>	21
American Woodcock	<i>Scolopax minor</i>	48
Ancient Murrelet	<i>Synthliboramphus antiquus</i>	51
Anna's Hummingbird	<i>Calypte anna</i>	38

Common Name	Scientific Name	Page No.
Arctic Tern	<i>Sterna paradisaea</i>	57
Ash-throated Flycatcher	<i>Myiarchus cinerascens</i>	80
Baird's Sandpiper	<i>Calidris bairdii</i>	46
Baird's Sparrow	<i>Ammodramus bairdii</i>	119
Bald Eagle	<i>Haliaeetus leucocephalus</i>	65
Baltimore Oriole	<i>Icterus galbula</i>	128
Band-tailed Pigeon	<i>Patagioenas fasciata</i>	33
Bank Swallow	<i>Riparia riparia</i>	88
Barn Owl	<i>Tyto alba</i>	68
Barn Swallow	<i>Hirundo rustica</i>	89
Barred Owl	<i>Strix varia</i>	71
Barrow's Goldeneye	<i>Bucephala islandica</i>	27
Bay-breasted Warbler	<i>Setophaga castanea</i>	110
Beaver	<i>Castor canadensis</i>	147
Belted Kingfisher	<i>Megaceryle alcyon</i>	72
Bewick's Wren	<i>Thryomanes bewickii</i>	93
Big Brown Bat	<i>Eptesicus fuscus</i>	136
Big Free-tailed Bat	<i>Nyctinomops macrotis</i>	138
Bighorn Sheep (Mountain Sheep)	<i>Ovis canadensis</i>	161
Bison	<i>Bos bison</i>	160
Black Bear	<i>Ursus americanus</i>	154
Black Hills Red-bellied Snake	<i>Storeria occipitomaculata pahasapae</i>	170
Black Rail	<i>Laterallus jamaicensis</i>	39
Black Rosy-Finch	<i>Leucosticte atrata</i>	99
Black Scoter	<i>Melanitta americana</i>	26
Black Swift	<i>Cypseloides niger</i>	36
Black Tern	<i>Chlidonias niger</i>	57
Black Vulture	<i>Coragyps atratus</i>	64
Black-and-white Warbler	<i>Mniotilta varia</i>	106
Black-backed Woodpecker	<i>Picoides arcticus</i>	75

Common Name	Scientific Name	Page No.
Black-bellied Plover	<i>Pluvialis squatarola</i>	41
Black-billed Cuckoo	<i>Coccyzus erythrophthalmus</i>	35
Black-billed Magpie	<i>Pica hudsonia</i>	86
Blackburnian Warbler	<i>Setophaga fusca</i>	111
Black-capped Chickadee	<i>Poecile atricapillus</i>	89
Black-chinned Hummingbird	<i>Archilochus alexandri</i>	37
Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>	63
Black-footed Ferret	<i>Mustela nigripes</i>	156
Black-headed Grosbeak	<i>Pheucticus melanocephalus</i>	124
Black-headed Gull	<i>Chroicocephalus ridibundus</i>	52
Black-legged Kittiwake	<i>Rissa tridactyla</i>	52
Black-necked Stilt	<i>Himantopus mexicanus</i>	41
Blackpoll Warbler	<i>Setophaga striata</i>	111
Black-tailed Jackrabbit	<i>Lepus californicus</i>	139
Black-tailed Prairie Dog	<i>Cynomys ludovicianus</i>	143
Black-throated Blue Warbler	<i>Setophaga caerulescens</i>	112
Black-throated Gray Warbler	<i>Setophaga nigrescens</i>	113
Black-throated Green Warbler	<i>Setophaga virens</i>	114
Black-throated Sparrow	<i>Amphispiza bilineata</i>	118
Blue Grosbeak	<i>Passerina caerulea</i>	124
Blue Jay	<i>Cyanocitta cristata</i>	85
Blue-gray Gnatcatcher	<i>Polioptila caerulea</i>	93
Blue-headed Vireo	<i>Vireo solitarius</i>	83
Blue-winged Teal	<i>Anas discors</i>	22
Blue-winged Warbler	<i>Vermivora cyanoptera</i>	106
Bobcat	<i>Lynx rufus</i>	159
Bobolink	<i>Dolichonyx oryzivorus</i>	125
Bohemian Waxwing	<i>Bombycilla garrulus</i>	103
Bonaparte's Gull	<i>Chroicocephalus philadelphia</i>	52
Boreal Chorus Frog	<i>Pseudacris maculata</i>	165

Common Name	Scientific Name	Page No.
Boreal Owl	<i>Aegolius funereus</i>	72
Brambling	<i>Fringilla montifringilla</i>	99
Brant	<i>Branta bernicla</i>	19
Brazilian Free-tailed Bat	<i>Tadarida brasiliensis</i>	137
Brewer's Blackbird	<i>Euphagus cyanocephalus</i>	127
Brewer's Sparrow	<i>Spizella breweri</i>	117
Broad-tailed Hummingbird	<i>Selasphorus platycercus</i>	38
Broad-winged Hawk	<i>Buteo platypterus</i>	67
Brown Creeper	<i>Certhia americana</i>	91
Brown Pelican	<i>Pelecanus occidentalis</i>	60
Brown Thrasher	<i>Toxostoma rufum</i>	97
Brown-capped Rosy-Finch	<i>Leucosticte australis</i>	100
Brown-headed Cowbird	<i>Molothrus ater</i>	128
Buff-breasted Sandpiper	<i>Calidris subruficollis</i>	46
Bufflehead	<i>Bucephala albeola</i>	26
Bullock's Oriole	<i>Icterus bullockii</i>	128
Bullsnake	<i>Pituophis catenifer sayi</i>	171
Burrowing Owl	<i>Athene cunicularia</i>	70
Bushtit	<i>Psaltiriparus minimus</i>	90
Bushy-tailed Woodrat	<i>Neotoma cinerea</i>	149
Cackling Goose	<i>Branta hutchinsii</i>	19
California Gull	<i>Larus californicus</i>	54
California Myotis	<i>Myotis californicus</i>	133
Calliope Hummingbird	<i>Selasphorus calliope</i>	38
Canada Goose	<i>Branta canadensis</i>	19
Canada Lynx	<i>Lynx canadensis</i>	158
Canada Warbler	<i>Cardellina canadensis</i>	114
Canvasback	<i>Aythya valisineria</i>	24
Canyon Deermouse	<i>Peromyscus crinitus</i>	147
Canyon Towhee	<i>Melospiza fusca</i>	116

Common Name	Scientific Name	Page No.
Canyon Wren	<i>Catherpes mexicanus</i>	91
Cape May Warbler	<i>Setophaga tigrina</i>	110
Carolina Wren	<i>Thryothorus ludovicianus</i>	93
Caspian Tern	<i>Hydroprogne caspia</i>	57
Cassin's Finch	<i>Haemorhous cassinii</i>	101
Cassin's Kingbird	<i>Tyrannus vociferans</i>	81
Cassin's Sparrow	<i>Peucaea cassinii</i>	116
Cassin's Vireo	<i>Vireo cassinii</i>	83
Cattle Egret	<i>Bubulcus ibis</i>	62
Cedar Waxwing	<i>Bombycilla cedrorum</i>	104
Chestnut-collared Longspur	<i>Calcarius ornatus</i>	104
Chestnut-sided Warbler	<i>Setophaga pensylvanica</i>	111
Chimney Swift	<i>Chaetura pelagica</i>	36
Chipping Sparrow	<i>Spizella passerina</i>	116
Chukar	<i>Alectoris chukar</i>	29
Cinnamon Teal	<i>Anas cyanoptera</i>	22
Clark's Grebe	<i>Aechmophorus clarkii</i>	33
Clark's Nutcracker	<i>Nucifraga columbiana</i>	86
Clay-colored Sparrow	<i>Spizella pallida</i>	117
Cliff Chipmunk	<i>Tamias dorsalis</i>	140
Cliff Swallow	<i>Petrochelidon pyrrhonota</i>	88
Columbia Spotted Frog	<i>Rana luteiventris</i>	165
Common Gallinule	<i>Gallinula galeata</i>	40
Common Goldeneye	<i>Bucephala clangula</i>	27
Common Grackle	<i>Quiscalus quiscula</i>	127
Common Gray Fox	<i>Urocyon cinereoargenteus</i>	154
Common Loon	<i>Gavia immer</i>	58
Common Merganser	<i>Mergus merganser</i>	27
Common Muskrat	<i>Ondatra zibethicus</i>	151
Common Nighthawk	<i>Chordeiles minor</i>	35

Common Name	Scientific Name	Page No.
Common Poorwill	<i>Phalaenoptilus nuttallii</i>	36
Common Raven	<i>Corvus corax</i>	87
Common Redpoll	<i>Acanthis flammea</i>	101
Common Shelduck	<i>Tadorna tadorna</i>	28
Common Tern	<i>Sterna hirundo</i>	57
Common Yellowthroat	<i>Geothlypis trichas</i>	109
Connecticut Warbler	<i>Oporornis agilis</i>	108
Cooper's Hawk	<i>Accipiter cooperii</i>	66
Cordilleran Flycatcher	<i>Empidonax occidentalis</i>	79
Coyote	<i>Canis latrans</i>	153
Crested Caracara	<i>Caracara cheriway</i>	76
Dark-eyed Junco	<i>Junco hyemalis</i>	122
Desert Cottontail	<i>Sylvilagus audubonii</i>	138
Desert Striped Whipsnake	<i>Coluber taeniatus taeniatus</i>	170
Dickcissel	<i>Spiza americana</i>	125
Double-crested Cormorant	<i>Phalacrocorax auritus</i>	60
Downy Woodpecker	<i>Picoides pubescens</i>	74
Dunlin	<i>Calidris alpina</i>	45
Dusky Flycatcher	<i>Empidonax oberholseri</i>	79
Dusky Grouse	<i>Dendragapus obscurus</i>	30
Dusky Shrew	<i>Sorex monticolus</i>	132
Dwarf Shrew	<i>Sorex nanus</i>	132
Eared Grebe	<i>Podiceps nigricollis</i>	32
Eastern Bluebird	<i>Sialia sialis</i>	94
Eastern Cottontail	<i>Sylvilagus floridanus</i>	139
Eastern Fox Squirrel	<i>Sciurus niger</i>	143
Eastern Gray Squirrel	<i>Sciurus carolinensis</i>	143
Eastern Kingbird	<i>Tyrannus tyrannus</i>	81
Eastern Meadowlark	<i>Sturnella magna</i>	126
Eastern Mole	<i>Scalopus aquaticus</i>	133

Common Name	Scientific Name	Page No.
Eastern Phoebe	<i>Sayornis phoebe</i>	80
Eastern Red Bat	<i>Lasiurus borealis</i>	135
Eastern Screech-Owl	<i>Megascops asio</i>	69
Eastern Spiny Softshell	<i>Apalone spinifera spinifera</i>	166
Eastern Spotted Skunk	<i>Spilogale putorius</i>	157
Eastern Wood-Pewee	<i>Contopus virens</i>	78
Eastern Yellow-bellied Racer	<i>Coluber constrictor flaviventris</i>	169
Eurasian Collared-Dove	<i>Streptopelia decaocto</i>	34
Eurasian Wigeon	<i>Anas penelope</i>	21
European Goldfinch	<i>Carduelis carduelis</i>	103
European Starling	<i>Sturnus vulgaris</i>	98
Evening Grosbeak	<i>Coccothraustes vespertinus</i>	103
Ferruginous Hawk	<i>Buteo regalis</i>	68
Field Sparrow	<i>Spizella pusilla</i>	117
Fisher	<i>Pekania pennanti</i>	155
Flammulated Owl	<i>Psilosops flammeolus</i>	69
Forster's Tern	<i>Sterna forsteri</i>	58
Fox Sparrow	<i>Passerella iliaca</i>	120
Franklin's Gull	<i>Leucophaeus pipixcan</i>	53
Fringed Myotis	<i>Myotis thysanodes</i>	135
Fulvous Whistling-Duck	<i>Dendrocygna bicolor</i>	18
Gadwall	<i>Anas strepera</i>	21
Garganey	<i>Anas querquedula</i>	23
Glaucous Gull	<i>Larus hyperboreus</i>	56
Glaucous-winged Gull	<i>Larus glaucescens</i>	56
Glossy Ibis	<i>Plegadis falcinellus</i>	63
Golden Eagle	<i>Aquila chrysaetos</i>	68
Golden-crowned Kinglet	<i>Regulus satrapa</i>	94
Golden-crowned Sparrow	<i>Zonotrichia atricapilla</i>	122
Golden-mantled Ground Squirrel	<i>Callospermophilus lateralis</i>	142

Common Name	Scientific Name	Page No.
Golden-winged Warbler	<i>Vermivora chrysoptera</i>	106
Grasshopper Sparrow	<i>Ammodramus savannarum</i>	119
Gray Catbird	<i>Dumetella carolinensis</i>	97
Gray Flycatcher	<i>Empidonax wrightii</i>	79
Gray Jay	<i>Perisoreus canadensis</i>	85
Gray Partridge	<i>Perdix perdix</i>	29
Gray Vireo	<i>Vireo vicinior</i>	83
Gray Wolf	<i>Canis lupus</i>	153
Gray-cheeked Thrush	<i>Catharus minimus</i>	95
Gray-crowned Rosy-Finch	<i>Leucosticte tephrocotis</i>	99
Great Basin Gophersnake	<i>Pituophis catenifer deserticola</i>	171
Great Basin Pocket Mouse	<i>Perognathus mollipilosus</i>	146
Great Basin Skink	<i>Plestiodon skiltonianus utahensis</i>	167
Great Basin Spadefoot	<i>Spea intermontana</i>	163
Great Black-backed Gull	<i>Larus marinus</i>	56
Great Blue Heron	<i>Ardea herodias</i>	61
Great Crested Flycatcher	<i>Myiarchus crinitus</i>	81
Great Egret	<i>Ardea alba</i>	61
Great Gray Owl	<i>Strix nebulosa</i>	71
Great Horned Owl	<i>Bubo virginianus</i>	69
Great Plains Earless Lizard	<i>Holbrookia maculata maculata</i>	169
Great Plains Toad	<i>Anaxyrus cognatus</i>	164
Greater Prairie-Chicken	<i>Tympanuchus cupido</i>	31
Greater Sage-Grouse	<i>Centrocercus urophasianus</i>	30
Greater Scaup	<i>Aythya marila</i>	25
Greater Short-horned Lizard	<i>Phrynosoma hernandesi</i>	168
Greater White-fronted Goose	<i>Anser albifrons</i>	18
Greater Yellowlegs	<i>Tringa melanoleuca</i>	49
Great-tailed Grackle	<i>Quiscalus mexicanus</i>	127
Green Heron	<i>Butorides virescens</i>	62

Common Name	Scientific Name	Page No.
Green-tailed Towhee	<i>Pipilo chlorurus</i>	115
Green-winged Teal	<i>Anas crecca</i>	23
Grizzly Bear	<i>Ursus arctos</i>	154
Gyr Falcon	<i>Falco rusticolus</i>	77
Hairy Woodpecker	<i>Picoides villosus</i>	74
Hammond's Flycatcher	<i>Empidonax hammondii</i>	79
Harlequin Duck	<i>Histrionicus histrionicus</i>	25
Harris's Hawk	<i>Parabuteo unicinctus</i>	66
Harris's Sparrow	<i>Zonotrichia querula</i>	121
Hayden's Shrew	<i>Sorex haydeni</i>	131
Heermann's Gull	<i>Larus heermanni</i>	54
Hepatic Tanager	<i>Piranga flava</i>	122
Hermit Thrush	<i>Catharus guttatus</i>	96
Hermit Warbler	<i>Setophaga occidentalis</i>	114
Herring Gull	<i>Larus argentatus</i>	55
Hispid Pocket Mouse	<i>Chaetodipus hispidus</i>	146
Hoary Bat	<i>Lasiurus cinereus</i>	136
Hoary Redpoll	<i>Acanthis hornemanni</i>	102
Hooded Merganser	<i>Lophodytes cucullatus</i>	27
Hooded Warbler	<i>Setophaga citrina</i>	109
Horned Grebe	<i>Podiceps auritus</i>	32
Horned Lark	<i>Eremophila alpestris</i>	87
House Finch	<i>Haemorhous mexicanus</i>	100
House Mouse	<i>Mus musculus</i>	151
House Sparrow	<i>Passer domesticus</i>	98
House Wren	<i>Troglodytes aedon</i>	91
Hudsonian Godwit	<i>Limosa haemastica</i>	44
Iceland Gull	<i>Larus glaucoides</i>	55
Idaho Pocket Gopher	<i>Thomomys idahoensis</i>	144
Indigo Bunting	<i>Passerina cyanea</i>	125

Common Name	Scientific Name	Page No.
Juniper Titmouse	<i>Baeolophus ridgwayi</i>	89
Kentucky Warbler	<i>Geothlypis formosus</i>	109
Killdeer	<i>Charadrius vociferus</i>	43
Lapland Longspur	<i>Calcarius lapponicus</i>	104
Lark Bunting	<i>Calamospiza melanocorys</i>	118
Lark Sparrow	<i>Chondestes grammacus</i>	118
Laughing Gull	<i>Leucophaeus atricilla</i>	53
Lawrence's Goldfinch	<i>Spinus lawrencei</i>	102
Lazuli Bunting	<i>Passerina amoena</i>	124
Le Conte's Sparrow	<i>Ammodramus leconteii</i>	119
Least Bittern	<i>Ixobrychus exilis</i>	61
Least Chipmunk	<i>Tamias minimus</i>	140
Least Flycatcher	<i>Empidonax minimus</i>	78
Least Sandpiper	<i>Calidris minutilla</i>	46
Least Tern	<i>Sternula antillarum</i>	56
Least Weasel	<i>Mustela nivalis</i>	156
Lesser Black-backed Gull	<i>Larus fuscus</i>	55
Lesser Frigatebird	<i>Fregata ariel</i>	59
Lesser Goldfinch	<i>Spinus psaltria</i>	102
Lesser Nighthawk	<i>Chordeiles acutipennis</i>	35
Lesser Scaup	<i>Aythya affinis</i>	25
Lesser Yellowlegs	<i>Tringa flavipes</i>	49
Lewis's Woodpecker	<i>Melanerpes lewis</i>	72
Lincoln's Sparrow	<i>Melospiza lincolnii</i>	120
Little Blue Heron	<i>Egretta caerulea</i>	62
Little Brown Myotis	<i>Myotis lucifugus</i>	134
Little Gull	<i>Hydrocoloeus minutus</i>	53
Loggerhead Shrike	<i>Lanius ludovicianus</i>	82
Long-billed Curlew	<i>Numenius americanus</i>	44
Long-billed Dowitcher	<i>Limnodromus scolopaceus</i>	48

Common Name	Scientific Name	Page No.
Long-billed Murrelet	<i>Brachyramphus perdix</i>	51
Long-eared Myotis	<i>Myotis evotis</i>	134
Long-eared Owl	<i>Asio otus</i>	71
Long-legged Myotis	<i>Myotis volans</i>	135
Long-tailed Duck	<i>Clangula hyemalis</i>	26
Long-tailed Jaeger	<i>Stercorarius longicaudus</i>	51
Long-tailed Vole	<i>Microtus longicaudus</i>	149
Long-tailed Weasel	<i>Mustela frenata</i>	156
MacGillivray's Warbler	<i>Geothlypis tolmiei</i>	108
Magnificent Hummingbird	<i>Eugenes fulgens</i>	37
Magnolia Warbler	<i>Setophaga magnolia</i>	110
Mallard	<i>Anas platyrhynchos</i>	22
Marbled Godwit	<i>Limosa fedoa</i>	44
Marsh Wren	<i>Cistothorus palustris</i>	92
Masked Shrew	<i>Sorex cinereus</i>	131
McCown's Longspur	<i>Rhynchophanes mccownii</i>	105
Meadow Jumping Mouse	<i>Zapus hudsonius</i>	152
Meadow Vole	<i>Microtus pennsylvanicus</i>	150
Merlin	<i>Falco columbarius</i>	76
Merriam's Shrew	<i>Sorex merriami</i>	132
Mew Gull	<i>Larus canus</i>	54
Midget Faded Rattlesnake	<i>Crotalus oreganus concolor</i>	173
Mississippi Kite	<i>Ictinia mississippiensis</i>	65
Montane Vole	<i>Microtus montanus</i>	150
Moose	<i>Alces americanus</i>	160
Mottled Duck	<i>Anas fulvigula</i>	22
Mountain Bluebird	<i>Sialia currucoides</i>	95
Mountain Chickadee	<i>Poecile gambeli</i>	89
Mountain Cottontail	<i>Sylvilagus nutallii</i>	139
Mountain Goat	<i>Oreamnos americanus</i>	160

Common Name	Scientific Name	Page No.
Mountain Lion (Puma)	<i>Puma concolor</i>	158
Mountain Plover	<i>Charadrius montanus</i>	43
Mourning Dove	<i>Zenaida macroura</i>	34
Mourning Warbler	<i>Geothlypis philadelphia</i>	108
Mule Deer (Black-tailed Deer)	<i>Odocoileus hemionus</i>	159
Mute Swan	<i>Cygnus olor</i>	19
Nashville Warbler	<i>Oreothlypis ruficapilla</i>	107
Nelson's Sparrow	<i>Ammodramus nelsoni</i>	120
North American Deermouse	<i>Peromyscus maniculatus</i>	148
North American Porcupine	<i>Erethizon dorsatum</i>	152
Northern Bobwhite	<i>Colinus virginianus</i>	29
Northern Cardinal	<i>Cardinalis cardinalis</i>	123
Northern Flicker	<i>Colaptes auratus</i>	75
Northern Flying Squirrel	<i>Glaucomys sabrinus</i>	144
Northern Goshawk	<i>Accipiter gentilis</i>	66
Northern Grasshopper Mouse	<i>Onychomys leucogaster</i>	148
Northern Harrier	<i>Circus cyaneus</i>	65
Northern Hawk Owl	<i>Surnia ulula</i>	70
Northern Leopard Frog	<i>Lithobates pipiens</i>	165
Northern Long-eared Myotis	<i>Myotis septentrionalis</i>	134
Northern Many-lined Skink	<i>Plestiodon multivirgatus multivirgatus</i>	167
Northern Mockingbird	<i>Mimus polyglottos</i>	98
Northern Parula	<i>Setophaga americana</i>	110
Northern Pintail	<i>Anas acuta</i>	23
Northern Pocket Gopher	<i>Thomomys talpoides</i>	145
Northern Pygmy-Owl	<i>Glaucidium gnoma</i>	70
Northern Raccoon	<i>Procyon lotor</i>	155
Northern River Otter	<i>Lontra canadensis</i>	158
Northern Rough-winged Swallow	<i>Stelgidopteryx serripennis</i>	88
Northern Rubber Boa	<i>Charina bottae</i>	169

Common Name	Scientific Name	Page No.
Northern Sagebrush Lizard	<i>Sceloporus graciosus graciosus</i>	167
Northern Saw-whet Owl	<i>Aegolius acadicus</i>	72
Northern Shoveler	<i>Anas clypeata</i>	23
Northern Shrike	<i>Lanius excubitor</i>	82
Northern Tree Lizard	<i>Urosaurus ornatus wrighti</i>	168
Northern Waterthrush	<i>Parkesia noveboracensis</i>	106
Norway Rat	<i>Rattus norvegicus</i>	151
Olive-backed Pocket Mouse	<i>Perognathus fasciatus</i>	145
Olive-sided Flycatcher	<i>Contopus cooperi</i>	77
Orange-crowned Warbler	<i>Oreothlypis celata</i>	107
Orchard Oriole	<i>Icterus spurius</i>	128
Ord's Kangaroo Rat	<i>Dipodomys ordii</i>	146
Osprey	<i>Pandion haliaetus</i>	64
Ovenbird	<i>Seiurus aurocapilla</i>	105
Pacific Loon	<i>Gavia pacifica</i>	58
Pacific Marten	<i>Martes caurina</i>	155
Pacific Wren	<i>Troglodytes pacificus</i>	92
Painted Bunting	<i>Passerina ciris</i>	125
Pale Milksnake	<i>Lampropeltis triangulum multistriata</i>	170
Pallid Bat	<i>Antrozous pallidus</i>	137
Palm Warbler	<i>Setophaga palmarum</i>	112
Parasitic Jaeger	<i>Stercorarius parasiticus</i>	51
Passenger Pigeon	<i>Ectopistes migratorius</i>	34
Pectoral Sandpiper	<i>Calidris melanotos</i>	47
Peregrine Falcon	<i>Falco peregrinus</i>	77
Philadelphia Vireo	<i>Vireo philadelphicus</i>	84
Pied-billed Grebe	<i>Podilymbus podiceps</i>	31
Pileated Woodpecker	<i>Dryocopus pileatus</i>	76
Pine Grosbeak	<i>Pinicola enucleator</i>	100
Pine Siskin	<i>Spinus pinus</i>	102

Common Name	Scientific Name	Page No.
Pine Warbler	<i>Setophaga pinus</i>	112
Piñon Deermouse	<i>Peromyscus truei</i>	148
Pinyon Jay	<i>Gymnorhinus cyanocephalus</i>	85
Piping Plover	<i>Charadrius melodus</i>	42
Plains Black-headed Snake	<i>Tantilla nigriceps</i>	171
Plains Box Turtle	<i>Terrapene ornata ornata</i>	166
Plains Gartersnake	<i>Thamnophis radix</i>	172
Plains Harvest Mouse	<i>Reithrodontomys montanus</i>	147
Plains Hog-nosed Snake	<i>Heterodon nasicus</i>	169
Plains Pocket Mouse	<i>Perognathus flavescens</i>	145
Plains Spadefoot	<i>Spea bombifrons</i>	163
Plateau Fence Lizard	<i>Sceloporus tristichus</i>	168
Plumbeous Vireo	<i>Vireo plumbeus</i>	84
Pomarine Jaeger	<i>Stercorarius pomarinus</i>	50
Prairie Falcon	<i>Falco mexicanus</i>	77
Prairie Lizard	<i>Sceloporus consobrinus</i>	168
Prairie Racerunner	<i>Aspidoscelis sexlineata viridis</i>	167
Prairie Rattlesnake	<i>Crotalus viridis</i>	172
Prairie Vole	<i>Microtus ochrogaster</i>	150
Prairie Warbler	<i>Setophaga discolor</i>	113
Preble's Meadow Jumping Mouse	<i>Zapus hudsonius preblei</i>	152
Preble's Shrew	<i>Sorex preblei</i>	133
Pronghorn	<i>Antilocapra americana</i>	160
Prothonotary Warbler	<i>Protonotaria citrea</i>	107
Purple Finch	<i>Haemorhous purpureus</i>	100
Purple Gallinule	<i>Porphyrio martinicus</i>	40
Purple Martin	<i>Progne subis</i>	87
Pygmy Nuthatch	<i>Sitta pygmaea</i>	90
Pygmy Rabbit	<i>Brachylagus idahoensis</i>	138
Red Crossbill	<i>Loxia curvirostra</i>	101

Common Name	Scientific Name	Page No.
Red Fox	<i>Vulpes vulpes</i>	153
Red Knot	<i>Calidris canutus</i>	45
Red Phalarope	<i>Phalaropus fulicarius</i>	50
Red Squirrel	<i>Tamiasciurus hudsonicus</i>	144
Red-bellied Woodpecker	<i>Melanerpes carolinus</i>	73
Red-breasted Merganser	<i>Mergus serrator</i>	28
Red-breasted Nuthatch	<i>Sitta canadensis</i>	90
Red-eyed Vireo	<i>Vireo olivaceus</i>	84
Red-faced Warbler	<i>Cardellina rubrifrons</i>	115
Redhead	<i>Aythya americana</i>	24
Red-headed Woodpecker	<i>Melanerpes erythrocephalus</i>	73
Red-naped Sapsucker	<i>Sphyrapicus nuchalis</i>	74
Red-necked Grebe	<i>Podiceps grisegena</i>	32
Red-necked Phalarope	<i>Phalaropus lobatus</i>	50
Red-shouldered Hawk	<i>Buteo lineatus</i>	67
Red-sided Gartersnake	<i>Thamnophis sirtalis parietalis</i>	172
Red-tailed Hawk	<i>Buteo jamaicensis</i>	67
Red-throated Loon	<i>Gavia stellata</i>	58
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	126
Ring-billed Gull	<i>Larus delawarensis</i>	54
Ring-necked Duck	<i>Aythya collaris</i>	24
Ring-necked Pheasant	<i>Phasianus colchicus</i>	29
Ringtail	<i>Bassariscus astutus</i>	154
Rock Pigeon	<i>Columba livia</i>	33
Rock Wren	<i>Salpinctes obsoletus</i>	91
Rocky Mountain Toad (Woodhouse's Toad)	<i>Anaxyrus woodhousii woodhousii</i>	164
Rose-breasted Grosbeak	<i>Pheucticus ludovicianus</i>	124
Ross's Goose	<i>Chen rossii</i>	18
Ross's Gull	<i>Rhodostethia rosea</i>	53
Rough-legged Hawk	<i>Buteo lagopus</i>	68

Common Name	Scientific Name	Page No.
Ruby-crowned Kinglet	<i>Regulus calendula</i>	94
Ruby-throated Hummingbird	<i>Archilochus colubris</i>	37
Ruddy Duck	<i>Oxyura jamaicensis</i>	28
Ruddy Shelduck	<i>Tadorna ferruginea</i>	28
Ruddy Turnstone	<i>Arenaria interpres</i>	44
Ruffed Grouse	<i>Bonasa umbellus</i>	30
Rufous Hummingbird	<i>Selasphorus rufus</i>	38
Rusty Blackbird	<i>Euphagus carolinus</i>	127
Sabine's Gull	<i>Xema sabini</i>	52
Sage Thrasher	<i>Oreoscoptes montanus</i>	97
Sagebrush Sparrow	<i>Artemisospiza nevadensis</i>	118
Sagebrush Vole	<i>Lemmyscus curtatus</i>	151
Sand Hills Pocket Gopher	<i>Geomys lutescens</i>	145
Sanderling	<i>Calidris alba</i>	45
Sandhill Crane	<i>Antigone canadensis</i>	40
Savannah Sparrow	<i>Passerculus sandwichensis</i>	119
Say's Phoebe	<i>Sayornis saya</i>	80
Scarlet Tanager	<i>Piranga olivacea</i>	123
Scissor-tailed Flycatcher	<i>Tyrannus forficatus</i>	82
Scott's Oriole	<i>Icterus parisorum</i>	129
Sedge Wren	<i>Cistothorus platensis</i>	92
Semipalmated Plover	<i>Charadrius semipalmatus</i>	42
Semipalmated Sandpiper	<i>Calidris pusilla</i>	47
Sharp-shinned Hawk	<i>Accipiter striatus</i>	66
Sharp-tailed Grouse	<i>Tympanuchus phasianellus</i>	31
Short-billed Dowitcher	<i>Limnodromus griseus</i>	47
Short-eared Owl	<i>Asio flammeus</i>	71
Short-tailed Weasel (Ermine)	<i>Mustela erminea</i>	155
Silky Pocket Mouse	<i>Perognathus flavus</i>	146
Silver-haired Bat	<i>Lasionycteris noctivagans</i>	136

Common Name	Scientific Name	Page No.
Smith's Longspur	<i>Calcarius pictus</i>	104
Smooth Greensnake	<i>Opheodrys vernalis</i>	170
Snapping Turtle	<i>Chelydra serpentina</i>	166
Snow Bunting	<i>Plectrophenax nivalis</i>	105
Snow Goose	<i>Chen caerulescens</i>	18
Snowshoe Hare	<i>Lepus americanus</i>	139
Snowy Egret	<i>Egretta thula</i>	61
Snowy Owl	<i>Bubo scandiacus</i>	70
Snowy Plover	<i>Charadrius nivosus</i>	42
Solitary Sandpiper	<i>Tringa solitaria</i>	49
Song Sparrow	<i>Melospiza melodia</i>	120
Sora	<i>Porzana carolina</i>	39
Southern Red-backed Vole	<i>Myodes gapperi</i>	149
Spotted Bat	<i>Euderma maculatum</i>	137
Spotted Ground Squirrel	<i>Xerospermophilus spilosoma</i>	142
Spotted Sandpiper	<i>Actitis macularius</i>	48
Spotted Towhee	<i>Pipilo maculatus</i>	115
Sprague's Pipit	<i>Anthus spragueii</i>	99
Steller's Jay	<i>Cyanocitta stelleri</i>	85
Stilt Sandpiper	<i>Calidris himantopus</i>	45
Streaked Shearwater	<i>Calonectris leucomelas</i>	59
Striped Skunk	<i>Mephitis mephitis</i>	158
Summer Tanager	<i>Piranga rubra</i>	122
Surf Scoter	<i>Melanitta perspicillata</i>	25
Swainson's Hawk	<i>Buteo swainsoni</i>	67
Swainson's Thrush	<i>Catharus ustulatus</i>	96
Swamp Sparrow	<i>Melospiza georgiana</i>	121
Swift Fox	<i>Vulpes velox</i>	153
Tennessee Warbler	<i>Oreothlypis peregrina</i>	107
Thayer's Gull	<i>Larus thayeri</i>	55

Common Name	Scientific Name	Page No.
Thirteen-lined Ground Squirrel	<i>Ictidomys tridecemlineatus</i>	142
Tiger Salamander	<i>Ambystoma mavortium</i>	163
Townsend's Big-eared Bat	<i>Corynorhinus townsendii</i>	137
Townsend's Solitaire	<i>Myadestes townsendi</i>	95
Townsend's Warbler	<i>Setophaga townsendi</i>	113
Tree Swallow	<i>Tachycineta bicolor</i>	87
Tricolored Heron	<i>Egretta tricolor</i>	62
Trumpeter Swan	<i>Cygnus buccinator</i>	20
Tufted Duck	<i>Aythya fuligula</i>	24
Tundra Swan	<i>Cygnus columbianus</i>	20
Turkey Vulture	<i>Cathartes aura</i>	64
Uinta Chipmunk	<i>Tamias umbrinus</i>	141
Uinta Ground Squirrel	<i>Urocitellus armatus</i>	141
Upland Sandpiper	<i>Bartramia longicauda</i>	43
Vagrant Shrew	<i>Sorex vagrans</i>	133
Valley Gartersnake	<i>Thamnophis sirtalis fitchi</i>	172
Varied Thrush	<i>Ixoreus naevius</i>	97
Vaux's Swift	<i>Chaetura vauxi</i>	36
Veery	<i>Catharus fuscescens</i>	95
Vermilion Flycatcher	<i>Pyrocephalus rubinus</i>	80
Vesper Sparrow	<i>Poocetes gramineus</i>	117
Violet-green Swallow	<i>Tachycineta thalassina</i>	88
Virginia Opossum	<i>Didelphis virginiana</i>	131
Virginia Rail	<i>Rallus limicola</i>	39
Virginia's Warbler	<i>Oreothlypis virginiae</i>	108
Wandering Gartersnake	<i>Thamnophis elegans vagrans</i>	171
Wapiti (Elk)	<i>Cervus canadensis</i>	159
Warbling Vireo	<i>Vireo gilvus</i>	84
Water Vole	<i>Microtus richardsoni</i>	150
Western Bluebird	<i>Sialia mexicana</i>	94

Common Name	Scientific Name	Page No.
Western Grebe	<i>Aechmophorus occidentalis</i>	32
Western Harvest Mouse	<i>Reithrodontomys megalotis</i>	147
Western Heather Vole	<i>Phenacomys intermedius</i>	149
Western Jumping Mouse	<i>Zapus princeps</i>	152
Western Kingbird	<i>Tyrannus verticalis</i>	81
Western Meadowlark	<i>Sturnella neglecta</i>	126
Western Painted Turtle	<i>Chrysemys picta bellii</i>	166
Western Sandpiper	<i>Calidris mauri</i>	47
Western Screech-Owl	<i>Megascops kennicottii</i>	69
Western Small-footed Myotis	<i>Myotis ciliolabrum</i>	134
Western Spotted Skunk	<i>Spilogale gracilis</i>	157
Western Tanager	<i>Piranga ludoviciana</i>	123
Western Toad (Boreal Toad)	<i>Anaxyrus boreas</i>	163
Western Water Shrew	<i>Sorex navigator</i>	132
Western Wood-Pewee	<i>Contopus sordidulus</i>	78
Whimbrel	<i>Numenius phaeopus</i>	43
White Ibis	<i>Eudocimus albus</i>	63
White-breasted Nuthatch	<i>Sitta carolinensis</i>	90
White-crowned Sparrow	<i>Zonotrichia leucophrys</i>	121
White-eyed Vireo	<i>Vireo griseus</i>	82
White-faced Ibis	<i>Plegadis chihi</i>	64
White-footed Deermouse	<i>Peromyscus leucopus</i>	148
White-headed Woodpecker	<i>Picoides albolarvatus</i>	75
White-rumped Sandpiper	<i>Calidris fuscicollis</i>	46
White-tailed Deer	<i>Odocoileus virginianus</i>	159
White-tailed Jackrabbit	<i>Lepus townsendii</i>	140
White-tailed Kite	<i>Elanus leucurus</i>	65
White-tailed Prairie Dog	<i>Cynomys leucurus</i>	142
White-tailed Ptarmigan	<i>Lagopus leucura</i>	30
White-throated Sparrow	<i>Zonotrichia albicollis</i>	121

Common Name	Scientific Name	Page No.
White-throated Swift	<i>Aeronautes saxatalis</i>	37
White-winged Crossbill	<i>Loxia leucoptera</i>	101
White-winged Dove	<i>Zenaida asiatica</i>	34
White-winged Scoter	<i>Melanitta fusca</i>	26
Whooper Swan	<i>Cygnus cygnus</i>	20
Whooping Crane	<i>Grus americana</i>	41
Wild Turkey	<i>Meleagris gallopavo</i>	31
Willet	<i>Tringa semipalmata</i>	49
Williamson's Sapsucker	<i>Sphyrapicus thyroideus</i>	73
Willow Flycatcher	<i>Empidonax traillii</i>	78
Wilson's Phalarope	<i>Phalaropus tricolor</i>	50
Wilson's Snipe	<i>Gallinago delicata</i>	48
Wilson's Warbler	<i>Cardellina pusilla</i>	114
Winter Wren	<i>Troglodytes hiemalis</i>	92
Wolverine	<i>Gulo gulo</i>	157
Wood Duck	<i>Aix sponsa</i>	20
Wood Frog	<i>Lithobates sylvaticus</i>	165
Wood Stork	<i>Mycteria americana</i>	59
Wood Thrush	<i>Hylocichla mustelina</i>	96
Woodhouse's Scrub-Jay	<i>Aphelocoma woodhouseii</i>	86
Worm-eating Warbler	<i>Helmitheros vermivorum</i>	105
Wyoming Ground Squirrel	<i>Urocitellus elegans</i>	141
Wyoming Pocket Gopher	<i>Thomomys clusius</i>	144
Wyoming Toad	<i>Anaxyrus baxteri</i>	164
Yellow Grosbeak	<i>Pheucticus chrysopheplus</i>	123
Yellow Rail	<i>Coturnicops noveboracensis</i>	39
Yellow Warbler	<i>Setophaga petechia</i>	111
Yellow-bellied Marmot	<i>Marmota flaviventris</i>	141
Yellow-bellied Sapsucker	<i>Sphyrapicus varius</i>	74
Yellow-billed Cuckoo	<i>Coccyzus americanus</i>	35

Common Name	Scientific Name	Page No.
Yellow-billed Loon	<i>Gavia adamsii</i>	59
Yellow-breasted Chat	<i>Icteria virens</i>	115
Yellow-crowned Night-Heron	<i>Nyctanassa violacea</i>	63
Yellow-headed Blackbird	<i>Xanthocephalus xanthocephalus</i>	126
Yellow-pine Chipmunk	<i>Tamias amoenus</i>	140
Yellow-rumped Warbler	<i>Setophaga coronata</i>	112
Yellow-throated Vireo	<i>Vireo flavifrons</i>	83
Yellow-throated Warbler	<i>Setophaga dominica</i>	113
Yuma Myotis	<i>Myotis yumanensis</i>	135

Index of Scientific Names

Scientific Name	Common Name	Page No.
<i>Coragyps atratus</i>	Black Vulture	64
<i>Acanthis flammea</i>	Common Redpoll	101
<i>Acanthis hornemanni</i>	Hoary Redpoll	102
<i>Accipiter cooperii</i>	Cooper's Hawk	66
<i>Accipiter gentilis</i>	Northern Goshawk	66
<i>Accipiter striatus</i>	Sharp-shinned Hawk	66
<i>Actitis macularius</i>	Spotted Sandpiper	48
<i>Aechmophorus clarkii</i>	Clark's Grebe	33
<i>Aechmophorus occidentalis</i>	Western Grebe	32
<i>Aegolius acadicus</i>	Northern Saw-whet Owl	72
<i>Aegolius funereus</i>	Boreal Owl	72
<i>Aeronautes saxatalis</i>	White-throated Swift	37
<i>Agelaius phoeniceus</i>	Red-winged Blackbird	126
<i>Aix sponsa</i>	Wood Duck	20
<i>Alces americanus</i>	Moose	160
<i>Alectoris chukar</i>	Chukar	29
<i>Ambystoma mavortium</i>	Tiger Salamander	163
<i>Ammodramus bairdii</i>	Baird's Sparrow	119
<i>Ammodramus leconteii</i>	Le Conte's Sparrow	119
<i>Ammodramus nelsoni</i>	Nelson's Sparrow	120
<i>Ammodramus savannarum</i>	Grasshopper Sparrow	119
<i>Amphispiza bilineata</i>	Black-throated Sparrow	118
<i>Anas acuta</i>	Northern Pintail	23
<i>Anas americana</i>	American Wigeon	21
<i>Anas clypeata</i>	Northern Shoveler	23
<i>Anas crecca</i>	Green-winged Teal	23
<i>Anas cyanoptera</i>	Cinnamon Teal	22
<i>Anas discors</i>	Blue-winged Teal	22

Scientific Name	Common Name	Page No.
<i>Anas fulvigula</i>	Mottled Duck	22
<i>Anas penelope</i>	Eurasian Wigeon	21
<i>Anas platyrhynchos</i>	Mallard	22
<i>Anas querquedula</i>	Garganey	23
<i>Anas rubripes</i>	American Black Duck	21
<i>Anas strepera</i>	Gadwall	21
<i>Anaxyrus baxteri</i>	Wyoming Toad	164
<i>Anaxyrus boreas</i>	Western Toad (Boreal Toad)	163
<i>Anaxyrus cognatus</i>	Great Plains Toad	164
<i>Anaxyrus woodhousii woodhousii</i>	Rocky Mountain Toad (Woodhouse's Toad)	164
<i>Anser albifrons</i>	Greater White-fronted Goose	18
<i>Anthus rubescens</i>	American Pipit	98
<i>Anthus spragueii</i>	Sprague's Pipit	99
<i>Antigone canadensis</i>	Sandhill Crane	40
<i>Antilocapra americana</i>	Pronghorn	160
<i>Antrozous pallidus</i>	Pallid Bat	137
<i>Apalone spinifera spinifera</i>	Eastern Spiny Softshell	166
<i>Aphelocoma woodhouseii</i>	Woodhouse's Scrub-Jay	86
<i>Aquila chrysaetos</i>	Golden Eagle	68
<i>Archilochus alexandri</i>	Black-chinned Hummingbird	37
<i>Archilochus colubris</i>	Ruby-throated Hummingbird	37
<i>Ardea alba</i>	Great Egret	61
<i>Ardea herodias</i>	Great Blue Heron	61
<i>Arenaria interpres</i>	Ruddy Turnstone	44
<i>Artemisiospiza nevadensis</i>	Sagebrush Sparrow	118
<i>Asio flammeus</i>	Short-eared Owl	71
<i>Asio otus</i>	Long-eared Owl	71
<i>Aspidoscelis sexlineata viridis</i>	Prairie Racerunner	167
<i>Athene cunicularia</i>	Burrowing Owl	70
<i>Aythya affinis</i>	Lesser Scaup	25

Scientific Name	Common Name	Page No.
<i>Aythya americana</i>	Redhead	24
<i>Aythya collaris</i>	Ring-necked Duck	24
<i>Aythya fuligula</i>	Tufted Duck	24
<i>Aythya marila</i>	Greater Scaup	25
<i>Aythya valisineria</i>	Canvasback	24
<i>Baeolophus ridgwayi</i>	Juniper Titmouse	89
<i>Bartramia longicauda</i>	Upland Sandpiper	43
<i>Bassariscus astutus</i>	Ringtail	154
<i>Bombycilla cedrorum</i>	Cedar Waxwing	104
<i>Bombycilla garrulus</i>	Bohemian Waxwing	103
<i>Bonasa umbellus</i>	Ruffed Grouse	30
<i>Bos bison</i>	Bison	160
<i>Botaurus lentiginosus</i>	American Bittern	60
<i>Brachylagus idahoensis</i>	Pygmy Rabbit	138
<i>Brachyramphus perdix</i>	Long-billed Murrelet	51
<i>Branta bernicla</i>	Brant	19
<i>Branta canadensis</i>	Canada Goose	19
<i>Branta hutchinsii</i>	Cackling Goose	19
<i>Bubo scandiacus</i>	Snowy Owl	70
<i>Bubo virginianus</i>	Great Horned Owl	69
<i>Bubulcus ibis</i>	Cattle Egret	62
<i>Bucephala albeola</i>	Bufflehead	26
<i>Bucephala clangula</i>	Common Goldeneye	27
<i>Bucephala islandica</i>	Barrow's Goldeneye	27
<i>Buteo jamaicensis</i>	Red-tailed Hawk	67
<i>Buteo lagopus</i>	Rough-legged Hawk	68
<i>Buteo lineatus</i>	Red-shouldered Hawk	67
<i>Buteo platypterus</i>	Broad-winged Hawk	67
<i>Buteo regalis</i>	Ferruginous Hawk	68
<i>Buteo swainsoni</i>	Swainson's Hawk	67

Scientific Name	Common Name	Page No.
<i>Butorides virescens</i>	Green Heron	62
<i>Calamospiza melanocorys</i>	Lark Bunting	118
<i>Calcarius lapponicus</i>	Lapland Longspur	104
<i>Calcarius ornatus</i>	Chestnut-collared Longspur	104
<i>Calcarius pictus</i>	Smith's Longspur	104
<i>Calidris alba</i>	Sanderling	45
<i>Calidris alpina</i>	Dunlin	45
<i>Calidris bairdii</i>	Baird's Sandpiper	46
<i>Calidris canutus</i>	Red Knot	45
<i>Calidris fuscicollis</i>	White-rumped Sandpiper	46
<i>Calidris himantopus</i>	Stilt Sandpiper	45
<i>Calidris mauri</i>	Western Sandpiper	47
<i>Calidris melanotos</i>	Pectoral Sandpiper	47
<i>Calidris minutilla</i>	Least Sandpiper	46
<i>Calidris pusilla</i>	Semipalmated Sandpiper	47
<i>Calidris subruficollis</i>	Buff-breasted Sandpiper	46
<i>Callospermophilus lateralis</i>	Golden-mantled Ground Squirrel	142
<i>Calonectris leucomelas</i>	Streaked Shearwater	59
<i>Calypte anna</i>	Anna's Hummingbird	38
<i>Canis latrans</i>	Coyote	153
<i>Canis lupus</i>	Gray Wolf	153
<i>Caracara cheriway</i>	Crested Caracara	76
<i>Cardellina canadensis</i>	Canada Warbler	114
<i>Cardellina pusilla</i>	Wilson's Warbler	114
<i>Cardellina rubrifrons</i>	Red-faced Warbler	115
<i>Cardinalis cardinalis</i>	Northern Cardinal	123
<i>Carduelis carduelis</i>	European Goldfinch	103
<i>Castor canadensis</i>	Beaver	147
<i>Cathartes aura</i>	Turkey Vulture	64
<i>Catharus fuscescens</i>	Veery	95

Scientific Name	Common Name	Page No.
<i>Catharus guttatus</i>	Hermit Thrush	96
<i>Catharus minimus</i>	Gray-cheeked Thrush	95
<i>Catharus ustulatus</i>	Swainson's Thrush	96
<i>Catherpes mexicanus</i>	Canyon Wren	91
<i>Centrocercus urophasianus</i>	Greater Sage-Grouse	30
<i>Certhia americana</i>	Brown Creeper	91
<i>Cervus canadensis</i>	Wapiti (Elk)	159
<i>Chaetodipus hispidus</i>	Hispid Pocket Mouse	146
<i>Chaetura pelagica</i>	Chimney Swift	36
<i>Chaetura vauxi</i>	Vaux's Swift	36
<i>Charadrius melodus</i>	Piping Plover	42
<i>Charadrius montanus</i>	Mountain Plover	43
<i>Charadrius nivosus</i>	Snowy Plover	42
<i>Charadrius semipalmatus</i>	Semipalmated Plover	42
<i>Charadrius vociferus</i>	Killdeer	43
<i>Charina bottae</i>	Northern Rubber Boa	169
<i>Chelydra serpentina</i>	Snapping Turtle	166
<i>Chen caerulescens</i>	Snow Goose	18
<i>Chen rossii</i>	Ross's Goose	18
<i>Chlidonias niger</i>	Black Tern	57
<i>Chondestes grammacus</i>	Lark Sparrow	118
<i>Chordeiles acutipennis</i>	Lesser Nighthawk	35
<i>Chordeiles minor</i>	Common Nighthawk	35
<i>Chroicocephalus philadelphia</i>	Bonaparte's Gull	52
<i>Chroicocephalus ridibundus</i>	Black-headed Gull	52
<i>Chrysemys picta bellii</i>	Western Painted Turtle	166
<i>Cinclus mexicanus</i>	American Dipper	93
<i>Circus cyaneus</i>	Northern Harrier	65
<i>Cistothorus palustris</i>	Marsh Wren	92
<i>Cistothorus platensis</i>	Sedge Wren	92

Scientific Name	Common Name	Page No.
<i>Clangula hyemalis</i>	Long-tailed Duck	26
<i>Coccothraustes vespertinus</i>	Evening Grosbeak	103
<i>Coccyzus americanus</i>	Yellow-billed Cuckoo	35
<i>Coccyzus erythrophthalmus</i>	Black-billed Cuckoo	35
<i>Colaptes auratus</i>	Northern Flicker	75
<i>Colinus virginianus</i>	Northern Bobwhite	29
<i>Coluber constrictor flaviventris</i>	Eastern Yellow-bellied Racer	169
<i>Coluber taeniatus taeniatus</i>	Desert Striped Whipsnake	170
<i>Columba livia</i>	Rock Pigeon	33
<i>Contopus cooperi</i>	Olive-sided Flycatcher	77
<i>Contopus sordidulus</i>	Western Wood-Pewee	78
<i>Contopus virens</i>	Eastern Wood-Pewee	78
<i>Corvus brachyrhynchos</i>	American Crow	86
<i>Corvus corax</i>	Common Raven	87
<i>Corynorhinus townsendii</i>	Townsend's Big-eared Bat	137
<i>Coturnicops noveboracensis</i>	Yellow Rail	39
<i>Crotalus oreganus concolor</i>	Midget Faded Rattlesnake	173
<i>Crotalus viridis</i>	Prairie Rattlesnake	172
<i>Cyanocitta cristata</i>	Blue Jay	85
<i>Cyanocitta stelleri</i>	Steller's Jay	85
<i>Cygnus buccinator</i>	Trumpeter Swan	20
<i>Cygnus columbianus</i>	Tundra Swan	20
<i>Cygnus cygnus</i>	Whooper Swan	20
<i>Cygnus olor</i>	Mute Swan	19
<i>Cynomys leucurus</i>	White-tailed Prairie Dog	142
<i>Cynomys ludovicianus</i>	Black-tailed Prairie Dog	143
<i>Cypseloides niger</i>	Black Swift	36
<i>Dendragapus obscurus</i>	Dusky Grouse	30
<i>Dendrocygna bicolor</i>	Fulvous Whistling-Duck	18
<i>Didelphis virginiana</i>	Virginia Opossum	131

Scientific Name	Common Name	Page No.
<i>Dipodomys ordii</i>	Ord's Kangaroo Rat	146
<i>Dolichonyx oryzivorus</i>	Bobolink	125
<i>Dryocopus pileatus</i>	Pileated Woodpecker	76
<i>Dumetella carolinensis</i>	Gray Catbird	97
<i>Ectopistes migratorius</i>	Passenger Pigeon	34
<i>Egretta caerulea</i>	Little Blue Heron	62
<i>Egretta thula</i>	Snowy Egret	61
<i>Egretta tricolor</i>	Tricolored Heron	62
<i>Elanus leucurus</i>	White-tailed Kite	65
<i>Empidonax hammondi</i>	Hammond's Flycatcher	79
<i>Empidonax minimus</i>	Least Flycatcher	78
<i>Empidonax oberholseri</i>	Dusky Flycatcher	79
<i>Empidonax occidentalis</i>	Cordilleran Flycatcher	79
<i>Empidonax traillii</i>	Willow Flycatcher	78
<i>Empidonax wrightii</i>	Gray Flycatcher	79
<i>Eptesicus fuscus</i>	Big Brown Bat	136
<i>Eremophila alpestris</i>	Horned Lark	87
<i>Erethizon dorsatum</i>	North American Porcupine	152
<i>Euderma maculatum</i>	Spotted Bat	137
<i>Eudocimus albus</i>	White Ibis	63
<i>Eugenes fulgens</i>	Magnificent Hummingbird	37
<i>Euphagus carolinus</i>	Rusty Blackbird	127
<i>Euphagus cyanocephalus</i>	Brewer's Blackbird	127
<i>Falco columbarius</i>	Merlin	76
<i>Falco mexicanus</i>	Prairie Falcon	77
<i>Falco peregrinus</i>	Peregrine Falcon	77
<i>Falco rusticolus</i>	Gyrfalcon	77
<i>Falco sparverius</i>	American Kestrel	76
<i>Fregata ariel</i>	Lesser Frigatebird	59
<i>Fringilla montifringilla</i>	Brambling	99

Scientific Name	Common Name	Page No.
<i>Fulica americana</i>	American Coot	40
<i>Gallinago delicata</i>	Wilson's Snipe	48
<i>Gallinula galeata</i>	Common Gallinule	40
<i>Gavia adamsii</i>	Yellow-billed Loon	59
<i>Gavia immer</i>	Common Loon	58
<i>Gavia pacifica</i>	Pacific Loon	58
<i>Gavia stellata</i>	Red-throated Loon	58
<i>Geomys lutescens</i>	Sand Hills Pocket Gopher	145
<i>Geothlypis formosus</i>	Kentucky Warbler	109
<i>Geothlypis philadelphia</i>	Mourning Warbler	108
<i>Geothlypis tolmiei</i>	MacGillivray's Warbler	108
<i>Geothlypis trichas</i>	Common Yellowthroat	109
<i>Glaucidium gnoma</i>	Northern Pygmy-Owl	70
<i>Glaucomys sabrinus</i>	Northern Flying Squirrel	144
<i>Grus americana</i>	Whooping Crane	41
<i>Gulo gulo</i>	Wolverine	157
<i>Gymnorhinus cyanocephalus</i>	Pinyon Jay	85
<i>Haemorhous cassinii</i>	Cassin's Finch	101
<i>Haemorhous mexicanus</i>	House Finch	100
<i>Haemorhous purpureus</i>	Purple Finch	100
<i>Haliaeetus leucocephalus</i>	Bald Eagle	65
<i>Helmitheros vermivorum</i>	Worm-eating Warbler	105
<i>Heterodon nasicus</i>	Plains Hog-nosed Snake	169
<i>Himantopus mexicanus</i>	Black-necked Stilt	41
<i>Hirundo rustica</i>	Barn Swallow	89
<i>Histrionicus histrionicus</i>	Harlequin Duck	25
<i>Holbrookia maculata maculata</i>	Great Plains Earless Lizard	169
<i>Hydrocoloeus minutus</i>	Little Gull	53
<i>Hydroprogne caspia</i>	Caspian Tern	57
<i>Hylocichla mustelina</i>	Wood Thrush	96

Scientific Name	Common Name	Page No.
<i>Icteria virens</i>	Yellow-breasted Chat	115
<i>Icterus bullockii</i>	Bullock's Oriole	128
<i>Icterus galbula</i>	Baltimore Oriole	128
<i>Icterus parisorum</i>	Scott's Oriole	129
<i>Icterus spurius</i>	Orchard Oriole	128
<i>Ictidomys tridecemlineatus</i>	Thirteen-lined Ground Squirrel	142
<i>Ictinia mississippiensis</i>	Mississippi Kite	65
<i>Ixobrychus exilis</i>	Least Bittern	61
<i>Ixoreus naevius</i>	Varied Thrush	97
<i>Junco hyemalis</i>	Dark-eyed Junco	122
<i>Lagopus leucura</i>	White-tailed Ptarmigan	30
<i>Lampropeltis triangulum multistriata</i>	Pale Milksnake	170
<i>Lanius excubitor</i>	Northern Shrike	82
<i>Lanius ludovicianus</i>	Loggerhead Shrike	82
<i>Larus argentatus</i>	Herring Gull	55
<i>Larus californicus</i>	California Gull	54
<i>Larus canus</i>	Mew Gull	54
<i>Larus delawarensis</i>	Ring-billed Gull	54
<i>Larus fuscus</i>	Lesser Black-backed Gull	55
<i>Larus glaucescens</i>	Glaucous-winged Gull	56
<i>Larus glaucoides</i>	Iceland Gull	55
<i>Larus heermanni</i>	Heermann's Gull	54
<i>Larus hyperboreus</i>	Glaucous Gull	56
<i>Larus marinus</i>	Great Black-backed Gull	56
<i>Larus thayeri</i>	Thayer's Gull	55
<i>Lasionycteris noctivagans</i>	Silver-haired Bat	136
<i>Lasiurus borealis</i>	Eastern Red Bat	135
<i>Lasiurus cinereus</i>	Hoary Bat	136
<i>Laterallus jamaicensis</i>	Black Rail	39
<i>Lemmiscus curtatus</i>	Sagebrush Vole	151

Scientific Name	Common Name	Page No.
<i>Lepus americanus</i>	Snowshoe Hare	139
<i>Lepus californicus</i>	Black-tailed Jackrabbit	139
<i>Lepus townsendii</i>	White-tailed Jackrabbit	140
<i>Leucophaeus atricilla</i>	Laughing Gull	53
<i>Leucophaeus pipixcan</i>	Franklin's Gull	53
<i>Leucosticte atrata</i>	Black Rosy-Finch	99
<i>Leucosticte australis</i>	Brown-capped Rosy-Finch	100
<i>Leucosticte tephrocotis</i>	Gray-crowned Rosy-Finch	99
<i>Limnodromus griseus</i>	Short-billed Dowitcher	47
<i>Limnodromus scolopaceus</i>	Long-billed Dowitcher	48
<i>Limosa fedoa</i>	Marbled Godwit	44
<i>Limosa haemastica</i>	Hudsonian Godwit	44
<i>Lithobates catesbeianus</i>	American Bullfrog	164
<i>Lithobates pipiens</i>	Northern Leopard Frog	165
<i>Lithobates sylvaticus</i>	Wood Frog	165
<i>Lontra canadensis</i>	Northern River Otter	158
<i>Lophodytes cucullatus</i>	Hooded Merganser	27
<i>Loxia curvirostra</i>	Red Crossbill	101
<i>Loxia leucoptera</i>	White-winged Crossbill	101
<i>Lynx canadensis</i>	Canada Lynx	158
<i>Lynx rufus</i>	Bobcat	159
<i>Marmota flaviventris</i>	Yellow-bellied Marmot	141
<i>Martes caurina</i>	Pacific Marten	155
<i>Megaceryle alcyon</i>	Belted Kingfisher	72
<i>Megascops asio</i>	Eastern Screech-Owl	69
<i>Megascops kennicottii</i>	Western Screech-Owl	69
<i>Melanerpes carolinus</i>	Red-bellied Woodpecker	73
<i>Melanerpes erythrocephalus</i>	Red-headed Woodpecker	73
<i>Melanerpes formicivorus</i>	Acorn Woodpecker	73
<i>Melanerpes lewis</i>	Lewis's Woodpecker	72

Scientific Name	Common Name	Page No.
<i>Melanitta americana</i>	Black Scoter	26
<i>Melanitta fusca</i>	White-winged Scoter	26
<i>Melanitta perspicillata</i>	Surf Scoter	25
<i>Meleagris gallopavo</i>	Wild Turkey	31
<i>Melospiza georgiana</i>	Swamp Sparrow	121
<i>Melospiza lincolni</i>	Lincoln's Sparrow	120
<i>Melospiza melodia</i>	Song Sparrow	120
<i>Melospiza fusca</i>	Canyon Towhee	116
<i>Mephitis mephitis</i>	Striped Skunk	158
<i>Mergus merganser</i>	Common Merganser	27
<i>Mergus serrator</i>	Red-breasted Merganser	28
<i>Microtus longicaudus</i>	Long-tailed Vole	149
<i>Microtus montanus</i>	Montane Vole	150
<i>Microtus ochrogaster</i>	Prairie Vole	150
<i>Microtus pennsylvanicus</i>	Meadow Vole	150
<i>Microtus richardsoni</i>	Water Vole	150
<i>Mimus polyglottos</i>	Northern Mockingbird	98
<i>Mniotilta varia</i>	Black-and-white Warbler	106
<i>Molothrus ater</i>	Brown-headed Cowbird	128
<i>Mus musculus</i>	House Mouse	151
<i>Mustela erminea</i>	Short-tailed Weasel (Ermine)	155
<i>Mustela frenata</i>	Long-tailed Weasel	156
<i>Mustela nigripes</i>	Black-footed Ferret	156
<i>Mustela nivalis</i>	Least Weasel	156
<i>Myadestes townsendi</i>	Townsend's Solitaire	95
<i>Mycteria americana</i>	Wood Stork	59
<i>Myiarchus cinerascens</i>	Ash-throated Flycatcher	80
<i>Myiarchus crinitus</i>	Great Crested Flycatcher	81
<i>Myodes gapperi</i>	Southern Red-backed Vole	149
<i>Myotis californicus</i>	California Myotis	133

Scientific Name	Common Name	Page No.
<i>Myotis ciliolabrum</i>	Western Small-footed Myotis	134
<i>Myotis evotis</i>	Long-eared Myotis	134
<i>Myotis lucifugus</i>	Little Brown Myotis	134
<i>Myotis septentrionalis</i>	Northern Long-eared Myotis	134
<i>Myotis thysanodes</i>	Fringed Myotis	135
<i>Myotis volans</i>	Long-legged Myotis	135
<i>Myotis yumanensis</i>	Yuma Myotis	135
<i>Neotoma cinerea</i>	Bushy-tailed Woodrat	149
<i>Nucifraga columbiana</i>	Clark's Nutcracker	86
<i>Numenius americanus</i>	Long-billed Curlew	44
<i>Numenius phaeopus</i>	Whimbrel	43
<i>Nyctanassa violacea</i>	Yellow-crowned Night-Heron	63
<i>Nycticorax nycticorax</i>	Black-crowned Night-Heron	63
<i>Nyctinomops macrotis</i>	Big Free-tailed Bat	138
<i>Ochotona princeps</i>	American Pika	138
<i>Odocoileus hemionus</i>	Mule Deer (Black-tailed Deer)	159
<i>Odocoileus virginianus</i>	White-tailed Deer	159
<i>Ondatra zibethicus</i>	Common Muskrat	151
<i>Onychomys leucogaster</i>	Northern Grasshopper Mouse	148
<i>Opheodrys vernalis</i>	Smooth Greensnake	170
<i>Oporornis agilis</i>	Connecticut Warbler	108
<i>Oreamnos americanus</i>	Mountain Goat	160
<i>Oreoscoptes montanus</i>	Sage Thrasher	97
<i>Oreothlypis celata</i>	Orange-crowned Warbler	107
<i>Oreothlypis peregrina</i>	Tennessee Warbler	107
<i>Oreothlypis ruficapilla</i>	Nashville Warbler	107
<i>Oreothlypis virginiae</i>	Virginia's Warbler	108
<i>Ovis canadensis</i>	Bighorn Sheep (Mountain Sheep)	161
<i>Oxyura jamaicensis</i>	Ruddy Duck	28
<i>Pandion haliaetus</i>	Osprey	64

Scientific Name	Common Name	Page No.
<i>Parabuteo unicinctus</i>	Harris's Hawk	66
<i>Parkesia noveboracensis</i>	Northern Waterthrush	106
<i>Passer domesticus</i>	House Sparrow	98
<i>Passerculus sandwichensis</i>	Savannah Sparrow	119
<i>Passerella iliaca</i>	Fox Sparrow	120
<i>Passerina amoena</i>	Lazuli Bunting	124
<i>Passerina caerulea</i>	Blue Grosbeak	124
<i>Passerina ciris</i>	Painted Bunting	125
<i>Passerina cyanea</i>	Indigo Bunting	125
<i>Patagioenas fasciata</i>	Band-tailed Pigeon	33
<i>Pekania pennanti</i>	Fisher	155
<i>Pelecanus erythrorhynchos</i>	American White Pelican	60
<i>Pelecanus occidentalis</i>	Brown Pelican	60
<i>Perdix perdix</i>	Gray Partridge	29
<i>Perimyotis subflavus</i>	American Perimyotis (Tri-colored Bat)	136
<i>Perisoreus canadensis</i>	Gray Jay	85
<i>Perognathus fasciatus</i>	Olive-backed Pocket Mouse	145
<i>Perognathus flavescens</i>	Plains Pocket Mouse	145
<i>Perognathus flavus</i>	Silky Pocket Mouse	146
<i>Perognathus mollipilosus</i>	Great Basin Pocket Mouse	146
<i>Peromyscus crinitus</i>	Canyon Deermouse	147
<i>Peromyscus leucopus</i>	White-footed Deermouse	148
<i>Peromyscus maniculatus</i>	North American Deermouse	148
<i>Peromyscus truei</i>	Piñon Deermouse	148
<i>Petrochelidon pyrrhonota</i>	Cliff Swallow	88
<i>Peucaea cassinii</i>	Cassin's Sparrow	116
<i>Phalacrocorax auritus</i>	Double-crested Cormorant	60
<i>Phalaenoptilus nuttallii</i>	Common Poorwill	36
<i>Phalaropus fulicarius</i>	Red Phalarope	50
<i>Phalaropus lobatus</i>	Red-necked Phalarope	50

Scientific Name	Common Name	Page No.
<i>Phalaropus tricolor</i>	Wilson's Phalarope	50
<i>Phasianus colchicus</i>	Ring-necked Pheasant	29
<i>Phenacomys intermedius</i>	Western Heather Vole	149
<i>Pheucticus chrysopheplus</i>	Yellow Grosbeak	123
<i>Pheucticus ludovicianus</i>	Rose-breasted Grosbeak	124
<i>Pheucticus melanocephalus</i>	Black-headed Grosbeak	124
<i>Phrynosoma hernandesi</i>	Greater Short-horned Lizard	168
<i>Pica hudsonia</i>	Black-billed Magpie	86
<i>Picoides albolarvatus</i>	White-headed Woodpecker	75
<i>Picoides arcticus</i>	Black-backed Woodpecker	75
<i>Picoides dorsalis</i>	American Three-toed Woodpecker	75
<i>Picoides pubescens</i>	Downy Woodpecker	74
<i>Picoides villosus</i>	Hairy Woodpecker	74
<i>Pinicola enucleator</i>	Pine Grosbeak	100
<i>Pipilo chlorurus</i>	Green-tailed Towhee	115
<i>Pipilo maculatus</i>	Spotted Towhee	115
<i>Piranga flava</i>	Hepatic Tanager	122
<i>Piranga ludoviciana</i>	Western Tanager	123
<i>Piranga olivacea</i>	Scarlet Tanager	123
<i>Piranga rubra</i>	Summer Tanager	122
<i>Pituophis catenifer deserticola</i>	Great Basin Gophersnake	171
<i>Pituophis catenifer sayi</i>	Bullsnake	171
<i>Plectrophenax nivalis</i>	Snow Bunting	105
<i>Plegadis chihi</i>	White-faced Ibis	64
<i>Plegadis falcinellus</i>	Glossy Ibis	63
<i>Plestiodon multivirgatus multivirgatus</i>	Northern Many-lined Skink	167
<i>Plestiodon skiltonianus utahensis</i>	Great Basin Skink	167
<i>Pluvialis dominica</i>	American Golden-Plover	42
<i>Pluvialis squatarola</i>	Black-bellied Plover	41
<i>Podiceps auritus</i>	Horned Grebe	32

Scientific Name	Common Name	Page No.
<i>Podiceps grisegena</i>	Red-necked Grebe	32
<i>Podiceps nigricollis</i>	Eared Grebe	32
<i>Podilymbus podiceps</i>	Pied-billed Grebe	31
<i>Poecile atricapillus</i>	Black-capped Chickadee	89
<i>Poecile gambeli</i>	Mountain Chickadee	89
<i>Poliophtila caerulea</i>	Blue-gray Gnatcatcher	93
<i>Pooecetes gramineus</i>	Vesper Sparrow	117
<i>Porphyrio martinicus</i>	Purple Gallinule	40
<i>Porzana carolina</i>	Sora	39
<i>Procyon lotor</i>	Northern Raccoon	155
<i>Progne subis</i>	Purple Martin	87
<i>Protonotaria citrea</i>	Prothonotary Warbler	107
<i>Psaltriparus minimus</i>	Bushtit	90
<i>Pseudacris maculata</i>	Boreal Chorus Frog	165
<i>Psilosops flammeolus</i>	Flammulated Owl	69
<i>Puma concolor</i>	Mountain Lion (Puma)	158
<i>Pyrocephalus rubinus</i>	Vermilion Flycatcher	80
<i>Quiscalus mexicanus</i>	Great-tailed Grackle	127
<i>Quiscalus quiscula</i>	Common Grackle	127
<i>Rallus limicola</i>	Virginia Rail	39
<i>Rana luteiventris</i>	Columbia Spotted Frog	165
<i>Rattus norvegicus</i>	Norway Rat	151
<i>Recurvirostra americana</i>	American Avocet	41
<i>Regulus calendula</i>	Ruby-crowned Kinglet	94
<i>Regulus satrapa</i>	Golden-crowned Kinglet	94
<i>Reithrodontomys megalotis</i>	Western Harvest Mouse	147
<i>Reithrodontomys montanus</i>	Plains Harvest Mouse	147
<i>Rhodostethia rosea</i>	Ross's Gull	53
<i>Rhynchophanes mccownii</i>	McCown's Longspur	105
<i>Riparia riparia</i>	Bank Swallow	88

Scientific Name	Common Name	Page No.
<i>Rissa tridactyla</i>	Black-legged Kittiwake	52
<i>Salpinctes obsoletus</i>	Rock Wren	91
<i>Sayornis phoebe</i>	Eastern Phoebe	80
<i>Sayornis saya</i>	Say's Phoebe	80
<i>Scalopus aquaticus</i>	Eastern Mole	133
<i>Sceloporus consobrinus</i>	Prairie Lizard	168
<i>Sceloporus graciosus graciosus</i>	Northern Sagebrush Lizard	167
<i>Sceloporus tristichus</i>	Plateau Fence Lizard	168
<i>Sciurus aberti</i>	Abert's Squirrel	143
<i>Sciurus carolinensis</i>	Eastern Gray Squirrel	143
<i>Sciurus niger</i>	Eastern Fox Squirrel	143
<i>Scolopax minor</i>	American Woodcock	48
<i>Seiurus aurocapilla</i>	Ovenbird	105
<i>Selasphorus calliope</i>	Calliope Hummingbird	38
<i>Selasphorus platycercus</i>	Broad-tailed Hummingbird	38
<i>Selasphorus rufus</i>	Rufous Hummingbird	38
<i>Setophaga americana</i>	Northern Parula	110
<i>Setophaga caerulescens</i>	Black-throated Blue Warbler	112
<i>Setophaga castanea</i>	Bay-breasted Warbler	110
<i>Setophaga citrina</i>	Hooded Warbler	109
<i>Setophaga coronata</i>	Yellow-rumped Warbler	112
<i>Setophaga discolor</i>	Prairie Warbler	113
<i>Setophaga dominica</i>	Yellow-throated Warbler	113
<i>Setophaga fusca</i>	Blackburnian Warbler	111
<i>Setophaga magnolia</i>	Magnolia Warbler	110
<i>Setophaga nigrescens</i>	Black-throated Gray Warbler	113
<i>Setophaga occidentalis</i>	Hermit Warbler	114
<i>Setophaga palmarum</i>	Palm Warbler	112
<i>Setophaga pensylvanica</i>	Chestnut-sided Warbler	111
<i>Setophaga petechia</i>	Yellow Warbler	111

Scientific Name	Common Name	Page No.
<i>Setophaga pinus</i>	Pine Warbler	112
<i>Setophaga ruticilla</i>	American Redstart	109
<i>Setophaga striata</i>	Blackpoll Warbler	111
<i>Setophaga tigrina</i>	Cape May Warbler	110
<i>Setophaga townsendi</i>	Townsend's Warbler	113
<i>Setophaga virens</i>	Black-throated Green Warbler	114
<i>Sialia currucoides</i>	Mountain Bluebird	95
<i>Sialia mexicana</i>	Western Bluebird	94
<i>Sialia sialis</i>	Eastern Bluebird	94
<i>Sitta canadensis</i>	Red-breasted Nuthatch	90
<i>Sitta carolinensis</i>	White-breasted Nuthatch	90
<i>Sitta pygmaea</i>	Pygmy Nuthatch	90
<i>Sorex cinereus</i>	Masked Shrew	131
<i>Sorex haydeni</i>	Hayden's Shrew	131
<i>Sorex hoyi</i>	American Pygmy Shrew	131
<i>Sorex merriami</i>	Merriam's Shrew	132
<i>Sorex monticolus</i>	Dusky Shrew	132
<i>Sorex nanus</i>	Dwarf Shrew	132
<i>Sorex navigator</i>	Western Water Shrew	132
<i>Sorex preblei</i>	Preble's Shrew	133
<i>Sorex vagrans</i>	Vagrant Shrew	133
<i>Spea bombifrons</i>	Plains Spadefoot	163
<i>Spea intermontana</i>	Great Basin Spadefoot	163
<i>Sphyrapicus nuchalis</i>	Red-naped Sapsucker	74
<i>Sphyrapicus thyroideus</i>	Williamson's Sapsucker	73
<i>Sphyrapicus varius</i>	Yellow-bellied Sapsucker	74
<i>Spilogale gracilis</i>	Western Spotted Skunk	157
<i>Spilogale putorius</i>	Eastern Spotted Skunk	157
<i>Spinus lawrencei</i>	Lawrence's Goldfinch	102
<i>Spinus pinus</i>	Pine Siskin	102

Scientific Name	Common Name	Page No.
<i>Spinus psaltria</i>	Lesser Goldfinch	102
<i>Spinus tristis</i>	American Goldfinch	103
<i>Spiza americana</i>	Dickcissel	125
<i>Spizella breweri</i>	Brewer's Sparrow	117
<i>Spizella pallida</i>	Clay-colored Sparrow	117
<i>Spizella passerina</i>	Chipping Sparrow	116
<i>Spizella pusilla</i>	Field Sparrow	117
<i>Spizelloides arborea</i>	American Tree Sparrow	116
<i>Stelgidopteryx serripennis</i>	Northern Rough-winged Swallow	88
<i>Stercorarius longicaudus</i>	Long-tailed Jaeger	51
<i>Stercorarius parasiticus</i>	Parasitic Jaeger	51
<i>Stercorarius pomarinus</i>	Pomarine Jaeger	50
<i>Sterna forsteri</i>	Forster's Tern	58
<i>Sterna hirundo</i>	Common Tern	57
<i>Sterna paradisaea</i>	Arctic Tern	57
<i>Sternula antillarum</i>	Least Tern	56
<i>Storeria occipitomaculata pahasapae</i>	Black Hills Red-bellied Snake	170
<i>Streptopelia decaocto</i>	Eurasian Collared-Dove	34
<i>Streptopelia roseogrisea</i>	African Collared-Dove	33
<i>Strix nebulosa</i>	Great Gray Owl	71
<i>Strix varia</i>	Barred Owl	71
<i>Sturnella magna</i>	Eastern Meadowlark	126
<i>Sturnella neglecta</i>	Western Meadowlark	126
<i>Sturnus vulgaris</i>	European Starling	98
<i>Surnia ulula</i>	Northern Hawk Owl	70
<i>Sylvilagus audubonii</i>	Desert Cottontail	138
<i>Sylvilagus floridanus</i>	Eastern Cottontail	139
<i>Sylvilagus nutallii</i>	Mountain Cottontail	139
<i>Synthliboramphus antiquus</i>	Ancient Murrelet	51
<i>Tachycineta bicolor</i>	Tree Swallow	87

Scientific Name	Common Name	Page No.
<i>Tachycineta thalassina</i>	Violet-green Swallow	88
<i>Tadarida brasiliensis</i>	Brazilian Free-tailed Bat	137
<i>Tadorna ferruginea</i>	Ruddy Shelduck	28
<i>Tadorna tadorna</i>	Common Shelduck	28
<i>Tamias amoenus</i>	Yellow-pine Chipmunk	140
<i>Tamias dorsalis</i>	Cliff Chipmunk	140
<i>Tamias minimus</i>	Least Chipmunk	140
<i>Tamias umbrinus</i>	Uinta Chipmunk	141
<i>Tamiasciurus hudsonicus</i>	Red Squirrel	144
<i>Tantilla nigriceps</i>	Plains Black-headed Snake	171
<i>Taxidea taxus</i>	American Badger	157
<i>Terrapene ornata ornata</i>	Plains Box Turtle	166
<i>Thamnophis elegans vagrans</i>	Wandering Gartersnake	171
<i>Thamnophis radix</i>	Plains Gartersnake	172
<i>Thamnophis sirtalis fitchi</i>	Valley Gartersnake	172
<i>Thamnophis sirtalis parietalis</i>	Red-sided Gartersnake	172
<i>Thomomys clusius</i>	Wyoming Pocket Gopher	144
<i>Thomomys idahoensis</i>	Idaho Pocket Gopher	144
<i>Thomomys talpoides</i>	Northern Pocket Gopher	145
<i>Thryomanes bewickii</i>	Bewick's Wren	93
<i>Thryothorus ludovicianus</i>	Carolina Wren	93
<i>Toxostoma rufum</i>	Brown Thrasher	97
<i>Tringa flavipes</i>	Lesser Yellowlegs	49
<i>Tringa melanoleuca</i>	Greater Yellowlegs	49
<i>Tringa semipalmata</i>	Willet	49
<i>Tringa solitaria</i>	Solitary Sandpiper	49
<i>Troglodytes aedon</i>	House Wren	91
<i>Troglodytes hiemalis</i>	Winter Wren	92
<i>Troglodytes pacificus</i>	Pacific Wren	92
<i>Turdus migratorius</i>	American Robin	96

Scientific Name	Common Name	Page No.
<i>Tympanuchus cupido</i>	Greater Prairie-Chicken	31
<i>Tympanuchus phasianellus</i>	Sharp-tailed Grouse	31
<i>Tyrannus forficatus</i>	Scissor-tailed Flycatcher	82
<i>Tyrannus tyrannus</i>	Eastern Kingbird	81
<i>Tyrannus verticalis</i>	Western Kingbird	81
<i>Tyrannus vociferans</i>	Cassin's Kingbird	81
<i>Tyto alba</i>	Barn Owl	68
<i>Urocitellus elegans</i>	Wyoming Ground Squirrel	141
<i>Urocitellus armatus</i>	Uinta Ground Squirrel	141
<i>Urocyon cinereoargenteus</i>	Common Gray Fox	154
<i>Urosaurus ornatus wrighti</i>	Northern Tree Lizard	168
<i>Ursus americanus</i>	Black Bear	154
<i>Ursus arctos</i>	Grizzly Bear	154
<i>Vermivora chrysoptera</i>	Golden-winged Warbler	106
<i>Vermivora cyanoptera</i>	Blue-winged Warbler	106
<i>Vireo cassinii</i>	Cassin's Vireo	83
<i>Vireo flavifrons</i>	Yellow-throated Vireo	83
<i>Vireo gilvus</i>	Warbling Vireo	84
<i>Vireo griseus</i>	White-eyed Vireo	82
<i>Vireo olivaceus</i>	Red-eyed Vireo	84
<i>Vireo philadelphicus</i>	Philadelphia Vireo	84
<i>Vireo plumbeus</i>	Plumbeous Vireo	84
<i>Vireo solitarius</i>	Blue-headed Vireo	83
<i>Vireo vicinior</i>	Gray Vireo	83
<i>Vison vison</i>	American Mink	156
<i>Vulpes velox</i>	Swift Fox	153
<i>Vulpes vulpes</i>	Red Fox	153
<i>Xanthocephalus xanthocephalus</i>	Yellow-headed Blackbird	126
<i>Xema sabini</i>	Sabine's Gull	52
<i>Xerospermophilus spilosoma</i>	Spotted Ground Squirrel	142

Scientific Name	Common Name	Page No.
<i>Zapus hudsonius</i>	Meadow Jumping Mouse	152
<i>Zapus hudsonius preblei</i>	Preble's Meadow Jumping Mouse	152
<i>Zapus princeps</i>	Western Jumping Mouse	152
<i>Zenaida asiatica</i>	White-winged Dove	34
<i>Zenaida macroura</i>	Mourning Dove	34
<i>Zonotrichia albicollis</i>	White-throated Sparrow	121
<i>Zonotrichia atricapilla</i>	Golden-crowned Sparrow	122
<i>Zonotrichia leucophrys</i>	White-crowned Sparrow	121
<i>Zonotrichia querula</i>	Harris's Sparrow	121

Appendix I

Rare and Unusual Bird Sighting Form
WYOMING BIRD RECORDS COMMITTEE
260 BUENA VISTA, LANDER, WY 82520

The Wyoming Bird Records Committee sincerely thanks all observers for submitting this form to help keep Wyoming's bird records up-to-date and accurate. It is not necessary to complete every block if some details are lacking, but please provide all the details you can and attach photographs, if available.

Common Name:	Specific location of observation:
Scientific Name:	
Observation Date:	
Observation Time:	
Length of Observation:	UTM E _____ UTM N _____ Datum _____ Zone _____ T
Distance from Bird:	Latitude _____° _____' _____" N Longitude _____° _____' _____" W
Light Conditions:	Weather at time of observation:
Optical Equipment:	
Notes made: during sighting _____ from memory _____	Prior weather and number of days since last change:
Date report prepared	

Give a general description of the bird seen and any other details of interest relating to the observation.

SEX	AGE	PLUMAGE		PHOTO/AUDIO/DRAWING
Male:	Adult:	Breeding:	Juvenal:	Enclosed:
Female:	Juvenile/Immature:	Winter:	Dark Morph:	Available:
Unknown:	2-3 year bird:	Eclipse:	Light Morph:	Please submit a copy of your field drawings.
Total Number:	Unknown:	Other:		

If possible, please include in the sections below details of the specific body parts actually observed during the sighting.

BILL:
HEAD:
NECK:
UPPERPARTS:
UNDERPARTS:

Please do not write below here; for WBRC use only

Form updated March 2016

Record Number	Category	Latilong	Date Received
----------------------	-----------------	-----------------	----------------------

WINGS:

TAIL:

LEGS & FEET:

List similar species and describe how or why you eliminated them.

Describe the behavior of this bird and the interaction with others.

What is the habitat at this location?

If heard, describe the bird's song or vocalizations.

Reporter's name, address, phone number, and e-mail address.

How many years have you birded?

Corroborating observers who are not reporting separately.

Have you observed this species before?

Please do not write below here; for WBRC use only

Form updated March 2016

Record Number

Category

Latilong

Date Received

THANK YOU FOR SUBMITTING YOUR SIGHTING TO THE WYOMING BIRD RECORDS COMMITTEE!

Rare and Unusual Bird Sighting Form – Photographs Only
WYOMING BIRD RECORDS COMMITTEE
260 BUENA VISTA, LANDER, WY 82520

The Wyoming Bird Records Committee sincerely thanks all observers for submitting this form and your photographs to help keep Wyoming's bird records up-to-date and accurate.

Common Name:	Specific location of observation:			
Scientific Name:				
Observation Date:				
Legal Description:				
UTM E	UTM N	Datum	Zone	T
Observer (name and contact information):				

Please do not write below here; for WBRC use only

Form updated March 2016

Record Number:	Category:	Latilong:	Date Received:
Atlas Update:		Sighting Entered in WGFD WOS Database:	

HOW TO DOCUMENT SIGHTINGS OF RARE AND UNUSUAL BIRDS

To improve the accuracy and breadth of Wyoming's ornithological record, the Wyoming Bird Records Committee (WBRC) offers the following recommendations to assist with documentation of bird sightings.

- 1) Read the excellent article "How to Document Rare Birds", by Donna L. Dittman and Greg W. Lasley (pages 145-149 in the 1992 issue of *Birding*, Volume 2, Number 3; http://www.losbird.org/dittman_lasley.htm).
- 2) Acceptable documentation must eliminate all similar species. Remember that immatures or juveniles of one species can be very similar to adults of another species. Examples that might cause confusion are gulls, jaegers, sparrows, and longspurs. Species that exhibit multiple color morphs can also be problematic.
- 3) Study and learn bird topography. Most field guides provide a schematic of avian body parts and feather groups. Specialized identification guides also provide specific structural and anatomical detail. A thorough grasp of this subject will heighten your general birding skill and facilitate accurate, detailed documentation.
- 4) Take meticulous and thorough field notes during or immediately after the observation. Alternatively, you can also use an audio recorder or digital camera to capture identification details. If the bird is cooperative, write your notes during the observation period. Try not to consult your field guide while you are writing to avoid predisposing your identification. Do not rely on memory to document a rare or unusual bird.
- 5) A good physical description of the bird is most crucial and you should include everything you observe. Include all details concerning plumage, shape, relative size, eyes, legs, and bill. Note the colors, including color distribution, color density, and color contrast between different feather groups. It is helpful to consider proportional details, i.e., bill length compared to head width, tail length as a proportion of body length, and length of primary projection compared to tail length. Record plumage characteristics such as degree of wear or signs of molt. When describing size, try to compare nearby known species or some other object of known dimensions. Avoid trying to estimate size in actual units (inches, feet) since this is very subjective.
- 6) Observe and record the subject bird's behavior. While behavior is seldom diagnostic by itself, in combination with other details, it is often conclusive. The observations of the Connecticut Warbler in Wyoming were accepted by the WBRC, in part, because the subjects were observed walking, not hopping.
- 7) If possible, take photographs. Take a number of shots to capture a complete portrait. The advent of digital photography has greatly increased the number of records the WBRC receives with photographs, which greatly helps our work. However, a photograph should be accompanied by a written description, if possible, as one photograph will likely not display all the diagnostic features.
- 8) If you do not have a camera, make a sketch. You do not need the talent of Rembrandt or Audubon to draw a convincing and diagnostic sketch. Utilize your knowledge of bird topography and you will be surprised how well you do. Draw this sketch in the field during the observation. Do not rely on memory.
- 9) Vocalizations can be extremely important for identification, especially for some groups such as flycatchers. Describe what you hear while you are listening. Better yet, record the bird, even if you do not have professional recording equipment. A hand-held recorder or cellular phone with a built in microphone or a digital camera with a video recording feature can provide reliable results. There are many excellent collections of bird songs and calls that can prepare birders for the unexpected find by ear alone.
- 10) Specimens that can be identified and repositied at the University of Wyoming Zoological Museum are still the most convincing evidence of an occurrence. If you encounter a dead rare or unusual bird in the field, please report or deliver the specimen to an appropriate authority, e.g. Wyoming Game and Fish Department, National Park Service, or U.S. Fish and Wildlife Service personnel. Note the exact date and location of the discovery. Freeze the specimen if delivery is to be delayed. Always use gloves or a plastic bag to pick up dead birds, and double-bag them with a note inside that includes the date and location and your contact information.

For questions about the Wyoming Bird Records Committee or clarification of these recommendations, please contact the Wyoming Game and Fish Department Nongame Bird Biologist at the Lander Regional Office, 260 Buena Vista, Lander, WY 82520; 307-332-2688; Andrea.Orabona@wyo.gov.

MAMMAL OBSERVATION RECORD

PLEASE RETURN TO: Nongame Mammal Biologist
Wyoming Game and Fish Department
260 Buena Vista
Lander, Wyoming 82520

Mammal Observed: _____
Number Observed: _____ Photograph Taken: _____
Observer's Name: _____ Telephone: _____
Address: _____
Occupation: _____
Name of Other Observers: _____
Address of Other Observers: _____
Agency/Organization: _____
Reporting Date: _____
Location (direction and distance from the nearest town or identifiable locations, and legal description of the site):

UTM Coordinates: _____ E _____ N Zone _____ Datum (e.g. NAD83) _____
Latitude/Longitude Coordinates: _____
_____ 1/4 _____ 1/4 Section _____ Township _____ Range _____
Describe Site (details about land use, habitat type, etc.): _____

COLLECTION OF DEAD ANIMAL:
Currently Held Where? _____
How Was Specimen Acquired? _____

OBSERVATION OF LIVE ANIMAL:
Time and Duration of Sighting: _____
Weather Conditions: _____
Distance of Observation: _____
Binoculars Used? _____ Spotting Scope Used? _____
Description of Animal(s) (color, size, markings):
Body: _____
Legs and Feet: _____
Tail: _____
Head and Face: _____
Other Comments: _____

Behavior (describe in observer's words): _____

Similar Species and How Observer Eliminated Them: _____

PLEASE DO NOT WRITE IN THIS SPACE:
Classification: _____
Confirmed: _____ Probable: _____ Unconfirmed: _____

Appendix II

DOCUMENTATION REQUESTED FOR ALL SIGHTINGS (AS) OF THE FOLLOWING SPECIES

Acorn Woodpecker	Gray Vireo	Purple Finch
American Black Duck	Gray-cheeked Thrush	Purple Gallinule
American Woodcock	Great Black-backed Gull	Purple Martin
Ancient Murrelet	Great Crested Flycatcher	Red Knot
Anna's Hummingbird	Greater Prairie-Chicken	Red Phalarope
Arctic Tern	Green Heron	Red-bellied Woodpecker
Baird's Sparrow	Gyr Falcon	Red-faced Warbler
Baltimore Oriole	Harris's Hawk	Red-necked Grebe
Band-tailed Pigeon	Heermann's Gull	Red-shouldered Hawk
Barn Owl	Hepatic Tanager	Red-throated Loon
Barred Owl	Hermit Warbler	Ross's Gull
Bay-breasted Warbler	Hoary Redpoll	Ruby-throated Hummingbird
Black Rail	Hooded Warbler	Rusty Blackbird
Black Scoter	Hudsonian Godwit	Scarlet Tanager
Black Swift	Iceland Gull	Scissor-tailed Flycatcher
Black Vulture	Kentucky Warbler	Scott's Oriole
Blackburnian Warbler	Laughing Gull	Sedge Wren
Black-headed Gull	Lawrence's Goldfinch	Short-billed Dowitcher
Black-legged Kittiwake	Le Conte's Sparrow	Smith's Longspur
Black-throated Green Warbler	Least Bittern	Snowy Owl
Black-throated Sparrow	Least Tern	Snowy Plover
Blue-headed Vireo	Lesser Black-backed Gull ²	Sprague's Pipit
Blue-winged Warbler	Lesser Frigatebird	Streaked Shearwater
Brambling	Lesser Nighthawk	Thayer's Gull
Brant	Little Blue Heron	Tricolored Heron
Brown Pelican	Little Gull	Tufted Duck
Buff-breasted Sandpiper	Long-billed Murrelet	Varied Thrush
Canada Warbler	Long-tailed Jaeger	Vaux's Swift
Canyon Towhee	Magnificent Hummingbird	Vermilion Flycatcher
Cape May Warbler	Mew Gull	Western Bluebird
Carolina Wren	Mississippi Kite	Western Screech-Owl
Cassin's Sparrow ¹	Mottled Duck	White Ibis
Cassin's Vireo	Mourning Warbler	White-eyed Vireo
Common Gallinule	Mute Swan	White-headed Woodpecker
Connecticut Warbler	Nelson's Sparrow	White-tailed Kite
Crested Caracara	Northern Bobwhite	White-tailed Ptarmigan
Eastern Meadowlark	Northern Cardinal	Whooper Swan
Eastern Phoebe	Northern Hawk Owl	Whooping Crane
Eastern Wood-Pewee	Pacific Wren	Winter Wren
Eurasian Wigeon	Painted Bunting	Wood Stork
Field Sparrow	Palm Warbler	Wood Thrush
Flammulated Owl	Parasitic Jaeger	Worm-eating Warbler
Fulvous Whistling-Duck	Philadelphia Vireo	Yellow Grosbeak
Garganey	Pileated Woodpecker	Yellow Rail
Glaucous Gull	Pine Warbler	Yellow-bellied Sapsucker
Glaucous-winged Gull	Piping Plover	Yellow-billed Loon
Glossy Ibis	Pomarine Jaeger	Yellow-crowned Night-Heron
Golden-crowned Sparrow	Prairie Warbler	Yellow-throated Vireo
Golden-winged Warbler	Prothonotary Warbler	Yellow-throated Warbler

¹ Except confirmed breeding in Torrington area ² Except L19 & L27

**DOCUMENTATION REQUESTED FOR THE FIRST LATILONG SIGHTINGS (FL)
AND ALL NESTING OBSERVATIONS OF THE FOLLOWING SPECIES**

American Bittern	Hammond's Flycatcher
American Golden-Plover	Indigo Bunting
Ash-throated Flycatcher	Juniper Titmouse
Bewick's Wren	Least Flycatcher
Black-and-white Warbler	Lesser Goldfinch
Black-backed Woodpecker	Long-tailed Duck
Black-billed Cuckoo	Magnolia Warbler
Black-chinned Hummingbird	Nashville Warbler
Blackpoll Warbler	Northern Parula
Black-throated Blue Warbler	Northern Pygmy-Owl
Black-throated Gray Warbler	Northern Saw-whet Owl
Bobolink	Orchard Oriole
Boreal Owl	Pacific Loon
Broad-winged Hawk	Peregrine Falcon
Brown-capped Rosy-Finch	Red-headed Woodpecker
Bushtit	Rose-breasted Grosbeak
Cackling Goose	Ross's Goose
Cassin's Kingbird	Ruddy Turnstone
Cattle Egret	Sabine's Gull
Chestnut-collared Longspur	Summer Tanager
Chestnut-sided Warbler	Surf Scoter
Chimney Swift	Swamp Sparrow
Common Tern	Tennessee Warbler
Dickcissel	Trumpeter Swan
Dunlin	Upland Sandpiper
Eastern Bluebird	Virginia's Warbler
Eastern Screech-Owl	Western Scrub-Jay
Gray Flycatcher	Whimbrel
Great Egret	White-rumped Sandpiper
Greater Scaup	White-winged Crossbill
Greater White-fronted Goose	White-winged Dove
Great-tailed Grackle	White-winged Scoter
Great-tailed Grackle	Yellow-billed Cuckoo

Appendix III

HABITAT CODES AND DESCRIPTIONS

Habitat Code	Habitat Type
01.00	Conifer Forests
01.10	Lodgepole Pine
01.20	Douglas Fir
01.30	Engelmann Spruce-Subalpine Fir
01.40	Ponderosa Pine (savannah)
01.50	Ponderosa Pine-Douglas Fir
01.60	Limber Pine
01.70	Whitebark Pine
01.80	Pine-Juniper
01.81	Pinyon Pine-Juniper
01.82	Limber Pine-Juniper
01.83	Ponderosa Pine-Juniper
01.90	Other or Mixed Conifer Forests
02.00	Deciduous Forests
02.10	Aspen
02.11	Aspen-Upland
02.12	Aspen-Riparian
02.13	Aspen-Conifer
02.20	Riparian-Cottonwood
02.21	Cottonwood-Upland (dry site)
02.22	Cottonwood-Riparian
02.23	Balsam Poplar
02.30	Cottonwood-Dryland
02.31	Plains Cottonwood
02.32	Narrowleaf Cottonwood
02.40	Maple
02.41	Boxelder Maple
02.42	Rocky Mountain Maple
02.43	Bigtooth Maple
02.50	Oak
02.60	Paper Birch
02.70	Maple-Elm-Ash
02.80	Other or Mixed Deciduous Forests
03.00	Woodland-Chaparral
03.10	Gambel Oak

Habitat Code	Habitat Type
03.20	Juniper
03.21	Juniper-Sagebrush
03.22	Juniper-Mountain Mahogany
03.23	Juniper-Other Shrub Understory
04.00	Basin-Prairie Shrub-Shrub Steppe
04.10	Sagebrush-Grassland
04.11	Basin Big Sagebrush
04.12	Wyoming Big Sagebrush
04.13	Mountain Big Sagebrush
04.14	Black Sagebrush
04.15	Silver Sagebrush
04.16	Threetip Sagebrush
04.17	Low Sagebrush (<i>Artemisia arbuscula</i>)
04.18	Sand Sagebrush (<i>Artemisia filifolia</i>)
04.19	Other or Mixed Sagebrush or Sagewort
04.20	Greasewood
04.21	Greasewood-Sagebrush
04.22	Greasewood-Saltbush
04.30	Rabbitbrush
04.31	Green Rabbitbrush
04.32	Rubber Rabbitbrush
04.33	Rabbitbrush-Sagebrush-Mixed Shrub
04.40	Saltbush
04.41	Gardner Saltbush
04.42	Fourwing Saltbush
04.43	Shadscale
04.44	Saltbush-Sagebrush-Mixed Shrub
04.50	Winterfat
04.60	Woody Aster (alkali aster) <i>Xylorhiza</i>
04.70	Other or Mixed Shrub-Shrub Steppe
05.00	Mountain-Foothills Shrub-Shrub Steppe
05.10	Sagebrush-Grassland
05.11	Basin Big Sagebrush
05.12	Wyoming Big Sagebrush
05.13	Mountain Big Sagebrush
05.14	Black Sagebrush
05.15	Silver Sagebrush

Habitat Code	Habitat Type
05.16	Threetip Sagebrush
05.17	Low Sagebrush (<i>Artemisia arbuscula</i>)
05.18	Sand Sagebrush (<i>Artemisia filifolia</i>)
05.19	Other or Mixed Sagebrush
05.20	Rabbitbrush
05.21	Green Rabbitbrush
05.22	Rubber Rabbitbrush
05.23	Rabbitbrush-Sagebrush-Mixed Shrub
05.30	Mountain Mahogany
05.31	True Mountain Mahogany
05.32	True Mountain Mahogany-Sagebrush-Mixed Shrub
05.33	Curleaf Mountain Mahogany
05.34	Curleaf Mountain Mahogany-Sagebrush-Mixed Shrub
05.40	Bitterbrush
05.41	Bitterbrush-Sagebrush-Mixed Shrub
05.50	Serviceberry
05.51	Serviceberry-Sagebrush-Mixed Shrub
05.60	Chokecherry
05.61	Chokecherry-Sagebrush-Mixed Shrub
05.70	Hawthorne
05.80	Skunkbush Sumac
05.90	Other or Mixed Shrub-Shrub Steppe
06.00	Riparian Shrub-Shrub Steppe
06.10	Willow
06.11	Alpine Willow
06.12	Willow-Other Shrubs
06.20	Hawthorne-Wild Plum-Dogwood
06.30	Water Birch-Bog Birch
06.31	Birch-Alder
06.32	Birch-Willow
06.33	Birch-Other Shrubs
06.40	Alder
06.50	Shrubby Cinquefoil
06.60	Buffaloberry
06.70	Tamarisk
06.80	Russian Olive
06.90	Other or Mixed Shrub-Shrub Steppe

Habitat Code	Habitat Type
07.00	Grasslands
07.10	Eastern Great Plains Area Grasslands
07.11	Shortgrass (blue grama-buffalo grass)
07.12	Midgrass (blue grama-needlegrass-Western wheatgrass)
07.13	Draws (bluebunch wheatgrass-needlegrass-wildrye)
07.14	Sandy (muhly-sand dropseed-prairie sandreed)
07.15	Tallgrass, Rough Breaks (little bluestem-sideoats grama-switchgrass punice)
07.16	Tallgrass (big, sand, and little bluestem-prairie sandreed-needleandthread)
07.17	Annual Forb
07.20	Great Basin-Foothills Grasslands
07.21	Bluebunch Wheatgrass-Bluegrass
07.22	Thickspike-Western Wheatgrass-Needleandthread-Bluegrass-Blue Grama
07.23	Annual Forb
07.30	Mountain-Foothills Grassland
07.31	Bluebunch Wheatgrass-Idaho Fescue-Bluegrass-Needlegrass
07.32	Mountain Park-Meadow (bent grasses, Idaho fescue-reed grasses-slender wheatgrass)
07.33	Foothills-Mountain Forb (Northwest cinquefoil-lupine-geranium-elkweed)
07.34	Annual Forb
07.40	Alpine Grasslands (bent grass-sheep fescue-icegrass)
07.50	Alpine Moss-Lichen-Forb (silene-avens-sundivort-forget-me-not)
07.60	Wet-Moist Meadow Grasslands
07.61	Tufted Hairgrass-Reedgrass (freshwater)
07.62	Inland Saltgrass-Alkali Sacaton (alkaline)
07.63	Common Reed (phragmites)
07.64	Grass-Sedge-Rush Meadow
07.70	Kentucky Bluegrass Grasslands
07.80	Annual Grasslands
08.00	Grasslike Types
08.10	Sedge
08.20	Sedge-Grass-Rush Meadow
09.00	Marsh-Swamp Wetlands
09.10	Cattail
09.20	Bulrush
09.30	Cattail-Bulrush-Sedge
10.00	Aquatic
10.10	Lentic or Standing Water (pond, lake, reservoir)

Habitat Code	Habitat Type
10.11	Large Lentic (>5 surface acres)
10.12	Small Lentic (<5 surface acres)
10.13	Intermittent Lentic
10.20	Lotic or Running Water (spring, stream, river)
10.21	Large Lotic (>50 ft. channel width)
10.22	Small Lotic (<50 ft. channel width)
10.23	Spring
10.24	Intermittent
11.00	Cropland-Agricultural Lands
11.10	Row Crop
11.11	Corn
11.12	Beans
11.13	Potatoes
11.14	Beets
11.15	Sunflower
11.16	Other Row Crop
11.20	Alfalfa
11.21	Standing (in field)
11.22	Stacked
11.23	Seed Crop
11.30	Irrigated Native Meadow (hayed or pastured)
11.40	Irrigated Introduced Meadow (hayed or pastured)
11.50	Small Grains
11.51	Wheat
11.52	Barley
11.53	Oats
11.54	Rye
11.55	Other Small Grains
11.60	Dryland Grass Pastures
11.61	Crested Wheatgrass
11.62	Russian Wildrye
11.63	Other-Mixed Dryland Grass Pastures
11.70	Fallow Lands
11.80	Stored Crops
11.90	Rural Development
11.91	Ranch-Farm Facilities
11.92	Shelterbelt

Habitat Code	Habitat Type
12.00	Barren-Special Features
12.10	Cave
12.20	Cliffs (canyon, mountain)
12.30	Spires
12.40	Rock Outcrop-Rock Piles
12.50	Talus Slope-Talus Fields
12.60	Sand Dune-Sand Blowouts
12.70	Shoreline (mud flats, sand beach, gravel bar)
12.80	Glacier
99.00	Disturbed Areas
99.10	Roadside-Railroad Banks
99.20	Mined Area (unreclaimed)
99.30	Mined Area (reclaimed)
99.31	Grass (native species)
99.32	Grass (introduced species)
99.33	Shrub-Shrub-grass
99.34	Deciduous Tree
99.35	Coniferous Tree
99.40	Sprayed Area
99.41	Sagebrush
99.42	Other Shrub
99.43	Other
99.50	Burned Area
99.51	Grass
99.52	Sagebrush
99.53	Other Shrub
99.54	Woodland or Chaparral
99.55	Deciduous Tree
99.56	Coniferous Tree
99.57	Marsh
99.60	Logged-Clearcut Area
99.61	Lodgepole Pine
99.62	Ponderosa Pine
99.63	Douglas Fir
99.64	Spruce-Fir
99.65	Other or Mixed Conifer
99.66	Deciduous

Habitat Code	Habitat Type
99.70	Logged-Thinned
99.71	Lodgepole Pine
99.72	Ponderosa Pine
99.73	Douglas Fir
99.74	Spruce-Fir
99.75	Other or Mixed Conifer
99.76	Deciduous
99.80	Oil and Gas Sites
99.90	Urban-Built Up
99.91	Garbage Dump
99.92	Other